

ASPIRIN LOW DOSE- aspirin tablet, chewable

Chain Drug Consortium

Premier Value 44-218

Active ingredient (in each chewable tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- have had stomach ulcers or bleeding problems

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - feel faint
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 chewable tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

- see end flap for expiration date and lot number

Inactive ingredients

corn starch, dextrans hydrated, FD&C yellow #6 aluminum lake, flavor, maltodextrin, sucralose

Questions or comments?

1-800-426-9391

Principal Display Panel

VALUE PACK

108 Tablets

(3 bottles of 36 tablets each)

†COMPARE TO THE ACTIVE INGREDIENT IN
BAYER® CHEWABLE LOW DOSE ASPIRIN

Premier Value®

Low Dose

Aspirin 81 mg

Chewable Tablets

PAIN RELIEVER (NSAID)

Aspirin regimen**

actual size

108 Orange flavored

Chewable Tablets

**Chew or crush tablets completely
before swallowing.**

****Talk to your doctor or other healthcare provider
before using this product for your heart.**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY
SEAL UNDER CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer®

Chewable Low Dose Aspirin. 50844 REV0122B21807

Distributed By:

Pharmacy Value Alliance, LLC

407 East Lancaster Avenue

Wayne, PA 19087

**If for any reason you are
not satisfied with this product,**

**please return it to the store
where purchased for a full refund.**

Drug Facts

Warnings Read on and be aware of the risks of aspirin. Aspirin should not be used in children under 12 years of age because of the risk of Reye's syndrome, a rare but serious illness. Aspirin should not be used in children under 12 years of age because of the risk of Reye's syndrome, a rare but serious illness. Aspirin should not be used in children under 12 years of age because of the risk of Reye's syndrome, a rare but serious illness.

Uses For the temporary relief of minor aches and pains or as recommended by your doctor.

Active ingredient (in each chewable tablet) Aspirin 81 mg (NSAID) (nonsteroidal anti-inflammatory drug)

Purpose Pain reliever

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

VALUE PACK
108 Tablets
(3 bottles of 36 tablets each)



Low Dose Aspirin 81 mg Chewable Tablets
PAIN RELIEVER (NSAID)

Aspirin regimen**

108 Orange flavored Chewable Tablets
Chew or crush tablets completely before swallowing.

COMPARE TO THE ACTIVE INGREDIENT IN BAYER® CHEWABLE LOW DOSE ASPIRIN

Low Dose Aspirin 81 mg Chewable Tablets
PAIN RELIEVER (NSAID)

108 Orange flavored Chewable Tablets
Chew or crush tablets completely before swallowing.

108 Orange flavored Chewable Tablets
Chew or crush tablets completely before swallowing.

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

**Talk to your doctor or other healthcare provider before using this product for your heart.

no print / no varnish area
lot no. & exp. date



8 40986 03221 9

Drug Facts (continued)

Active ingredients Aspirin 81 mg (NSAID) (nonsteroidal anti-inflammatory drug)

Questions or comments? 1-800-426-9391

Other information

- Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
- If pregnant or breast-feeding, ask a health professional before use. It is especially important to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
- Directions Do not take more than directed.
- Drink a full glass of water with each dose.
- Chew or crush tablets completely before swallowing. Do not swallow tablets whole.
- Adults and children 12 years and over: Take 4 to 6 chewable tablets every 4 hours not to exceed 6 tablets in 24 hours unless directed by a doctor.
- Children under 12 years: Do not use unless directed by a doctor.
- Store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F).
- See end flap for expiration date and lot number.

Disturbed By: Pharmacy Value Alliance, LLC
407 East Lancaster Avenue
Wayne, PA 19087

If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

©2014 Bayer AG, owner of the registered trademark Bayer®
Chewable Low Dose Aspirin, 50544 8970122821807

B-1590-218A-07-TB
REV0122821807

Drug Facts (continued)

Allergy alert: Aspirin may cause a severe allergic reaction, which may include hives, facial swelling, or throat swelling.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you take a blood thinning (anticoagulant) or steroid drug, are age 60 or older, or have other drugs containing prescription or over-the-counter NSAIDs (aspirin, ibuprofen, naproxen, or others).

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver disease, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- warfarin

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - feel faint
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present

Premier Value 44-218

ASPIRIN LOW DOSE

aspirin tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68016-690
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
DEXTROSE MONOHYDRATE (UNII: LX22YL083G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	orange	Score	no score
Shape	ROUND	Size	8mm
Flavor	ORANGE	Imprint Code	44;218
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68016-690-36	1 in 1 CARTON	09/12/1990	
1		36 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:68016-690-18	3 in 1 CARTON	09/12/1990	
2		36 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	09/12/1990	

Labeler - Chain Drug Consortium (101668460)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(68016-690)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(68016-690)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(68016-690)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(68016-690)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(68016-690)

Revised: 9/2023

Chain Drug Consortium