SU PHEDRINE PE NON-DROWSY, MAXIMUM STRENGTH- phenylephrine hcl tablet, film coated Salado Sales. Inc.

CVP 44-453

Active ingredient (in each tablet)

Phenylephrine HCl 10 mg

Purpose

Nasal decongestant

Uses

- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves sinus congestion and pressure

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- diabetes
- thyroid disease
- high blood pressure
- difficulty in urination due to enlargement of the prostate gland

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with fever

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- adults and children 12 years and over: take 1 tablet every 4 hours. Do not take more than 6 tablets in 24 hours.
- children under 12 years: ask a doctor

Other information

- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

croscarmellose sodium, dextrose monohydrate, dibasic calcium phosphate dihydrate, FD&C red #40, lecithin, magnesium stearate, maltodextrin, microcrystalline cellulose, silicon dioxide, sodium carboxymethylcellulose, sodium citrate dihydrate, titanium dioxide

Questions or comments?

1-800-426-9391

Principal Display Panel

CVP® HEALTH

COMPARE TO SUDAFED PE® CONGESTION ACTIVE INGREDIENT*

NON-DROWSY
SU•PHEDRINE PE
PHENYLEPHRINE HYDROCHLORIDE
NASAL DECONGESTANT

18 TABLETS 10 mg EACH 1 PILL PER DOSE

MAXIMUM STRENGTH

RELIEVES NASAL &
SINUS CONGESTION
CONTAINS NO PSEUDOEPHEDRINE

TAMPER EVIDENT: DO NOT USE IF

PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Sudafed PE® Congestion. 50844 REV0118M45344

Distributed by Consumer Value Products, Inc. P.O. Box 6115, Temple, Texas 76502 CVPproducts.com

NON-DROWSY SU·PHEDRINE PE

PHENYLEPHRINE HYDROCHLORIDE



no. & Expiration Date

ot

No Print Area




NON-DROWSY

PHENYLEPHRINE HYDROCHLORIDE NASAL DECONGESTANT

MAXIMUM STRENGTH

RELIEVES NASAL & SINUS CONGESTION

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING



CONTAINS NO PSEUDOEPHEDRINE

CVPproducts.com Distributed by Consumer Value Products, Inc. P.O. Box 6115, Temple, Texas 76502

Stop use and ask a doctor if

■ heart disease

■ thyroid disease ■ high blood pressure adiab etes ■

Ask a doctor before use if you have

20844 Sudafed PE® Congestion. REVOT18M45344 Johnson Corporation, owner of the registered trademark *This product is not manufactured or distributed by Johnson &

■ symptoms do not improve within 7 days or occur with fever ■ nervousness, dizziness, or sleeplessness occur

When using this product do not exceed recommended dosage.

■ difficulty in urination due to enlargement of the prostate gland



B-0226-453-44-RR REV0118M45344

Questions or comments? 1-800-426-9391

citrate dihydrate, titanium dioxide cellulose, silicon dioxide, sodium carboxymethylcellulose, sodium lecithin, magnesium stearate, maltodextrin, microcrystalline monohydrate, dibasic calcium phosphate dihydrate, FD&C red #40, Inactive ingredients croscamellose sodium, dextrose

- see end flap for expiration date and lot number (4°88-°83)
- store at 25°C (77°F); excursions permitted between 15°-30°C OPENED OR BLISTER IS TORN OR BROKEN
 - TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS Other information
 - children under 12 years: ask a doctor hours. Do not take more than 6 tablets in 24 hours.
 - adults and children 12 years and over: take 1 tablet every 4 Directions

help or contact a Poison Control Center (1-800-222-1222) right Keep out of reach of children. In case of overdose, get medical

If pregnant or breast-feeding, ask a health professional before Drug Facts (continued)

Do not use it you are now taking a prescription monoamine

oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Warnings

■ remporanly relieves sinus congestion and pressure hay fever or other upper respiratory allergies

 femborarity relieves nasal congestion due to the common cold, sasn

nasai decongestant Phenylephrine HCI 10 mg. Active ingredient (in each tablet)

Drug Facts

Purpose

SU PHEDRINE PE NON-DROWSY, MAXIMUM STRENGTH

phenylephrine hcl tablet, film coated

Pro	duct	Inform	ation
FIU	uuct		Iativii

Product Type HUMAN OTC DRUG Item Code (Source) NDC:57243-453

Route of Administration ORAL

Active Ingredient/Active Moiety Ingredient Name Basis of Strength PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE UNII:1WS297W6MV) PHENYLEPHRINE HYDROCHLORIDE 10 mg

Inactive Ingredients	
Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)	
DEXTROSE MONOHYDRATE (UNII: LX22YL083G)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K6790BS311)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics				
Color	red	Score	no score	
Shape	ROUND	Size	7mm	
Flavor		Imprint Code	44;453	
Contains				

F	Packaging						
#	tem Code	Package Description	Marketing Start Date	Marketing End Date			
1	NDC:57243- 453-44	1 in 1 CARTON	01/14/2005				
1		18 in 1 BLISTER PACK; Type 0: Not a Combination Product					

Marketing Information					
	Marketing	Application Number or Monograph	Marketing Start	Marketing End	

Category	Citation	Date	Date
OTC Monograph Drug	M012	01/14/2005	

Labeler - Salado Sales, Inc. (009830555)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(57243-453) , pack(57243-453)

Establishment					
Name	Address	ID/FEI	Business Operations		
LNK International, Inc.		832867894	manufacture(57243-453)		

Establishment					
Name	Address	ID/FEI	Business Operations		
LNK International, Inc.		868734088	manufacture(57243-453)		

Establishment					
Name	Address	ID/FEI	Business Operations		
LNK International, Inc.		117025878	manufacture(57243-453)		

Revised: 12/2023 Salado Sales, Inc.