

TOPCARE ULTRA STRENGTH STOMACH RELIEF - bismuth subsalicylate liquid
Topco Associates, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Topcare Ultra Strength Stomach Relief Liquid

ACTIVE INGREDIENT(in each 15 mL)

Bismuth subsalicylate 525 mg

PURPOSE

Upset stomach reliever and anti-diarrheal

USE(S)

relieves:

- diarrhea
- heartburn
- indigestion
- nausea
- upset stomach associated with these symptoms

WARNINGS

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

DO NOT USE IF YOU HAVE

- an ulcer
- bloody or black stool
- a bleeding problem

ASK A DOCTOR BEFORE USE IF YOU HAVE

- fever
- mucus in the stool

ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE

taking any drug for

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

WHEN USING THIS PRODUCT

a temporary, but harmless darkening of the stool and/or tongue may occur

STOP USE AND ASK DOCTOR IF

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

IF PREGNANT OR BREAST-FEEDING,

ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN

In case of overdose, get medical help or contact a Poison Control Center immediately.

DIRECTIONS

- **shake well before use**
- mL = milliliter
- TBSP = tablespoon
- adults and children 12 years and over: 1 dose (1 TBSP or 15 mL) every 1/2 hour as needed
- do not exceed 8 doses (120 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of fluids to help prevent dehydration caused by diarrhea

OTHER INFORMATION

- **each 15 mL or 1 TBSP contains:**

- potassium 15 mg
- salicylate 236 mg
- sodium 5 mg
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)
- dosage cup provided

INACTIVE INGREDIENTS

benzoic acid, D&C red # 22, D&C red # 28, flavor, hydroxyethyl cellulose, potassium hydroxide, purified water, saccharin sodium, salicylic acid, simethicone, xanthan gum

PRINCIPAL DISPLAY PANEL

NDC 76162-022-05

COMPARE TO PEPTO-BISMOL® ULTRA STRENGTH ACTIVE INGREDIENT*

TopCare

ULTRA STRENGTH

Stomach Relief

BISMUTH SUBSALICYLATE 525 mg per 15 mL

UPSET STOMACH RELIEVER/ANTIDIARRHEAL

5 SYMPTOM DIGESTIVE RELIEF

- Heartburn
- Indigestion
- Nausea
- Upset stomach
- Diarrhea

12 FL OZ (354 mL)

Drug Facts **DO NOT USE IF IMPRINTED SHRINKBRAND IS MISSING OR BROKEN.**

Active ingredient (in each 15 mL) *Purpose*
 Bismuth subsalicylate 525 mgUpset stomach reliever and anti-diarrheal

Uses relieves: ■ diarrhea ■ heartburn ■ indigestion ■ nausea ■ upset stomach associated with these symptoms

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. **Do not take if you are**
 ■ allergic to salicylates (including aspirin)
 ■ taking other salicylate products

Do not use if you have ■ bloody or black stool ■ an ulcer ■ a bleeding problem

Ask a doctor before use if you have ■ fever ■ mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for
 ■ anticoagulation (thinning of the blood)
 ■ diabetes ■ gout ■ arthritis

When using this product a temporary, but harmless darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if ■ symptoms get worse ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center immediately.

This product is not manufactured or distributed by Procter & Gamble, Inc., the distributor of Pepto-Bismol® Ultra Strength.

NDC 76162-022-05

COMPARE TO PEPTO-BISMOL® ULTRA STRENGTH ACTIVE INGREDIENT*

ULTRA STRENGTH Stomach Relief

BISMUTH SUBSALICYLATE 525mg per 15 mL
 UPSET STOMACH RELIEVER/ANTIDIARRHEAL
 5 SYMPTOM DIGESTIVE RELIEF

- Heartburn
- Indigestion
- Upset Stomach
- Nausea
- Diarrhea

12 FL OZ (354 mL)

Drug Facts (continued)

Directions ■ shake well before use ■ mL = milliliter
 ■ TBSP = tablespoon ■ adults and children 12 years and over: 1 dose (1 TBSP or 15 mL) every 1/2 hour as needed ■ do not exceed 8 doses (120 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ children under 12 years: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information ■ each 15 mL or 1 TBSP contains:
 • potassium 15 mg • salicylate 236 mg • sodium 5 mg
 ■ protect from freezing. ■ avoid excessive heat (over 104°F or 40°C). ■ dosage cap provided

Inactive ingredients benzoic acid, D&C red # 22, D&C red # 28, flavor, hydroxyethyl cellulose, potassium hydroxide, purified water, saccharin sodium, salicylic acid, simethicone, xanthan gum.

DISTRIBUTED BY TOPCO ASSOCIATES LLC, ELK GROVE VILLAGE, IL 60007
 ©TOPCO GUAA1021 QUESTIONS? 1-888-423-0139
 topcare@topco.com www.topcarebrand.com

Scan here for more information or call 1-888-423-0139

✓ **QUALITY GUARANTEED**

MFR# 53041 REV 1021

80% UPC SYMBOL FOR POSITION ONLY

verify UPC number against previous printing and packaging change form

0 36800 13585 7

◀ **Drug Facts continued**

TOPCARE ULTRA STRENGTH STOMACH RELIEF

bismuth subsalicylate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:76162-022
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
BENZOIC ACID (UNII: 8SKN0B0MIM)	
D&C RED NO. 22 (UNII: 1678RKX8RT)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SALICYLIC ACID (UNII: O414PZ4LPZ)	
HYDROXYETHYL CELLULOSE (1500 MPA.S AT 1%) (UNII: L605B5892V)	

DIMETHICONE (UNII: 92RU3N3Y1O)

XANTHAN GUM (UNII: TTV12P4NEE)

Product Characteristics

Color	PINK	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76162-022-05	354 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/05/2022	
2	NDC:76162-022-04	236 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/05/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part335	01/05/2022	

Labeler - Topco Associates, LLC (006935977)

Revised: 11/2022

Topco Associates, LLC