

**METHAMPHETAMINE HYDROCHLORIDE- methamphetamine hydrochloride tablet**  
**West-Ward Pharmaceuticals Corp.**


-----  
**Methamphetamine Hydrochloride Tablets USP, CII**

**Rx only**

**METHAMPHETAMINE HAS A HIGH POTENTIAL FOR ABUSE. PARTICULAR ATTENTION SHOULD BE PAID TO THE POSSIBILITY OF SUBJECTS OBTAINING METHAMPHETAMINE FOR NON-THERAPEUTIC USE OR DISTRIBUTION TO OTHERS, AND THE DRUG SHOULD BE PRESCRIBED OR DISPENSED SPARINGLY. MISUSE OF METHAMPHETAMINE MAY CAUSE SUDDEN DEATH AND SERIOUS CARDIOVASCULAR ADVERSE EVENTS.**

**DESCRIPTION**

Methamphetamine hydrochloride, USP, chemically known as Benzeneethanamine, N,  $\alpha$ -dimethyl-, hydrochloride (1:1), ( $\alpha$ S)-, is a member of the amphetamine group of sympathomimetic amines. It has the following structural formula:



Methamphetamine hydrochloride tablets USP, contain 5 mg of methamphetamine hydrochloride USP for oral administration. In addition each tablet contains the following inactive ingredients: crospovidone, magnesium stearate, microcrystalline cellulose, povidone and purified water.

**CLINICAL PHARMACOLOGY**

Methamphetamine is a sympathomimetic amine with CNS stimulant activity. Peripheral actions include elevation of systolic and diastolic blood pressures and weak bronchodilator and respiratory stimulant action. Other central nervous system actions, or metabolic effects, may be involved, for example.

The mechanism of action involved in producing the beneficial behavioral changes seen in hyperkinetic children receiving methamphetamine is unknown.

In humans, methamphetamine is rapidly absorbed from the gastrointestinal tract. The primary site of metabolism is in the liver by aromatic hydroxylation, N-dealkylation and deamination. At least seven metabolites have been identified in the urine. The biological half-life has been reported in the range of 4 to 5 hours. Excretion occurs primarily in the urine and is dependent on urine pH. Alkaline urine will significantly increase the drug half-life. Approximately 62% of an oral dose is eliminated in the urine

within the first 24 hours with about one-third as intact drug and the remainder as metabolites.

## INDICATIONS AND USAGE

### Attention Deficit Disorder with Hyperactivity

Methamphetamine hydrochloride tablets are indicated as an integral part of a total treatment program which typically includes other remedial measures (psychological, educational, social) for a stabilizing effect in children over 6 years of age with a behavioral syndrome characterized by the following group of developmentally inappropriate symptoms: moderate to severe distractibility, short attention span, hyperactivity, emotional lability, and impulsivity. The diagnosis of this syndrome should not be made with finality when these symptoms are only of comparatively recent origin. Nonlocalizing (soft) neurological signs, learning disability, and abnormal EEG may or may not be present, and a diagnosis of central nervous system dysfunction may or may not be warranted.

## CONTRAINDICATIONS

In patients known to be hypersensitive to amphetamine, or other components of methamphetamine. Hypersensitivity reactions such as angioedema and anaphylactic reactions have been reported in patients treated with other amphetamine products (see ADVERSE REACTIONS).

Patients taking monoamine oxidase inhibitors (MAOIs), or within 14 days of stopping MAOIs (including MAOIs such as linezolid or intravenous methylene blue), because of an increased risk of hypertensive crisis (see WARNINGS and Drug Interactions). It is also contraindicated in patients with glaucoma, advanced arteriosclerosis, symptomatic cardiovascular disease, moderate to severe hypertension, hyperthyroidism or known hypersensitivity or idiosyncrasy to sympathomimetic amines. Methamphetamine should not be given to patients who are in an agitated state or who have a history of drug abuse.

## WARNINGS

### Serious Cardiovascular Events

#### *Sudden Death and Pre-existing Structural Cardiac Abnormalities or Other Serious Heart Problems*

- **Children and Adolescents:** Sudden death has been reported in association with CNS stimulant treatment at usual doses in children and adolescents with structural cardiac abnormalities or other serious heart problems. Although some serious heart problems alone carry an increased risk of sudden death, stimulant products generally should not be used in children or adolescents with known serious structural cardiac abnormalities, cardiomyopathy, serious heart rhythm abnormalities, or other serious cardiac problems that may place them at increased vulnerability to the sympathomimetic effects of a stimulant drug.
- **Adults:** Sudden deaths, stroke, and myocardial infarction have been reported in adults taking stimulant drugs at usual doses for ADHD. Although the role of stimulants in these adult cases is also unknown, adults have a greater likelihood than children of having serious structural cardiac abnormalities, cardiomyopathy, serious heart rhythm abnormalities, coronary artery disease, or other serious cardiac problems. Adults with such abnormalities should also generally not be treated with stimulant drugs.

#### *Hypertension and other Cardiovascular Conditions*

Stimulant medications cause a modest increase in average blood pressure (about 2 to 4 mmHg) and average heart rate (about 3 to 6 bpm), and individuals may have larger increases. While the mean changes alone would not be expected to have short-term consequences, all patients should be monitored

for larger changes in heart rate and blood pressure. Caution is indicated in treating patients whose underlying medical conditions might be compromised by increases in blood pressure or heart rate, e.g., those with pre-existing hypertension, heart failure, recent myocardial infarction, or ventricular arrhythmia.

### *Assessing Cardiovascular Status in Patients being Treated with Stimulant Medications*

Children, adolescents, or adults who are being considered for treatment with stimulant medications should have a careful history (including assessment for a family history of sudden death or ventricular arrhythmia) and physical exam to assess for the presence of cardiac disease, and should receive further cardiac evaluation if findings suggest such disease (e.g., electrocardiogram and echocardiogram). Patients who develop symptoms such as exertional chest pain, unexplained syncope, or other symptoms suggestive of cardiac disease during stimulant treatment should undergo a prompt cardiac evaluation.

### *Psychiatric Adverse Events*

Pre-existing Psychosis: Administration of stimulants may exacerbate symptoms of behavior disturbance and thought disorder in patients with a pre-existing psychotic disorder.

Bipolar Illness: Particular care should be taken in using stimulants to treat ADHD in patients with comorbid bipolar disorder because of concern for possible induction of a mixed/manic episode in such patients. Prior to initiating treatment with a stimulant, patients with comorbid depressive symptoms should be adequately screened to determine if they are at risk for bipolar disorder; such screening should include a detailed psychiatric history, including a family history of suicide, bipolar disorder, and depression.

Emergence of New Psychotic or Manic Symptoms: Treatment emergent psychotic or manic symptoms, e.g., hallucinations, delusional thinking, or mania in children and adolescents without a prior history of psychotic illness or mania can be caused by stimulants at usual doses. If such symptoms occur, consideration should be given to a possible causal role of the stimulant, and discontinuation of treatment may be appropriate. In a pooled analysis of multiple short-term, placebo-controlled studies, such symptoms occurred in about 0.1% (4 patients with events out of 3482 exposed to methylphenidate or amphetamine for several weeks at usual doses) of stimulant-treated patients compared to 0 in placebo-treated patients.

Aggression: Aggressive behavior or hostility is often observed in children and adolescents with ADHD, and has been reported in clinical trials and the postmarketing experience of some medications indicated for the treatment of ADHD. Although there is no systematic evidence that stimulants cause aggressive behavior or hostility, patients beginning treatment for ADHD should be monitored for the appearance of or worsening of aggressive behavior or hostility.

Long-Term Suppression of Growth: Careful follow-up of weight and height in children ages 7 to 10 years who were randomized to either methylphenidate or non-medication treatment groups over 14 months, as well as in naturalistic subgroups of newly methylphenidate-treated and non-medication treated children over 36 months (to the ages of 10 to 13 years), suggests that consistently medicated children (i.e., treatment for 7 days per week throughout the year) have a temporary slowing in growth rate (on average, a total of about 2 cm less growth in height and 2.7 kg less growth in weight over 3 years), without evidence of growth rebound during this period of development. Published data are inadequate to determine whether chronic use of amphetamines may cause a similar suppression of growth, however, it is anticipated that they likely have this effect as well. Therefore, growth should be monitored during treatment with stimulants, and patients who are not growing or gaining height or weight as expected may need to have their treatment interrupted.

Seizures: There is some clinical evidence that stimulants may lower the convulsive threshold in patients with prior history of seizures, in patients with prior EEG abnormalities in absence of seizures, and, very rarely, in patients without a history of seizures and no prior EEG evidence of seizures. In the presence of seizures, the drug should be discontinued.

Peripheral Vasculopathy, including Raynaud's phenomenon: Stimulants, including methamphetamine, used to treat ADHD are associated with peripheral vasculopathy, including Raynaud's phenomenon. Signs and symptoms are usually intermittent and mild; however, very rare sequelae include digital ulceration and/or soft tissue breakdown. Effects of peripheral vasculopathy, including Raynaud's phenomenon, were observed in post-marketing reports at different times and at therapeutic doses in all age groups throughout the course of treatment. Signs and symptoms generally improve after reduction in dose or discontinuation of drug. Careful observation for digital changes is necessary during treatment with ADHD stimulants. Further clinical evaluation (e.g., rheumatology referral) may be appropriate for certain patients.

Serotonin Syndrome: Serotonin syndrome, a potentially life-threatening reaction, may occur when amphetamines are used in combination with other drugs that affect the serotonergic neurotransmitter systems such as monoamine oxidase inhibitors (MAOIs), selective serotonin reuptake inhibitors (SSRIs), serotonin norepinephrine reuptake inhibitors (SNRIs), triptans, tricyclic antidepressants, fentanyl, lithium, tramadol, tryptophan, buspirone, and St. John's Wort (see Drug Interactions). Amphetamines and amphetamine derivatives are known to be metabolized, to some degree, by cytochrome P450 2D6 (CYP2D6) and display minor inhibition of CYP2D6 metabolism (see CLINICAL PHARMACOLOGY). The potential for a pharmacokinetic interaction exists with the co-administration of CYP2D6 inhibitors which may increase the risk with increased exposure to methamphetamine. In these situations, consider an alternative nonserotonergic drug or an alternative drug that does not inhibit CYP2D6 (see Drug Interactions).

Serotonin syndrome symptoms may include mental status changes (e.g., agitation, hallucinations, delirium, and coma), autonomic instability (e.g., tachycardia, labile blood pressure, dizziness, diaphoresis, flushing, hyperthermia), neuromuscular symptoms (e.g., tremor, rigidity, myoclonus, hyperreflexia, incoordination), seizures, and/or gastrointestinal symptoms (e.g., nausea, vomiting, diarrhea).

Concomitant use of methamphetamine with MAOI drugs is contraindicated (see CONTRAINDICATIONS).

Discontinue treatment with methamphetamine and any concomitant serotonergic agents immediately if the above symptoms occur, and initiate supportive symptomatic treatment. If concomitant use of methamphetamine with other serotonergic drugs or CYP2D6 inhibitors is clinically warranted, initiate methamphetamine with lower doses, monitor patients for the emergence of serotonin syndrome during drug initiation or titration, and inform patients of the increased risk for serotonin syndrome.

Visual Disturbance: Difficulties with accommodation and blurring of vision have been reported with stimulant treatment.

## **PRECAUTIONS**

### **General**

Methamphetamine hydrochloride tablets should be used with caution in patients with even mild hypertension.

Methamphetamine should not be used to combat fatigue or to replace rest in normal persons.

Prescribing and dispensing of methamphetamine should be limited to the smallest amount that is feasible at one time in order to minimize the possibility of overdose.

### **Information for Patients**

The patient should be informed that methamphetamine may impair the ability to engage in potentially hazardous activities, such as, operating machinery or driving a motor vehicle.

*Circulation Problems in Fingers and Toes [Peripheral Vasculopathy, Including Raynaud's*

## Phenomenon]

- Instruct patients beginning treatment with methamphetamine about the risk of peripheral vasculopathy, including Raynaud's Phenomenon, and associated signs and symptoms: fingers or toes may feel numb, cool, painful, and/or may change color from pale, to blue, to red.
- Instruct patients to report to their physician any new numbness, pain, skin color change, or sensitivity to temperature in fingers or toes.
- **Instruct patients to call their physician immediately with any signs of unexplained wounds appearing on fingers or toes while taking methamphetamine.**
- Further clinical evaluation (e.g., rheumatology referral) may be appropriate for certain patients.

The patient should be cautioned not to increase dosage, except on advice of the physician.

Prescribers or other health professionals should inform patients, their families, and their caregivers about the benefits and risks associated with treatment with methamphetamine and should counsel them in its appropriate use. A patient Medication Guide is available for methamphetamine hydrochloride tablets. The prescriber or health professional should instruct patients, their families, and their caregivers to read the Medication Guide and should assist them in understanding its contents. Patients should be given the opportunity to discuss the contents of the Medication Guide and to obtain answers to any questions they may have. The complete text of the Medication Guide is available at <https://dailymed.nlm.nih.gov>.

### Drug Interactions

Insulin requirements in diabetes mellitus may be altered in association with the use of methamphetamine and the concomitant dietary regimen.

Methamphetamine may decrease the hypotensive effect of *guanethidine*.

Methamphetamine should not be used concurrently with *monoamine oxidase inhibitors* (see **CONTRAINDICATIONS**).

Concurrent administration of *tricyclic antidepressants* and indirect-acting sympathomimetic amines such as the amphetamines, should be closely supervised and dosage carefully adjusted.

*Phenothiazines* are reported in the literature to antagonize the CNS stimulant action of the amphetamines.

### Drug/Laboratory Test Interactions

Literature reports suggest that amphetamines may be associated with significant elevation of plasma corticosteroids. This should be considered if determination of plasma corticosteroid levels is desired in a person receiving amphetamines.

### Acidifying Agents

Lower blood levels and efficacy of amphetamines. Increase dose based on clinical response. Examples of acidifying agents include gastrointestinal acidifying agents (e.g., *guanethidine*, *reserpine*, *glutamic acid HCl*, *ascorbic acid*) and urinary acidifying agents (e.g., *ammonium chloride*, *sodium acid phosphate*, *methenamine salts*).

### Alkalinizing Agents

Increase blood levels and potentiate the action of amphetamine. Co-administration of methamphetamine and gastrointestinal alkalinizing agents should be avoided. Examples of alkalinizing agents include gastrointestinal alkalinizing agents (e.g., *sodium bicarbonate*) and urinary alkalinizing agents (e.g., *acetazolamide*, some *thiazides*).

### Tricyclic Antidepressants

May enhance the activity of tricyclic or sympathomimetic agents causing striking and sustained

increases in the concentration of d-amphetamine in the brain; cardiovascular effects can be potentiated. Monitor frequently and adjust or use alternative therapy based on clinical response. Examples of tricyclic antidepressants include desipramine, Protriptyline.

### **CYP2D6 Inhibitors**

The concomitant use of methamphetamine and CYP2D6 inhibitors may increase the exposure of methamphetamine compared to the use of the drug alone and increase the risk of serotonin syndrome. Initiate with lower doses and monitor patients for signs and symptoms of serotonin syndrome particularly during methamphetamine initiation and after a dosage increase. If serotonin syndrome occurs, discontinue methamphetamine and the CYP2D6 inhibitor (see WARNINGS, OVERDOSAGE). Examples of CYP2D6 Inhibitors include paroxetine and fluoxetine (also serotonergic drugs), quinidine, ritonavir.

### **Serotonergic Drugs**

The concomitant use of methamphetamine and serotonergic drugs increases the risk of serotonin syndrome. Initiate with lower doses and monitor patients for signs and symptoms of serotonin syndrome, particularly during methamphetamine initiation or dosage increase. If serotonin syndrome occurs, discontinue methamphetamine and the concomitant serotonergic drug(s) (see WARNINGS and PRECAUTIONS). Examples of serotonergic drugs include selective serotonin reuptake inhibitors (SSRI), serotonin norepinephrine reuptake inhibitors (SNRI), triptans, tricyclic antidepressants, fentanyl, lithium, tramadol, tryptophan, buspirone, St. John's Wort.

### **MAO Inhibitors**

Concomitant use of MAOIs and CNS stimulants can cause hypertensive crisis. Potential outcomes include death, stroke, myocardial infarction, aortic dissection, ophthalmological complications, eclampsia, pulmonary edema, and renal failure. Do not administer methamphetamine concomitantly or within 14 days after discontinuing MAOI (see CONTRAINDICATIONS and WARNINGS). Examples of MAOIs include selegiline, tranylcypromine, isocarboxazid, phenelzine, linezolid, methylene blue.

### **Proton Pump Inhibitors**

Time to maximum concentration ( $T_{max}$ ) of amphetamine is decreased compared to when administered alone. Monitor patients for changes in clinical effect and adjust therapy based on clinical response. An example of a proton pump inhibitor is omeprazole.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

Data are not available on long-term potential for carcinogenicity, mutagenicity, or impairment of fertility.

### **Pregnancy**

#### *Teratogenic Effects*

Pregnancy Category C: Methamphetamine has been shown to have teratogenic and embryocidal effects in mammals given high multiples of the human dose. There are no adequate and well-controlled studies in pregnant women. Methamphetamine hydrochloride tablets should not be used during pregnancy unless the potential benefit justifies the potential risk to the fetus.

#### *Nonteratogenic Effects*

Infants born to mothers dependent on amphetamines have an increased risk of premature delivery and low birth weight. Also, these infants may experience symptoms of withdrawal as demonstrated by dysphoria, including agitation and significant lassitude.

### **Usage in Nursing Mothers**

Amphetamines are excreted in human milk. Mothers taking amphetamines should be advised to refrain from nursing.

### **Pediatric Use**

Long-term effects of methamphetamine in children have not been established (see **WARNINGS**).

Drug treatment is not indicated in all cases of the behavioral syndrome characterized by moderate to severe distractibility, short attention span, hyperactivity, emotional lability and impulsivity. It should be considered only in light of the complete history and evaluation of the child. The decision to prescribe methamphetamine hydrochloride tablets should depend on the physician's assessment of the chronicity and severity of the child's symptoms and their appropriateness for his/her age. Prescription should not depend solely on the presence of one or more of the behavioral characteristics.

When these symptoms are associated with acute stress reactions, treatment with methamphetamine hydrochloride tablets is usually not indicated.

Clinical experience suggests that in psychotic children, administration of methamphetamine hydrochloride tablets may exacerbate symptoms of behavior disturbance and thought disorder.

Amphetamines have been reported to exacerbate motor and phonic tics and Tourette's syndrome. Therefore, clinical evaluation for tics and Tourette's syndrome in children and their families should precede use of stimulant medications.

### **Geriatric Use**

Clinical Studies of methamphetamine did not include sufficient numbers of subjects age 65 years and over to determine whether elderly subjects respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal or cardiac function, and of concomitant disease or other drug therapy observed in this population.

## **ADVERSE REACTIONS**

The following are adverse reactions in decreasing order of severity within each category that have been reported:

*Cardiovascular:* Elevation of blood pressure, tachycardia and palpitation. Fatal cardiorespiratory arrest has been reported, mostly in the context of abuse/misuse.

*Central Nervous System:* Psychotic episodes have been rarely reported at recommended doses. Dizziness, dysphoria, overstimulation, euphoria, insomnia, tremor, restlessness and headache. Exacerbation of motor and phonic tics and Tourette's syndrome.

*Gastrointestinal:* Diarrhea, constipation, dryness of mouth, unpleasant taste and other gastrointestinal disturbances.

*Hypersensitivity:* Urticaria.

*Endocrine:* Impotence and changes in libido; frequent or prolonged erections.

*Musculoskeletal:* Rhabdomyolysis.

*Miscellaneous:* Suppression of growth has been reported with the long-term use of stimulants in children (see **WARNINGS**).

*Skin and Subcutaneous Tissue Disorders:* Alopecia.

**To report SUSPECTED ADVERSE REACTIONS, contact West-Ward Pharmaceuticals Corp. at 1-800-962-8364 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

## **DRUG ABUSE AND DEPENDENCE**

### **Controlled Substance**

Methamphetamine hydrochloride tablets are subject to control under DEA schedule II.

### **Abuse**

Methamphetamine has been extensively abused. Tolerance, extreme psychological dependence, and severe social disability have occurred. There are reports of patients who have increased the dosage to many times that recommended. Abrupt cessation following prolonged high dosage administration results in extreme fatigue and mental depression; changes are also noted on the sleep EEG. Manifestations of chronic intoxication with methamphetamine include severe dermatoses, marked insomnia, irritability, hyperactivity, and personality changes. The most severe manifestation of chronic intoxication is psychosis often clinically indistinguishable from schizophrenia. Abuse and/or misuse of methamphetamine have resulted in death. Fatal cardiorespiratory arrest has been reported in the context of abuse and/or misuse of methamphetamine.

## **OVERDOSAGE**

Manifestations of amphetamine overdose include restlessness, tremor, hyperreflexia, rapid respiration, confusion, assaultiveness, hallucinations, panic states, hyperpyrexia, and rhabdomyolysis. Fatigue and depression usually follow the central nervous system stimulation. Serotonin syndrome has also been reported. Cardiovascular effects include arrhythmias, hypertension or hypotension and circulatory collapse. Gastrointestinal symptoms include nausea, vomiting, diarrhea and abdominal cramps. Fatal poisoning is usually preceded by convulsions and coma.

### **Treatment**

Consult with a Certified Poison Control Center for up to date guidance and advice.

## **DOSAGE AND ADMINISTRATION**

Methamphetamine hydrochloride tablets are given orally.

Methamphetamine should be administered at the lowest effective dosage, and dosage should be individually adjusted. Late evening medication should be avoided because of the resulting insomnia.

### **Attention Deficit Disorder with Hyperactivity**

For treatment of children 6 years or older with a behavioral syndrome characterized by moderate to severe distractibility, short attention span, hyperactivity, emotional lability and impulsivity:

An initial dose of 5 mg methamphetamine once or twice a day is recommended. Daily dosage may be raised in increments of 5 mg at weekly intervals until an optimum clinical response is achieved. The usual effective dose is 20 to 25 mg daily. The total daily dose may be given in two divided doses daily.

Where possible, drug administration should be interrupted occasionally to determine if there is a recurrence of behavioral symptoms sufficient to require continued therapy

## **HOW SUPPLIED**

### **Methamphetamine Hydrochloride Tablets USP**

**5 mg tablets are supplied as a round, white to off-white, biconvex, debossed tablet with "54" on one side and "681" on the other side.**

NDC 0054-0389-25: Bottle of 100 tablets



## Storage

Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]

Dispense in a tight, light resistant, child-resistant container as defined in the USP/NF.

Distr. by: **West-Ward**

**Pharmaceuticals Corp.**

Eatontown, NJ 07724

**10008165/05**

**Revised January 2020**

## Medication Guide

### **Methamphetamine Hydrochloride Tablets USP, CII**

(meth" am fet' a meen hye" droe klor' ide)

#### **Rx only**

Read the Medication Guide that comes with methamphetamine hydrochloride tablets before you or your child starts taking it and each time you get a refill.

There may be new information. This Medication Guide does not take the place of talking to your or your child's doctor about your or your child's treatment with methamphetamine hydrochloride tablets.

#### **What is the most important information I should know about?**

**The following have been reported with use of methamphetamine hydrochloride and other stimulant medicines.**

##### **1. Heart-related problems:**

- sudden death in patients who have heart problems or heart defects
- stroke and heart attack in adults
- increased blood pressure and heart rate

Tell your or your child's doctor if you or your child have any heart problems, heart defects, high blood pressure, or a family history of these problems.

Your or your child's doctor should check you or your child carefully for heart problems before starting methamphetamine hydrochloride tablets.

Your or your child's doctor should check your or your child's blood pressure and heart rate regularly during treatment with methamphetamine hydrochloride tablets.

**Call your or your child's doctor right away if you or your child has any signs of heart problems such as chest pain, shortness of breath, or fainting while taking methamphetamine hydrochloride tablets.**

##### **2. Mental (Psychiatric) problems:**

###### **All Patients**

- new or worse behavior and thought problems
- new or worse bipolar illness
- new or worse aggressive behavior or hostility

###### **Children and Teenagers**

- new psychotic symptoms (such as hearing voices, believing things that are not true, are suspicious) or new manic symptoms

Tell your or your child's doctor about any mental problems you or your child have, or about a family history of suicide, bipolar illness, or depression.

**Call your or your child's doctor right away if you or your child have any new or worsening mental symptoms or problems while taking methamphetamine hydrochloride tablets, especially seeing or hearing things that are not real, believing things that are not real, or are suspicious.**

**3. Circulation problems in fingers and toes** [Peripheral vasculopathy, including Raynaud's phenomenon]:

- fingers or toes may feel numb, cool, painful
- fingers or toes may change color from pale, to blue, to red

Tell your doctor if you have or your child has numbness, pain, skin color change, or sensitivity to temperature in your fingers or toes.

**Call your or your child's doctor right away if you have or your child has any signs of unexplained wounds appearing on fingers or toes while taking methamphetamine hydrochloride tablets.**

### **What are methamphetamine hydrochloride tablets?**

Methamphetamine hydrochloride tablets are a central nervous system stimulant prescription medicine. **It is used for the treatment of Attention-Deficit Hyperactivity Disorder; (ADHD).** Methamphetamine hydrochloride tablets may help increase attention and decrease impulsiveness and hyperactivity in patients with ADHD.

Methamphetamine hydrochloride tablets should be used as a part of a total treatment program for ADHD that may include counseling or other therapies.

**Methamphetamine hydrochloride tablets are a federally controlled substance (CII) because it can be abused or lead to dependence. Keep methamphetamine hydrochloride tablets in a safe place to prevent misuse and abuse. Selling or giving away methamphetamine hydrochloride tablets may harm others, and is against the law.**

Tell your or your child's doctor if you or your child have (or have a family history of) ever abused or been dependent on alcohol, prescription medicines or street drugs.

### **Who should not take methamphetamine hydrochloride tablets?**

Methamphetamine hydrochloride tablets should not be taken if you or your child:

- have heart disease or hardening of the arteries
- have moderate to severe high blood pressure
- have hyperthyroidism
- have an eye problem called glaucoma
- are agitated
- have a history of drug abuse
- are taking or have taken within the past 14 days an antidepressant medicine called a monoamine oxidase inhibitor or MAOI.
- are sensitive to, allergic to, or had a reaction to other stimulant medicines

Methamphetamine hydrochloride tablets are not recommended for use in children less than 6 years old in the treatment of ADHD.

**Methamphetamine hydrochloride tablets may not be right for you or your child. Before starting methamphetamine hydrochloride tablets tell your or your child's doctor about all health conditions (or a family history of) including:**

- heart problems, heart defects, high blood pressure
- mental problems including psychosis, mania, bipolar illness or depression
- tics or Tourette's syndrome
- thyroid problems
- diabetes
- seizures or have had an abnormal brain wave test (EEG)
- circulation problems in fingers and toes

Tell your or your child's doctor if you or your child is pregnant, planning to become pregnant, or breastfeeding.

### **Can methamphetamine hydrochloride tablets be taken with other medicines?**

**Tell your or your child's doctor about all of the medicines that you or your child take including prescription and nonprescription medicines, vitamins, and herbal supplements.**

Methamphetamine hydrochloride tablets and some medicines may interact with each other and cause serious side effects. Sometimes the doses of other medicines will need to be adjusted while taking methamphetamine hydrochloride tablets.

Your or your child's doctor will decide whether methamphetamine hydrochloride tablets can be taken with other medicines.

### **Especially tell your or your child's doctor if you or your child takes:**

- anti-depression medicines including MAOIs
- anti-psychotic medicines
- blood pressure medicines
- insulin
- seizure medicines

Know the medicines that you or your child takes. Keep a list of your medicines with you to show your doctor and pharmacist.

**Do not start any new medicine while taking methamphetamine hydrochloride tablets without talking to your or your child's doctor first.**

### **How should be taken?**

- **Take methamphetamine hydrochloride tablets exactly as prescribed.** Your or your child's doctor may adjust the dose until it is right for you or your child.
- Methamphetamine hydrochloride tablets are usually taken 1 or 2 times each day.
- From time to time, your or your child's doctor may stop methamphetamine hydrochloride tablets treatment for a while to check ADHD symptoms.
- Your or your child's doctor may do regular checks of the blood, heart, and blood pressure while taking methamphetamine hydrochloride tablets. Children should have their height and weight checked often while taking methamphetamine hydrochloride tablets. Methamphetamine hydrochloride tablets treatment may be stopped if a problem is found during these check-ups.
- **If you or your child takes too much methamphetamine hydrochloride tablets or overdoses, call your or your child's doctor or poison control center right away, or get emergency treatment.**

### **What are possible side effects of methamphetamine hydrochloride tablets?**

See "**What is the most important information I should know about methamphetamine hydrochloride tablets?**" for information on reported heart and mental problems.

**Other serious side effects include:**

- slowing of growth (height and weight) in children
- seizures, mainly in patients with a history of seizures
- eyesight changes or blurred vision

**Common side effects include:**

- fast heart beat
- decreased appetite
- tremors
- headache
- trouble sleeping
- dizziness
- stomach upset
- weight loss
- dry mouth

Methamphetamine hydrochloride tablets may affect your or your child's ability to drive or do other dangerous activities.

Talk to your or your child's doctor if you or your child has side effects that are bothersome or do not go away.

This is not a complete list of possible side effects. Ask your or your child's doctor or pharmacist for more information.

Call your or your child's doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**How should I store methamphetamine hydrochloride tablets?**

- Store methamphetamine hydrochloride tablets at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.] Protect from light.
- **Keep methamphetamine hydrochloride tablets and all medicines out of the reach of children.**

**General information about methamphetamine hydrochloride tablets**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use methamphetamine hydrochloride tablets for a condition for which it was not prescribed. Do not give methamphetamine hydrochloride tablets to other people, even if they have the same condition. It may harm them and it is against the law.

This Medication Guide summarizes the most important information about methamphetamine hydrochloride tablets. If you would like more information, talk with your or your child's doctor. You can ask your or your child's doctor or pharmacist for information about methamphetamine hydrochloride tablets that was written for healthcare professionals.

For more information about methamphetamine hydrochloride tablets, contact West-Ward Pharmaceuticals Corp. at 1-800-962-8364.

**What are the ingredients in methamphetamine hydrochloride tablets?**

**Active Ingredient:** methamphetamine hydrochloride USP

**Inactive Ingredients:** crospovidone, magnesium stearate, microcrystalline cellulose, povidone, and purified water.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Distr. by: **West-Ward**

**Pharmaceuticals Corp.**

Eatontown, NJ 07724

10008165/05

Revised January 2020

**PACKAGE/LABEL PRINCIPAL DISPLAY PANEL**

**USUAL DOSAGE:** See Package Insert for Complete Prescribing Information. Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.] Dispense in a tight, light-resistant, child-resistant container as defined in the USP/NF. Do not accept if seal over bottle opening is broken or missing. Distr. by: **West-Ward Pharmaceuticals Corp.** Eatontown, NJ 07724

NDC 0054-0389-25 100 Tablets

**Methamphetamine Hydrochloride**   
**Tablets USP**

**5 mg**

**PHARMACIST:** Dispense with attached Medication Guide. Each tablet contains 5 mg methamphetamine hydrochloride USP.

 **WEST-WARD**  
A HENKA COMPANY

**Rx only**

3 N  
0054-0389-25  
3

10008166/02

**METHAMPHETAMINE HYDROCHLORIDE**

methamphetamine hydrochloride tablet

**Product Information**

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0054-0389
<b>Route of Administration</b>	ORAL	<b>DEA Schedule</b>	CII

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>METHAMPHETAMINE HYDROCHLORIDE</b> (UNII: 997F43Z9CV) (METHAMPHETAMINE - UNII:44RAL3456C)	METHAMPHETAMINE HYDROCHLORIDE	5 mg

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>CROSPVIDONE (15 MPAS AT 5%)</b> (UNII: 68401960MK)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	

WATER (UNII: 059QF0KO0R)

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	6mm
<b>Flavor</b>		<b>Imprint Code</b>	54;681
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0054-0389-25	100 in 1 BOTTLE; Type 0: Not a Combination Product	12/06/2016	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA203846	12/06/2016	

**Labeler** - West-Ward Pharmaceuticals Corp. (080189610)

### Establishment

Name	Address	ID/FEI	Business Operations
West-Ward Columbus Inc.		058839929	MANUFACTURE(0054-0389)

Revised: 2/2020

West-Ward Pharmaceuticals Corp.