

**VICKS VAPOCOOL SORE THROAT- benzocaine and menthol lozenge**  
**The Procter & Gamble Manufacturing Company**

-----  
**VICKS® VapoCOOL™ Sore Throat Lozenges**

***Drug Facts***

**Active ingredients (per lozenge)**

Benzocaine 15 mg

Menthol 20 mg

**Purpose**

Oral anesthetic

Oral anesthetic

**Uses**

temporarily relieves occasional minor irritation and pain due to

- sore throat
- sore mouth

**Warnings**

**Methemoglobinemia Warning-**

Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

**Allergy alert:**

do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other “caine” anesthetics. If a skin reaction occurs, stop use and seek medical help right away.

**Sore Throat Warning -**

Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 12 years of age.

**Do not use**

- for teething
- in children under 2

**When using this product**

do not exceed recommended dosage

**Ask a doctor before use if you have**

- a severe sore throat accompanied by difficulty in breathing or that lasts more than 2 days
- a sore throat accompanied by fever, headache, rash, swelling, nausea or vomiting

**Stop use and ask a doctor if**

- sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persists or worsens

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- adults and children 12 years and over: dissolve 1 lozenge slowly in the mouth. Repeat every 2 hours as needed or as directed by a doctor.
- children under 12 years: do not use.

**Other information**

- store at room temperature; protect from moisture
- contains soy

**Inactive ingredients**

corn syrup, flavors, propylene glycol, sucrose

**Questions?**

**1-800-283-4879**

**DIST. BY PROCTER & GAMBLE,**

CINCINNATI OH 45202

**PRINCIPAL DISPLAY PANEL - 16 Lozenges Carton**

**VICKS®**

**VapoCOOL**

**SORE THROAT**

BENZOCAINE & MENTHOL - ORAL ANESTHETICS

COOLING + NUMBING

Soothes Throat Pain

Fast-acting

**WINTERFROST**

**16 LOZENGES**

**VICKS® VapoCOOL™ SORE THROAT**

**VICKS®**

**VapoCOOL™ SORE THROAT**  
BENZOCAINE & MENTHOL - ORAL ANESTHETICS

**COOLING + NUMBING**

Soothes Throat Pain  
Fast-acting

**WINTERFROST**  
**16 LOZENGES**

**Drug Facts**

Active ingredients (per lozenge)	Purpose
Benzocaine 15 mg	Oral anesthetic
Menthol 20 mg	Oral anesthetic

**Uses** temporarily relieves occasional minor irritation and pain due to • sore throat • sore mouth

**Warnings**  
**Methemoglobinemia Warning** – Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: • pale, gray or blue colored skin (cyanosis) • headache • rapid heart rate • shortness of breath • dizziness or lightheadedness • fatigue or lack of energy

**Allergy alert:** do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics. If a skin reaction occurs, stop use and seek medical help right away.

**Sore Throat Warning**– Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult a doctor promptly. Do not use more than 2 days or administer to children under 12 years of age.

**Do not use** • for teething • in children under 2

**When using this product** do not exceed recommended dosage

**Ask a doctor before use if you have**  
• a severe sore throat accompanied by difficulty in breathing or that lasts more than 2 days  
• a sore throat accompanied by fever, headache, rash, swelling, nausea or vomiting

**Stop use and ask a doctor if**  
• sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persists or worsens  
**If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
• adults and children 12 years and over: dissolve 1 lozenge slowly in the mouth. Repeat every 2 hours as needed or as directed by a doctor.  
• children under 12 years: do not use.

**Other information**  
• store at room temperature; protect from moisture • contains soy

**Inactive ingredients**  
corn syrup, flavors, propylene glycol, sucrose

**Questions?** 1-800-283-4879

DIST. BY: PROCTER & GAMBLE,  
CINCINNATI, OH 45202  
Patents: www.pg.com/patents

**P&G**  
www.pg.com  
90173550

239004050 2

**VICKS VAPOCOOL SORE THROAT**

benzocaine and menthol lozenge

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:37000-935
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>BENZOCAINE</b> (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	15 mg
<b>MENTHOL</b> (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	20 mg

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>CORN SYRUP</b> (UNII: 9G5L16BK6N)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	

**Product Characteristics**

<b>Color</b>	yellow	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	19mm
<b>Flavor</b>	MENTHOL	<b>Imprint Code</b>	Vicks;logo
<b>Contains</b>			

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:37000-935-16	2 in 1 CARTON	10/09/2019	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M022	10/09/2019	

**Labeler** - The Procter & Gamble Manufacturing Company (004238200)

Revised: 10/2023

The Procter &amp; Gamble Manufacturing Company