

**PAIN RELIEVER- acetaminophen, ibuprofen tablet, film coated**  
**CVS Pharmacy**

-----  
**CVS Pharmacy, Inc. Pain Reliever Drug Facts**

**Active ingredients (in each caplet)**

Acetaminophen 250 mg

Ibuprofen 125 mg (NSAID\*)

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever

**Uses**

- temporarily relieves minor aches and pains due to:
- headache
- toothache
- backache
- menstrual cramps
- muscular aches
- minor pain of arthritis

**Warnings**

**Acetaminophen liver damage warning:**

This product contains acetaminophen. Severe liver damage may occur if you take:

- with other drugs containing acetaminophen
- more than 6 caplets in 24 hours, which is the maximum daily amount for this product
- 3 or more alcoholic drinks every day while using this product

**Acetaminophen allergy alert:** may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If skin reaction occurs, stop use and seek medical help right away.

**NSAID allergy alert:** ibuprofen may cause a severe allergic reaction, especially in people allergic to

aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**NSAID stomach bleeding warning:**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if

you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you have ever had an allergic reaction to acetaminophen or any other pain reliever
- right before or after heart surgery

**Ask a doctor before use if**

- you have liver disease
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
  - pain gets worse or lasts more than 10 days
  - redness or swelling is present in the painful area
  - any new symptoms appear

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not take more than directed**

**Adults and children 12 years and over: take 2 caplets every 8 hours** while symptoms persist

**Children under 12 years:** ask a doctor

- do not take more than 6 caplets in 24 hours, unless directed by a doctor

### **Other information**

- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

### **Inactive ingredients**

colloidal silicon dioxide, corn starch, croscarmellose sodium, crospovidone, ferric oxide yellow, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, sucralose, talc, titanium dioxide

### **Questions or comments?**

**1-800-719-9260**

### **Package/Label Principal Display Panel**

CVS Health®

Compare to the active ingredients in Advil® Dual Action

Coated Caplets†

Dual Action

Pain Reliever

ACETAMINOPHEN 250 mg & IBUPROFEN (NSAID) 125 mg TABLETS

Pain reliever

Actual Size

CONTAINS 2 MEDICINES

36 CAPLETS†

†Capsule-shaped tablets

Acetaminophen + Ibuprofen

 **CVS Health.**

Compare to the active ingredients in Advil® Dual Action\*\*

Coated Caplets†

NDC 69842-675-68

Dual Action

# Pain Reliever

ACETAMINOPHEN 250 mg &  
IBUPROFEN (NSAID) 125 mg TABLETS

Pain reliever



Actual  
Size

CONTAINS **2** MEDICINES

Acetaminophen

+

Ibuprofen

**36 CAPLETS†**

†Capsule-shaped tablets

53J68 17 C1



#452079

Distributed by: CVS Pharmacy, Inc.  
One CVS Drive, Woonsocket, RI 02895  
© 2024 CVS/pharmacy CVS.com™  
1-800-SHOP CVS V-17160  
100% money back guarantee.  
CVS.com/returnpolicy

Do not use if printed seal under cap is broken or missing.

Do not use with other medicines containing ACETAMINOPHEN; can cause liver damage.

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION.

**Drug Facts**

**Active ingredients (in each caplet) Purposes**

Acetaminophen 250 mg.....Pain reliever  
Ibuprofen 125 mg (NSAID)\*.....Pain reliever  
\*nonsteroidal anti-inflammatory drug

**Uses** ■ temporarily relieves minor aches and pains due to: ■ headache ■ toothache ■ backache ■ menstrual cramps ■ muscular aches ■ minor pain of arthritis

**Warnings**

**Acetaminophen liver damage warning:**

This product contains acetaminophen. Severe liver damage may occur if you take: ■ with other drugs containing acetaminophen ■ more than 6 caplets in 24 hours, which is the maximum daily amount for this product ■ 3 or more alcoholic drinks every day while using this product

**Acetaminophen allergy alert:** may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If skin reaction occurs, stop use and seek medical help right away.

**NSAID allergy alert:** ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

**Drug Facts (continued)**

Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters  
If an allergic reaction occurs, stop use and seek medical help right away.

**NSAID stomach bleeding warning:**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist ■ if you have ever had an allergic reaction to acetaminophen or any other pain reliever ■ right before or after heart surgery

**Ask a doctor before use if** ■ you have liver disease ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are** ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

**Drug Facts (continued)**

**When using this product**

■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding:

■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ do not take more than directed  
Adults and children 12 years and over: take 2 caplets every 8 hours while symptoms persist  
Children under 12 years: ask a doctor ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor

\*\* This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Advil® Dual Action.

**Questions or comments? 1-800-719-9260**

**Inactive ingredients** colloidal silicon dioxide, corn starch, croscarmellose sodium, crospovidone, ferric oxide yellow, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, sucralose, talc, titanium dioxide

**Other information**  
■ read all warnings and directions before use.  
■ keep carton ■ store at 20-25°C (68-77°F)  
■ avoid excessive heat above 40°C (104°F)

**Drug Facts (continued)**

**PAIN RELIEVER**

acetaminophen, ibuprofen tablet, film coated

**Product Information**

**Product Type**

HUMAN OTC DRUG

**Item Code (Source)**

NDC:69842-675

Route of Administration ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	125 mg

### Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

### Product Characteristics

Color	YELLOW	Score	no score
Shape	OVAL	Size	14mm
Flavor		Imprint Code	l;l
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-675-68	1 in 1 CARTON	02/02/2024	
1		36 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:69842-675-23	1 in 1 CARTON	02/02/2024	
2		72 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA214836	02/02/2024	

**Labeler** - CVS Pharmacy (062312574)

Revised: 2/2024

CVS Pharmacy