

**MENSTRUAL COMPLETE MAXIMUM STRENGTH- acetaminophen, caffeine,  
pyrilamine maleate tablet, film coated  
CHAIN DRUG MARKETING ASSOCIATION INC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Quality Choice 44-390**

***Active ingredients (in each caplet)***

Acetaminophen 500 mg  
Caffeine 60 mg  
Pyrilamine maleate 15 mg

***Purpose***

Pain reliever  
Diuretic  
Antihistamine

***Uses***

for the temporary relief of these symptoms associated with menstrual periods:

- bloating
- headache
- muscle aches
- cramps
- backache
- fatigue
- water-weight gain

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- difficulty in urination due to enlargement of the prostate gland
- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **When using this product**

- avoid alcoholic beverages
- excitability may occur, especially in children
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- you may get drowsy
- limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat. The recommended dose of this product contains about as much caffeine as a cup of coffee.

### **Stop use and ask a doctor if**

- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not take more than the recommended dose**

- adults and children 12 years and over:
  - take 2 caplets with water
  - repeat every 6 hours, as needed
  - do not exceed 6 caplets per day
- children under 12 years: ask a doctor

### **Other information**

- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)
- see end flap for expiration date and lot number

### **Inactive ingredients**

corn starch, croscarmellose sodium, crospovidone, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, povidone, silicon dioxide, stearic acid, titanium dioxide, triacetin

### **Questions or comments?**

**1-800-426-9391**

### **Principal Display Panel**

**QC  
QUALITY®  
CHOICE**

**NDC 63868-032-24**

**\*Compare to the active ingredients  
in MIDOL® COMPLETE**

Maximum Strength  
**Menstrual Complete**

**Acetaminophen** 500 mg, Caffeine 60 mg  
Pyrilamine maleate 15 mg  
Pain Reliever / Diuretic / Antihistamine

Multi-Symptom Relief

**Actual  
Size**

**24** Caplets

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

\*This product is not manufactured or distributed by  
Bayer HealthCare LLC, owner of the registered  
trademark Midol® Complete.  
50844 REV0517A39008

**100% QC**  
SATISFACTION  
GUARANTEED

Distributed by C.D.M.A., Inc. ©  
43157 W 9 Mile Rd  
Novi, MI 48375  
[www.qualitychoice.com](http://www.qualitychoice.com)  
Questions: 248-449-9300

No Print / Varnish Area  
Lot no. & Exp. date



\* This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Midol® Complete.  
50844 REV0517A39008

B-0220-390-08  
REV0517A39008

**Drug Facts** (continued)

**Warnings**: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours

■ with other drugs containing acetaminophen  
■ 3 or more alcoholic drinks every day while using this product

**Allergy alert**: Acetaminophen may cause severe skin reactions. Symptoms may include:  
■ skin redness ■ blisters ■ rash

**Drug Facts**

**Active ingredients** (in each caplet)  
Acetaminophen 500 mg ..... Pain reliever  
Caffeine 60 mg ..... Diuretic  
Pyrilamine maleate 15 mg ..... Antihistamine

**Purpose**

**Uses**  
for the temporary relief of these symptoms associated with menstrual periods:  
■ bloating ■ headache ■ muscle aches  
■ cramps ■ backache ■ fatigue  
■ water-weight gain

**KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION**

NDC 63868-032-24

Compare to the active ingredients in MIDOL® COMPLETE

**QC QUALITY CHOICE**

**Maximum Strength Menstrual Complete**

**Acetaminophen 500 mg, Caffeine 60 mg  
Pyrilamine maleate 15 mg**

Pain Reliever / Diuretic / Antihistamine  
Multi-Symptom Relief

**24 Caplets** Actual Size

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING



**Drug Facts** (continued)

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
■ do not take more than the recommended dose  
■ adults and children 12 years and over:  
■ take 2 caplets with water  
■ repeat every 6 hours, as needed  
■ do not exceed 6 caplets per day  
■ children under 12 years: ask a doctor

**Other information**  
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)  
■ see end flap for expiration date and lot number

**Inactive ingredients**: corn starch, croscarmellose sodium, crospovidone, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, povidone, silicon dioxide, stearic acid, titanium dioxide, triacetin

**Questions or comments?**  
1-800-426-9391

**Drug Facts** (continued)

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**  
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**  
■ difficulty in urination due to enlargement of the prostate gland ■ liver disease  
■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma

**Ask a doctor or pharmacist before use if you are**  
■ taking the blood thinning drug warfarin  
■ taking sedatives or tranquilizers

**When using this product** ■ you may get drowsy  
■ avoid alcoholic beverages

**Drug Facts** (continued)

■ alcohol, sedatives, and tranquilizers may increase drowsiness  
■ excitability may occur, especially in children  
■ be careful when driving a motor vehicle or operating machinery  
■ limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat. The recommended dose of this product contains about as much caffeine as a cup of coffee.

**Stop use and ask a doctor if**  
■ new symptoms occur  
■ redness or swelling is present  
■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days

If pregnant or breast-feeding, ask a health professional before use.

Quality Choice 44-390

**MENSTRUAL COMPLETE MAXIMUM STRENGTH**  
acetaminophen, caffeine, pyrilamine maleate tablet, film coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63868-032
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
<b>CAFFEINE</b> (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	60 mg
<b>PYRILAMINE MALEATE</b> (UNII: R35D29L3ZA) (PYRILAMINE - UNII:HPE317O9TL)	PYRILAMINE MALEATE	15 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>CROSPVIDONE</b> (UNII: 2S7830E561)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6130)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	44;390
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:63868-032-24	1 in 1 CARTON	04/29/2002	
1		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph not			

OTC monograph not final	part343	04/29/2002	
-------------------------	---------	------------	--

**Labeler - CHAIN DRUG MARKETING ASSOCIATION INC (011920774)**

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(63868-032)

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(63868-032)

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	pack(63868-032)

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	pack(63868-032)

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(63868-032)

Revised: 10/2021

CHAIN DRUG MARKETING ASSOCIATION INC