

TOPCARE STOMACH RELIEF - bismuth subsalicylate liquid

Topco Associates, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Topcare Regular Strength Stomach Relief Liquid

ACTIVE INGREDIENT(in each 30 mL)

Bismuth subsalicylate 525 mg

PURPOSE

Upset stomach reliever and anti-diarrheal

USE(S)

relieves:

- diarrhea
- heartburn
- indigestion
- nausea
- upset stomach associated with these symptoms

WARNINGS

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

DO NOT USE IF YOU HAVE

- an ulcer
- bloody or black stool

- a bleeding problem

ASK A DOCTOR BEFORE USE IF YOU HAVE

- fever
- mucus in the stool

ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE

taking any drug for

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

WHEN USING THIS PRODUCT

a temporary, but harmless darkening of the stool and/or tongue may occur

STOP USE AND ASK DOCTOR IF

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

IF PREGNANT OR BREAST-FEEDING,

ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN

In case of overdose, get medical help or contact a Poison Control Center immediately.

DIRECTIONS

- **shake well before use**
- mL = milliliter
- TBSP = tablespoon
- adults and children 12 years and over: 1 dose (2 TBSP or 30 mL) every 1/2 to 1 hour as needed
- do not exceed 8 doses (16 TBSP or 240 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of fluids to help prevent dehydration caused by diarrhea

OTHER INFORMATION

- **each 30 mL or 2 TBSP contains:**
- potassium 25 mg
- salicylate 260 mg
- sodium 8 mg
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)
- dosage cup provided

INACTIVE INGREDIENTS

benzoic acid, D&C red # 22, D&C red # 28, flavor, hydroxyethyl cellulose, potassium hydroxide, purified water, saccharin sodium, salicylic acid, simethicone, xanthan gum

PRINCIPAL DISPLAY PANEL

NDC 76162-023-05

COMPARE TO PEPTO-BISMOL® REGULAR STRENGTH ACTIVE INGREDIENT*

TopCare

REGULAR STRENGTH

Stomach Relief

BISMUTH SUBSALICYLATE 525 mg per 30 mL

UPSET STOMACH RELIEVER/ANTIDIARRHEAL

5 SYMPTOM DIGESTIVE RELIEF

- Heartburn
- Indigestion
- Nausea
- Upset stomach
- Diarrhea

12 FL OZ (354 mL)

Drug Facts

Active ingredient (in each 30 mL) Bismuth subsalicylate 525 mg Purpose Upset stomach reliever and anti-diarrheal	DO NOT USE IF IMPRINTED SHRINKBAND IS MISSING OR BROKEN. Uses relieves: ■ diarrhea ■ heartburn ■ indigestion ■ nausea ■ upset stomach associated with these symptoms Warnings <i>Réye's syndrome:</i> Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Réye's syndrome, a rare but serious illness. Allergy alert: Contains salicylate. Do not take if you are allergic to salicylates (including aspirin) taking other salicylate products Do not use if you have ■ bloody or black stool ■ an ulcer ■ a bleeding problem Ask a doctor before use if you have ■ fever ■ mucus in the stool Ask a doctor or pharmacist before use if you are taking any drug or ■ anticoagulation (thinning of the blood) ■ diabetes ■ gout ■ arthritis When using this product a temporary, but harmless darkening of the stool and/or tongue may occur. Stop use and ask a doctor if ■ symptoms get worse ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center immediately.
--	--

NDC 76162-023-05

TOPCARE STOMACH RELIEF

bismuth subsalicylate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:76162-023
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
BENZOIC ACID (UNII: 8SKN0B0MIM)	
D&C RED NO. 22 (UNII: 1678RKX8RT)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SALICYLIC ACID (UNII: O414PZ4LPZ)	
HYDROXYETHYL CELLULOSE (1500 MPA.S AT 1%) (UNII: L605B5892V)	
DIMETHICONE (UNII: 92RU3N3Y1O)	

XANTHAN GUM (UNII: TTV12P4NEE)

Product Characteristics

Color	PINK	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76162-023-05	354 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/05/2022	
2	NDC:76162-023-04	236 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/05/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part335	01/05/2021	

Labeler - Topco Associates, LLC (006935977)

Revised: 11/2022

Topco Associates, LLC