

COLD AND FLU NIGHTTIME SEVERE- acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl tablet, film coated
L.N.K. International, Inc.

Convenience Valet 44-677

Active ingredients (in each caplet)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Doxylamine succinate 6.25 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Antihistamine
Nasal decongestant

Uses

- temporarily relieves common cold and flu symptoms:
 - minor aches and pains
 - sinus congestion and pressure
 - sore throat
 - fever
 - headache
 - nasal congestion
 - cough due to minor throat and bronchial irritation
 - cough to help you sleep
 - runny nose and sneezing
- reduces swelling of nasal passages
- promotes nasal and/or sinus drainage
- temporarily restores freer breathing through the nose

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters

- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- diabetes
- thyroid disease
- heart disease
- glaucoma
- high blood pressure
- difficulty in urination due to enlargement of the prostate gland
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic beverages
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- do not take more than 8 caplets in 24 hours
- adults and children 12 years and over: take 2 caplets with water every 4 hours
- children under 12 years: ask a doctor

Other information

- **each caplet contains:** sodium 3 mg
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

black iron oxide, corn starch, crospovidone, D&C yellow #10 aluminum lake, FD&C blue #1 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments?

1-877-507-6516

Principal display panel

24/7 life

BY 7-ELEVEN™

Cold & Flu

Nighttime Severe

Acetaminophen / Pain Reliever/Fever Reducer
Dextromethorphan HBr /Cough Suppressant
Doxylamine Succinate / Antihistamine

Phenylephrine HCl / Nasal Decongestant

Relieves: Headache, Fever, Sore Throat,
Minor Aches & Pains, Nasal Congestion,
Sinus Pressure, Sneezing, Runny Nose,
Cough

compare to Vicks® NyQuil®
VapoCOOL™ SEVERE Cold &
Flu + Congestion active ingredients*

Actual Size

16
CAPLETS

Satisfaction Guaranteed 1-800-255-0711

Distributed by: 7-Eleven, Inc.
Irving, TX 75063 www.7-ELEVEN.com

*This product is not manufactured or distributed by
The Procter & Gamble Company. Vicks®, NyQuil®, and
VapoCOOL™ are trademarks of The Procter & Gamble Company.
50844 REV0722B67721

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
OPENED OR IF BLISTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING**

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Drug Facts (continued)

Active ingredients (in each caplet)

Acetaminophen 325 mg. Pain reliever/fever reducer
 Dextromethorphan HBr 10 mg. Cough suppressant
 Doxylamine succinate 6.25 mg. Antihistamine
 Phenylephrine HCl 5 mg. Nasal decongestant

Uses ■ temporarily relieves common cold and flu symptoms:
 ■ minor aches and pains ■ sinus congestion and pressure
 ■ sore throat ■ fever ■ headache ■ nasal congestion
 ■ cough due to minor throat and bronchial irritation

Drug Facts (continued)

Caution: ■ temporarily relieves common cold and flu symptoms:
 ■ reduces swelling of nasal passages
 ■ promotes nasal and/or sinus drainage

No Print/No Varnish
 Lot # & Exp. Area

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
 OPENED OR IF BLISTER UNIT IS TORN, BROKEN
 OR SHOWS ANY SIGNS OF TAMPERING



Cold & Flu Nighttime Severe

Acetaminophen / Pain Reliever/Fever Reducer
Dextromethorphan HBr /Cough Suppressant
Doxylamine Succinate / Antihistamine
Phenylephrine HCl / Nasal Decongestant

Relieves: Headache, Fever, Sore Throat,
 Minor Aches & Pains, Nasal Congestion,
 Sinus Pressure, Sneezing, Runny Nose,
 Cough

compare to **Vicks® NyQuil®**
VapoCOOL™ SEVERE Cold &
Flu + Congestion active ingredients*

16
 CAPLETS

Actual Size



PARENTS:
 Learn about teen medicine abuse
www.StopMedicineAbuse.org

Satisfaction Guaranteed 1-800-255-0711

Distributed by: 7-Eleven, Inc.
 Irving, TX 75063 www.7-ELEVEN.com

*This product is not manufactured or distributed by
 The Procter & Gamble Company. Vicks®, NyQuil® and
 VapoCOOL™ are trademarks of The Procter & Gamble Company.
 65168C-SE-10-22-1 50844 REV0722867721

Drug Facts (continued)

Other information ■ each caplet contains: sodium 3 mg
 ■ TAMPER EVIDENT; DO NOT USE IF OUTER PACKAGE IS OPENED OR
 ■ BLISTER IS TORN OR BROKEN
 ■ store at 25°C (77°F); excursions permitted between 15°-30°C
 (59°-86°F) ■ see end flap for expiration date and lot number

Inactive ingredients black iron oxide, corn starch,
 (59°-86°F) ■ see end flap for expiration date and lot number

Questions or comments? 1-877-507-6516

Drug Facts (continued)

Warnings ■ temporarily restores free breathing through the nose
 ■ cough that occurs with too much phlegm (mucus)
 ■ Ask a doctor or pharmacist before use if you are
 ■ taking sedatives or tranquilizers
 ■ taking the blood thinning drug warfarin
 ■ When using this product
 ■ do not exceed recommended dosage
 ■ excitability may occur, especially in children
 ■ marked drowsiness may occur ■ avoid alcoholic beverages
 ■ be careful when driving a motor vehicle or operating machinery
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness
 ■ nervousness, dizziness, or sleeplessness occur
 ■ pain, nasal congestion, or cough gets worse or lasts more than 7
 ■ days
 ■ fever gets worse or lasts more than 3 days
 ■ new symptoms occur
 ■ cough comes back or occurs with rash or headache that lasts. These
 ■ could be signs of a serious condition.
 ■ If pregnant or breast-feeding, ask a health professional before use.
 ■ Keep out of reach of children. In case of overdose, get medical help or
 ■ contact a Poison Control Center right away. Prompt medical attention
 ■ is critical for adults as well as for children even if you do not notice any
 ■ signs or symptoms.

Directions
 ■ adults and children 12 years and over: take 2 caplets with water
 ■ do not take more than 8 caplets in 24 hours
 ■ do not take more than directed
 ■ children under 12 years: ask a doctor
 ■ every 4 hours

B-1205-677-21
 REV0722867721

Convenience Valet 44-677

COLD AND FLU NIGHTTIME SEVERE

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50844-677
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
FD&C YELLOW NO. 6 ALUMINUM LAKE (UNII: GYP6Z2JR6Q)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	green	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	44;677
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-677-21	2 in 1 CARTON	02/07/2022	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	02/07/2022	

Labeler - L.N.K. International, Inc. (038154464)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(50844-677) , pack(50844-677)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(50844-677)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(50844-677)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(50844-677)

Revised: 6/2024

L.N.K. International, Inc.