

BAYER GENUINE ASPIRIN- aspirin tablet
JC World Bell Wholesale Co., Inc.

BAYER 2'S BLISTER PACK

Drug Facts

Active ingredients (in each tablet)

Aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains due to:

- headache • muscle pain
- toothache • menstrual pain
- colds • minor pain of arthritis
- temporarily reduces fever

Warnings

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Reye's syndrome:**

Aspirin may cause a severe allergic reaction which may include: **Allergy alert:**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you **Stomach bleeding warning:**

- hives
- facial swelling
- asthma (wheezing)
- shock

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for • gout • diabetes • arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding,

ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
- children under 12 years: consult a doctor

Other information

- save carton for full directions and warnings
- store at room temperature


Inactive ingredients

corn starch, hypromellose, powdered cellulose, triacetin

Questions or comments?

1-800-331-4536 (Mon - Fri 9AM - 5PM EST)

Package Labeling:




<p>BAYER ASPIRIN (ACETASALIC ACID) PAIN RELIEVER/FEVER REDUCER</p>	
<p>Drug Facts</p> <p>Active ingredient (in each tablet) Aspirin 325 mg (NSAID) ... *Nonsteroidal anti-inflammatory drug</p> <p>Uses</p> <ul style="list-style-type: none"> temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> headache muscle pain menstrual pain colds temporarily reduces fever <p>Warnings</p> <p>People's symptoms: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Aspirin may cause a severe allergic reaction which may include:</p> <ul style="list-style-type: none"> hives facial swelling asthma (wheezing) shock <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:</p> <ul style="list-style-type: none"> are age 60 or older have had stomach ulcers or bleeding problems take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed <p>Do not use</p> <ul style="list-style-type: none"> if you are allergic to aspirin or any other pain reliever/fever reducer if you have ever had an allergic reaction to this product or any of its ingredients <p>Ask a doctor before use if</p> <ul style="list-style-type: none"> stomach bleeding warning applies to you you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, or kidney disease you are taking a diuretic you have asthma <p>Ask a doctor or pharmacist before use if you are taking a prescription drug for:</p> <ul style="list-style-type: none"> gold diabetes arthritis <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> an allergic reaction occurs. Seek medical help right away. 	<p>Purposes</p> <p>Pain reliever/fever reducer</p> <ul style="list-style-type: none"> fever stomach pain that does not get better pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur ringing in the ears or a loss of hearing occurs <p>Directions</p> <ul style="list-style-type: none"> drink a full glass of water with each dose adults and children 15 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours children under 12: see a doctor <p>Other information</p> <ul style="list-style-type: none"> use caution for full directions and warnings Store at room temperature <p>Inactive ingredients corn starch, hypromellose, powdered cellulose, triacetin</p> <p>Questions or comments? 1-800-331-4536 (Mon - Fri 9AM - 5PM EST)</p> <p>Do not use if printed safety seal bearing "Bayer HealthCare" under cap is torn or missing.</p>
<p>LOT: _____</p> <p>EXP: _____</p> <p>This product is repackaged by J.C. World Bell Wholesale Co, Inc; 376-380 Railroad Ave, Hackensack NJ 07601; 201-488-1096 (Mon-Fri 9AM-6PM EST), from a product manufactured by Bayer.</p>	

BACK

FRONT



FRONT CARD



BAYER GENUINE ASPIRIN

aspirin tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50269-005(NDC:0280-2000)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	

POWDERED CELLULOSE (UNII: SMD1X3XO9M)

TRIACETIN (UNII: XHX3C3X673)

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	BAYER
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50269-005-06	6 in 1 BOX	10/30/2019	
1		2 in 1 BLISTER PACK		
1		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	10/30/2019	

Labeler - JC World Bell Wholesale Co., Inc. (805257581)

Revised: 11/2023

JC World Bell Wholesale Co., Inc.