

**ANTI-DIARRHEAL- loperamide hcl tablet**  
**Discount Drug Mart**

-----  
**Drug Mart 44-375**

***Active ingredient (in each caplet)***

Loperamide HCl 2 mg

***Purpose***

Anti-diarrheal

***Use***

controls symptoms of diarrhea, including Travelers' Diarrhea

***Warnings***

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.

**Heart alert:** Taking more than directed can cause serious heart problems or death.

**Do not use**

if you have bloody or black stool.

**Ask a doctor before use if you have**

- a fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product,**

tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise; use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34-47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

**Other information**

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store between 20°-25°C (68°-77°F)
- see end flap for expiration date and lot number

**Inactive ingredients**

corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide

**Questions or comments?**

Call **1-800-426-9391** 8:30 AM-4:00 PM ET, Monday-Friday

**Principal Display Panel**

DISCOUNT  
drug

**mart**  
FOOD MARKET  
AN EMPLOYEE OWNERSHIP COMPANY

NDC 53943-375-08

MAY CONTAIN  
**ANTI-THEFT**  
DEVICE

**Anti-Diarrheal**  
**Loperamide HCl Tablets, 2 mg**  
**Anti-Diarrheal**

Controls The Symptoms of Diarrhea,  
Including Travelers' Diarrhea

\*Compare to the active ingredient in IMODIUM<sup>®</sup> A-D

**24** Caplets          actual size  
Each caplet contains 2 mg loperamide hydrochloride

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS  
TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

\*This product is not manufactured or distributed by Johnson & Johnson Corporation,  
owner of the registered trademark Imodium<sup>®</sup> A-D.

50844    REV0619A37508

DISTRIBUTED BY: DRUG MART - FOOD FAIR  
211 COMMERCE DRIVE, MEDINA, OH 44256  
[www.discount-drugmart.com](http://www.discount-drugmart.com)

---

**SATISFACTION GUARANTEED**

IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE  
TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO  
STORE, SEND REASON FOR DISSATISFACTION, NAME,  
ADDRESS AND EMPTY PACKAGE TO:

DISCOUNT DRUG MART  
211 COMMERCE DRIVE, MEDINA, OHIO 44256

---

1. To open, tear at perforations and remove a section.
2. Use ✂ to cut through plastic for caplet.

This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Includium™ A-D.

**Questions or comments?** Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

**Inactive ingredients** corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide

**Drug Facts (continued)** ■ store between 20°-25° C (68°-77° F) ■ see end flap for expiration date and lot number

**Other information**

■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

children under 2 years (up to 33 lbs)	do not use
children 2-5 years (34-47 lbs)	ask a doctor
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours

**Directions** ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

■ find right dose on chart. If possible, use weight to dose; otherwise, use age.  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.  
If pregnant or breast-feeding, ask a health professional before use.  
■ you get abdominal swelling or bloating. These may be signs of a serious condition.  
Stop use and ask a doctor if ■ symptoms get worse ■ diarrhea lasts for more than 2 days

**Warnings**  
Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.  
Heart alert: Taking more than directed can cause serious heart problems or death.  
Do not use if you have bloody or black stool.  
Ask a doctor before use if you have ■ a fever ■ mucus in the stool ■ a history of liver disease ■ a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.  
When using this product, tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

Use controls symptoms of diarrhea, including Traveler's Diarrhea

**Purpose** Loperamide HCl 2 mg ..... Anti-diarrheal

**Active ingredient (in each caplet)** Loperamide HCl 2 mg ..... Anti-diarrheal

**Drug Facts** KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION



DISTRIBUTED BY: DRUG MART - FOOD FAIR  
211 COMMERCE DRIVE, MEDINA, OH 44256  
www.discount-drugmart.com

**SATISFACTION GUARANTEED**  
IF DISSATISFIED, RETURN UNOPENED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO:  
DISCOUNT DRUG MART  
211 COMMERCE DRIVE, MEDINA, OH 44256

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNITS IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

50844 REV0619A37508

No print/No varnish  
Lot & Exp date



**Anti-Diarrheal**  
Loperamide HCl Tablets, 2 mg  
Anti-Diarrheal


B-0380-375-08  
REV0619A37508



MAY CONTAIN  
**ANTI-THEFT**  
DEVICE

NDC 53943-375-08

**Anti-Diarrheal**  
Loperamide HCl Tablets, 2 mg



**Drug Mart 44-375**

## ANTI-DIARRHEAL

loperamide hcl tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:53943-375
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>LOPERAMIDE HYDROCHLORIDE</b> (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>D&amp;C YELLOW NO. 10 ALUMINUM LAKE</b> (UNII: CQ3XH3DET6)	
<b>DIBASIC CALCIUM PHOSPHATE DIHYDRATE</b> (UNII: O7TSZ97GEP)	
<b>FD&amp;C BLUE NO. 1 ALUMINUM LAKE</b> (UNII: J9EQA3S2JM)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	

### Product Characteristics

<b>Color</b>	green	<b>Score</b>	2 pieces
<b>Shape</b>	OVAL	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;375
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53943-375-02	2 in 1 CARTON	12/13/2019	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:53943-375-08	4 in 1 CARTON	12/13/2019	
2		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076497	12/13/2019	

**Labeler** - Discount Drug Mart (047741335)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(53943-375) , pack(53943-375)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(53943-375)

Revised: 6/2023

Discount Drug Mart