# LEADER STOMACH RELIEF- bismuth subsalicylate tablet, chewable CARDINAL HEALTH 110, LLC. DBA LEADER

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

#### **Leader Stomach Relief**

## **ACTIVE INGREDIENT(in each tablet)**

Bismuth subsalicylate 262 mg (total salicylate 102 mg per tablet)

#### **PURPOSE**

Upset stomach reliever and anti-diarrheal

## USE(S)

relieves:

- diarrhea
- heartburn
- indigestion
- nausea
- upset stomach associated with these symptoms

#### WARNINGS

**Reye's Syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behaviour with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

#### DO NOT USE IF YOU HAVE

- bloody or black stool
- an ulcer

• a bleeding problem

#### **ASK A DOCTOR BEFORE USE IF YOU HAVE**

- fever
- mucus in the stool

#### ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE

taking any drug for

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

#### WHEN USING THIS PRODUCT

a temporary, but harmless darkening of the stool and/or tongue may occur

#### STOP USE AND ASK DOCTOR IF

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

### IF PREGNANT OR BREAST-FEEDING,

ask a health professional before use.

#### **KEEP OUT OF REACH OF CHILDREN**

In case of overdose, get medical help or contact a Poison Control Center right away.

#### **DIRECTIONS**

- chew or dissolve in mouth
- adults and children 12 years and over: 2 tablets every 1/2 to 1 hour as needed
- do not take more than 8 doses (16 tablets) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of fluids to help prevent dehydration which may accompany diarrhea

#### OTHER INFORMATION

- each tablet contains:
- sodium less than 1 mg
- salicylate 102 mg

- very low sodium
- avoid excessive heat (over 104°F or 40°C)
- TAMPER EVIDENT: Do not use if individual compartments are torn or open

#### **INACTIVE INGREDIENTS**

calcium carbonate, D&C red 27 aluminum lake, flavor, magnesium stearate, mannitol, pregelatinized starch, saccharin sodium

#### PRINCIPAL DISPLAY PANEL

NDC 70000-0433-2 **LEADER**™

## Stomach Relief

Bismuth Subsalicylate 262 mg Upset Stomach Reliever/Antidiarrheal

5 Symptom Digestive Relief

- Diarrhea
- Heartburn
- Indigestion
- Nausea
- Upset stomach

COMPATE TO PEPTO-BISMOL® active ingredient\*

**48 CHEWABLE TABLETS** 



## **LEADER STOMACH RELIEF**

bismuth subsalicylate tablet, chewable

| Product Information     |                |                    |                |
|-------------------------|----------------|--------------------|----------------|
| Product Type            | HUMAN OTC DRUG | Item Code (Source) | NDC:70000-0433 |
| Route of Administration | ORAL           |                    |                |
|                         |                |                    |                |

| Active Ingredient/Active Moiety   |                          |          |  |  |
|---|--------------------------|----------|--|--|
| Ingredient Name   | <b>Basis of Strength</b> | Strength |  |  |
| BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII: 0414PZ 4LPZ) | BISMUTH<br>SUBSALICYLATE | 262 mg   |  |  |

| Inactive Ingredients                         |          |  |
|--|----------|--|
| Ingredient Name                              | Strength |  |
| CALCIUM CARBONATE (UNII: H0G9379FGK)         |          |  |
| <b>D&amp;C RED NO. 27</b> (UNII: 2LRS185U6K) |          |  |

| MAGNESIUM STEARATE (UNII: 70097M6I30) |  |
|---------------------------------------|--|
| MANNITOL (UNII: 30WL53L36A)           |  |
| STARCH, CORN (UNII: O8232NY3SJ)       |  |
| SACCHARIN SODIUM (UNII: SB8ZUX40TY)   |  |

| Product Characteristics |       |              |          |
|-------------------------|-------|--------------|----------|
| Color                   | PINK  | Score        | no score |
| Shape                   | ROUND | Size         | 17mm     |
| Flavor                  |       | Imprint Code | GDC122   |
| Contains                |       |              |          |

| Packaging |                      |  |                         |                       |
|-----------|----------------------|--|-------------------------|-----------------------|
| #         | Item Code            | Package Description                                    | Marketing Start<br>Date | Marketing End<br>Date |
| 1         | NDC:70000-<br>0433-2 | 8 in 1 BOX   | 06/06/2019              |                       |
| 1         |                      | 6 in 1 BLISTER PACK; Type 0: Not a Combination Product |                         |                       |
| 2         | NDC:70000-<br>0433-1 | 5 in 1 BOX   | 06/06/2019              |                       |
| 2         |                      | 6 in 1 BLISTER PACK; Type 0: Not a Combination Product |                         |                       |

| Marketing Information  |   |                         |                       |
|------------------------|---|-------------------------|-----------------------|
| Marketing<br>Category  | Application Number or Monograph<br>Citation | Marketing Start<br>Date | Marketing End<br>Date |
| OTC MONOGRAPH<br>FINAL | part335                                     | 06/06/2019              |                       |
|                        |   |                         |                       |

## Labeler - CARDINAL HEALTH 110, LLC. DBA LEADER (063997360)

Revised: 1/2022 CARDINAL HEALTH 110, LLC. DBA LEADER