

**SAFCO SENSICAINA ULTRA TOPICAL ANESTHETIC GEL- benzocaine gel**  
**Safco Dental Supply Co.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Active Ingredients**

Benzocaine 200mg (in each g)

**Purpose**

Oral Anesthetic

**Use**

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

**Warnings**

**Methemoglobinemia warning:** Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

**Allergy Alert:** Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

**Do not use**

- for more than 7 days unless directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.
- for teething
- in children under 2 years of age

**When using this product** Avoid contact with eyes. If contact occurs, flush with water.

**Do not exceed recommended dosage.** If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

**If pregnant or breast feeding, ask a physician before use.**

**Keep out of reach of children.**

**Directions**

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

**Other Information**


Store at 59°-86°F (15°-30°C). Protect from freezing.

**Inactive ingredients**

flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.

**Questions or comments?**

safcodental.com \* 1-800-621-2178

<p>Distributed by:  <b>Safco Dental Supply Co.</b>          Buffalo Grove, IL 60089 • 800-621-2178          www.safcodental.com Made in USA</p>		<p><b>SensiCaine™ Ultra</b>          Topical Anesthetic Gel</p>
<p>For professional use only</p>	<p><b>Bubble Gum</b>  <b>REF 102-0457</b>          NDC 67239-0218-1</p>	<p>1 oz. (29.6 ml)  <b>Contains 20% Benzocaine</b></p>
<p>OPEN HERE </p>	<p>Gluten Free</p>	

<p>directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.</p> <ul style="list-style-type: none"> <li>• for teething</li> <li>• in children under 2 years of age</li> </ul>	<p><b>When using this product</b>          Avoid contact with eyes. If contact occurs, flush with water.</p> <p><b>Do not exceed recommended dosage.</b> If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.</p> <p><b>If pregnant or breast feeding,</b> ask a physician before use.</p> <p><b>Keep out of reach of children.</b></p>	<p><b>Directions</b></p> <ul style="list-style-type: none"> <li>• Apply only amount needed to the oral mucosa to prevent or relieve pain.</li> <li>• children under 2 years of age: do not use</li> </ul> <p><b>Other information</b>          Store at 59° - 86° F (15 - 30° C). Protect from freezing.</p> <p><b>Inactive ingredients</b>          flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.</p> <p><b>Questions or comments?</b>          safcodental.com • 800-621-2178</p> <p style="text-align: right;">99091778 Rev 03/2019</p>
--	--	--

<p><b>Drug Facts</b></p> <p><b>Active ingredients</b>          Benzocaine 200 mg (in each g)</p> <p><b>Purpose</b>          Oral Anesthetic</p> <p><b>Use</b>          For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.</p>	<p><b>Warnings</b></p> <p><b>Methemoglobinemia warning:</b> Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:</p> <ul style="list-style-type: none"> <li>• pale, gray or blue colored skin (cyanosis)</li> <li>• headache</li> <li>• rapid heart rate</li> <li>• shortness of breath</li> <li>• dizziness or lightheadedness</li> <li>• fatigue or lack of energy</li> </ul>	<p><b>Allergy Alert:</b> Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.</p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>• for more than 7 days unless</li> </ul>
--	--	---

Distributed by:  
**Safco Dental Supply Co.**  
 Buffalo Grove, IL 60089 • 800-621-2178  
 www.safcodental.com Made in USA

For professional use only

OPEN  
 HERE

Bubble Gum  
**REF 102-0457**  
 NDC 67239-0218-1  
 Gluten Free

**safco**

**SensiCaine™ Ultra**  
 Topical Anesthetic Gel

1 oz. (29.6 ml)  
 Contains 20% Benzocaine

directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.

- for teething
- in children under 2 years of age

**When using this product**

Avoid contact with eyes. If contact occurs, flush with water.

**Do not exceed recommended dosage.**

If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

**If pregnant or breast feeding,** ask a physician before use.

**Keep out of reach of children.**

**Directions**

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

**Other information**

Store at 59° - 86° F (15 - 30° C). Protect from freezing.

**Inactive ingredients**

flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.

**Questions or comments?**

safcodental.com • 800-621-2178

99081778 Rev 03/2019

**Drug Facts**

**Active ingredients**

Benzocaine 200 mg (in each g)

**Purpose**

Oral Anesthetic

**Use**

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

**Warnings**

**Methemoglobinemia warning:** Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

**Allergy Alert:** Do not use on

patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

**Do not use**

- for more than 7 days unless

Distributed by:  
**Safco Dental Supply Co.**  
 Buffalo Grove, IL 60089 • 800-621-2178  
 www.safcodental.com Made in USA

**safco**

Bubble Gum  
**REF 102-0457**  
 NDC 67239-0218-1  
 Gluten Free

**SensiCaine™ Ultra**  
 Topical Anesthetic Gel

1 oz. (29.6 ml)  
 Contains 20% Benzocaine

For professional use only



directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.

- for teething
- in children under 2 years of age

**When using this product**  
 Avoid contact with eyes. If contact occurs, flush with water.

**Do not exceed recommended dosage.** If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

**If pregnant or breast feeding,** ask a physician before use.

**Keep out of reach of children.**

**Directions**

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

**Other information**  
 Store at 59° - 86° F (15 - 30° C). Protect from freezing.

**Inactive ingredients**  
 flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #78, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.

**Questions or comments?**  
 safcodental.com • 800-621-2178

99081778 Rev 03/2019

**Drug Facts**

**Active ingredients**  
 Benzocaine 200 mg (in each g)

**Purpose**  
 Oral Anesthetic

**Use**  
 For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

**Warnings**

**Methemoglobinemia warning:** Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

**Allergy Alert:** Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

**Do not use**

- for more than 7 days unless

## SAFCO SENSICAIN ULTRA TOPICAL ANESTHETIC GEL

benzocaine gel

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:67239-0218
<b>Route of Administration</b>	DENTAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BENZOCAINE</b> (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g

### Inactive Ingredients

Ingredient Name	Strength
<b>POLYETHYLENE GLYCOL 3350</b> (UNII: G2M7P15E5P)	
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>FD&amp;C YELLOW NO. 5</b> (UNII: I753WB2F1M)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>FD&amp;C RED NO. 3</b> (UNII: PN2ZH5LOQY)	

FD&C RED NO. 40 (UNII: WZB9127XOA)

FD&C BLUE NO. 1 (UNII: H3R47K3TBD)

FD&C GREEN NO. 3 (UNII: 3P3ONR601S)

D&C RED NO. 28 (UNII: 767IP0Y5NH)

D&C GREEN NO. 5 (UNII: 8J6RDU8L9X)

### Product Characteristics

Color	pink	Score	
Shape		Size	
Flavor	BUBBLE GUM	Imprint Code	
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67239-0218-1	30 g in 1 JAR; Type 0: Not a Combination Product	03/05/2019	

Distributed by:  
**Safco Dental Supply Co.**  
 Buffalo Grove, IL 60089 • 800-621-2178  
 www.safcodental.com Made in USA



For professional use only

Bubble Gum

**REF** 102-0457

NDC 67239-0218-1

Gluten Free

**SensiCaine™ Ultra**  
 Topical Anesthetic Gel

1 oz. (29.6 ml)

Contains 20% Benzocaine



directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.

- for teething
- in children under 2 years of age

**When using this product**  
 Avoid contact with eyes. If contact occurs, flush with water.

**Do not exceed recommended dosage.** If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

**If pregnant or breast feeding,** ask a physician before use.

**Keep out of reach of children.**

**Directions**

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

**Other information**  
 Store at 59° - 86° F (15 - 30° C). Protect from freezing.

**Inactive ingredients**  
 flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.

**Questions or comments?**  
 safcodental.com • 800-621-2178

99091778 Rev 03/2019

<b>Drug Facts</b>
<b>Active ingredients</b> Benzocaine 200 mg (in each g)
<b>Purpose</b> Oral Anesthetic
<b>Use</b> For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.
<b>Warnings</b> <b>Methemoglobinemia</b> <b>warning:</b> Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: <ul style="list-style-type: none"> <li>• pale, gray or blue colored skin (cyanosis)</li> <li>• headache</li> <li>• rapid heart rate</li> <li>• shortness of breath</li> <li>• dizziness or lightheadedness</li> <li>• fatigue or lack of energy</li> </ul> <b>Allergy Alert:</b> Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.
<b>Do not use</b> • for more than 7 days unless

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part356	03/05/2019	

**Labeler** - Safco Dental Supply Co. (020417380)

Revised: 1/2022

Safco Dental Supply Co.