

**ACETAMINOPHEN 500MG, DEXTROMETHORPHAN HBR 10MG, GUAIFENESIN 200MG, PHENYLEPHRINE HCL 5MG- acetaminophen 500mg, dextromethorphan hbr 10mg, guaifenesin 200mg, phenylephrine hcl 5mg tablet**  
**Strive Pharmaceuticals Inc**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

**Cold & Flu - Grippa (Acetaminophen 325mg; Dextromethorphan HBr 10mg, Guaifenesin 200mg, Phenylephrine HCl 5mg)**

Acetaminophen 325mg; Dextromethorphan HBr 10mg; Gauifenesin 200mg; Phenylephrine HCl 5mg  
colloidal silicon dioxide, croscarmellose sodium, dibasic calcium phosphate, FD&C yellow#6, magnesium staerate, mica, microcrystalline cellulose, polyethylene glycol, polysorbate 80, polyvinyl alcohol, stearic acid, talc

- take with a full glass of water
- do not take more than directed
- do not exceed 8 caplets per 24 hours

---

adults & children 12 years of age & over	2 caplets with water every 4 hours
children 4 to 12 years of age	ask a doctor
children under 4 years of age	do not use

---

\*temporarily relief of occasional heaches and minor aches and pains with accompanying sleeplessness

**Liver warning:**

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 caplets in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

if a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly

**Do not use**

- with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking ths product
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Ask a doctor or a pharmacist before use if you are**

- taking the blood thinning drug warfarin

**When using this product**

- do not use more than directed

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition

**If pregnant or breast-feeding,**

ask a health professional before use

**Keep out of reach of children**


**Overdose warning:** In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Keep out of reach of children**

**Overdose Warning:** In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Pain reliever / fever reducer, cough suppressant, expectorant, nasal decongestant

- store at room temperature between 20 °C - 25 °C (68 °F - 77 °F)
- avoid excessive heat and humidity



LOT: \_\_\_\_\_  
EXP: \_\_\_\_\_

**DRUG FACTS**

Active ingredients (in each caplet)	Purpose
Acetaminophen 325mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guafenesin 200 mg	Expectorant
Phenylephrine HCl 5mg	Nasal decongestant

**Uses:**  
Temporarily relieves common cold & flu symptoms • nasal congestion • sinus congestion & pressure • cough due to minor throat & bronchial irritation • minor aches & pains • headache • fever • sore throat • reduces swelling of nasal passages • temporarily restores free breathing through the nose • promotes nasal and/or sinus drainage • helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings:**  
**Liver warning:**  
This product contains acetaminophen. Severe liver damage may occur if you take: • more than 8 caplets in 24 hrs, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
• Skin redness • itching • rash  
If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:**  
If sore throat is severe, persists for more than 3 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use:**  
• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
• if you have ever had an allergic reaction to this product or any of its ingredients.

**Ask a doctor before use if you have:**  
• liver disease • heart disease • high blood pressure • thyroid disease • diabetes

LIFT PANEL FOR CONTINUED DRUG FACTS

**DRUG FACTS (Continued)**

• trouble urinating due to an enlarged prostate gland • cough that occurs with too much phlegm (mucus)  
• persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or a pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if:  
• you get nervous, dizzy or sleepless • pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts.  
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions:**  
• Do not take more than directed. See Overdose warning.  
• take with a full glass of water • do not exceed 8 caplets per 24 hours  
adults and children 12 yrs of age & over 2 caplets with water every 4 hrs  
children 4 to 12 years of age ask a doctor  
children under 4 years of age do not use  
when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing.

**Other Information:**  
• Store at room temperature between 20-25°C (68-77°F) • Avoid excessive heat and humidity

**Inactive Ingredients:**  
colloidal silicon dioxide, croscarmellose sodium, dibasic calcium phosphate, FD&C yellow #6 aluminum lake, magnesium stearate, mica, microcrystalline cellulose, polyethylene glycol, polyorbital 80, polyvinyl alcohol, stearic acid, talc.

**DRUG FACTS (Continued)**

Questions Or Comments ?  
1-833-665-9444

**READ AND KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION.**

Distributed By: O4 GLOBAL TRADING USA LLC, 2241 S Watson RD, Suite 121 Arlington, TX 76010  
www.onlyfor.com

Packaged By: Slnve Pharmaceuticals Inc., East Brunswick, NJ 08816.

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF A BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGN OF TAMPERING**

Tear at arrows push through foil

www.onlyfor.com CT2GRP4-REV.102018

**ACETAMINOPHEN 500MG, DEXTROMETHORPHAN HBR 10MG,**

# GUAIFENESIN 200MG, PHENYLEPHRINE HCL 5MG

acetaminophen 500mg, dextromethorphan hbr 10mg, guaifenesin 200mg, phenylephrine hcl 5mg tablet

## Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70692-108
Route of Administration	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

## Inactive Ingredients

Ingredient Name	Strength
STEARIC ACID (UNII: 4ELV7Z65AP)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MICA (UNII: V8A1AW0880)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
TALC (UNII: 7SEV7J4R1U)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	

## Product Characteristics

Color	orange	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	S701
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70692-108-41	4 in 1 PACKAGE; Type 0: Not a Combination Product	12/20/2018	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	12/20/2018	

**Labeler** - Strive Pharmaceuticals Inc (080028013)

Revised: 11/2020

Strive Pharmaceuticals Inc