

**CICLOPIROX OLAMINE- ciclopirox olamine cream**  
**RPK Pharmaceuticals, Inc.**

-----  
**Ciclopirox Olamine**  
**Cream USP, 0.77%**

**Rx Only**

FOR DERMATOLOGIC USE ONLY  
NOT FOR OPHTHALMIC USE


**DESCRIPTION**

Ciclopirox Olamine Cream USP, 0.77% is for topical use.

Each gram of ciclopirox olamine cream USP contains 7.70 mg of ciclopirox (as ciclopirox olamine) in a water miscible vanishing cream base consisting of cetyl alcohol, cocamide DEA, lactic acid, mineral oil, myristyl alcohol, octyldodecanol, polysorbate 60, purified water, sorbitan monostearate, stearyl alcohol, and benzyl alcohol (1%) as preservative.

Ciclopirox olamine cream contains a synthetic, broad-spectrum, antifungal agent ciclopirox (as ciclopirox olamine). The chemical name is 6-cyclohexyl-1-hydroxy-4-methyl-2(1*H*)-pyridone, 2-aminoethanol salt.

The CAS Registry Number is 41621-49-2. The chemical structure is:



**CLINICAL PHARMACOLOGY**

**Mechanism of Action**

Ciclopirox is a hydroxypyridone antifungal agent that acts by chelation of polyvalent cations (Fe<sup>3+</sup> or Al<sup>3+</sup>), resulting in the inhibition of the metal-dependent enzymes that are responsible for the degradation of peroxides within the fungal cell.

**Pharmacokinetics**

Pharmacokinetic studies in men with tagged ciclopirox solution in polyethylene glycol 400 showed an average of 1.3% absorption of the dose when it was applied topically to 750 cm<sup>2</sup> on the back followed by occlusion for 6 hours. The biological half-life was 1.7 hours and excretion occurred via the kidney. Two days after application only 0.01% of the dose applied could be found in the urine. Fecal excretion was negligible.

Penetration studies in human cadaverous skin from the back, with ciclopirox with tagged ciclopirox

showed the presence of 0.8 to 1.6% of the dose in the stratum corneum 1.5 to 6 hours after application. The levels in the dermis were still 10 to 15 times above the minimum inhibitory concentrations.

Autoradiographic studies with human cadaverous skin showed that ciclopirox penetrates into the hair and through the epidermis and hair follicles into the sebaceous glands and dermis, while a portion of the drug remains in the stratum corneum.

Draize Human Sensitization Assay, 21-Day Cumulative Irritancy study, Phototoxicity study, and Photo-Draize study conducted in a total of 142 healthy male subjects showed no contact sensitization of the delayed hypersensitivity type, no irritation, no phototoxicity, and no photo-contact sensitization due to Ciclopirox Olamine Cream USP, 0.77%.

## **INDICATIONS AND USAGE**

Ciclopirox olamine cream is indicated for the topical treatment of the following dermal infections: tinea pedis, tinea cruris, and tinea corporis due to *Trichophyton rubrum*, *Trichophyton mentagrophytes*, *Epidermophyton floccosum*, and *Microsporum canis*; candidiasis (moniliasis) due to *Candida albicans*; and tinea (pityriasis) versicolor due to *Malassezia furfur*.

## **CONTRAINDICATIONS**

Ciclopirox olamine cream is contraindicated in individuals who have shown hypersensitivity to any of its components.

## **WARNINGS**

Ciclopirox olamine cream is not for ophthalmic use.

**Keep out of reach of children.**

## **PRECAUTIONS**

If a reaction suggesting sensitivity or chemical irritation should occur with the use of ciclopirox olamine cream, treatment should be discontinued and appropriate therapy instituted.

### **Information for Patients**

The patient should be told to:

1. Use the medication for the full treatment time even though symptoms may have improved and notify the physician if there is no improvement after four weeks.
2. Inform the physician if the area of application shows signs of increased irritation (redness, itching, burning, blistering, swelling, or oozing) indicative of possible sensitization.
3. Avoid the use of occlusive wrappings or dressings.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

A 104-week dermal carcinogenicity study in mice was conducted with ciclopirox cream applied at doses up to 1.93% (100 mg/kg/day or 300 mg/m<sup>2</sup>/day). No increase in drug related neoplasms was noted when compared to control.

The following *in vitro* genotoxicity tests have been conducted with ciclopirox: evaluation of gene mutation in the Ames Salmonella and E. coli assays (negative); chromosome aberration assays in V79 Chinese hamster lung fibroblast cells, with and without metabolic activation (positive); chromosome aberration assays in V79 Chinese hamster lung fibroblast cells in the presence of supplemental Fe<sup>3+</sup>, with and without metabolic activation (negative); gene mutation assays in the HGPRT-test with V79 Chinese hamster lung fibroblast cells (negative); and a primary DNA damage assay (i.e., unscheduled

DNA synthesis assay in A549 human cells) (negative). An *in vitro* cell transformation assay in BALB/c 3T3 cells was negative for cell transformation. In an *in vivo* Chinese hamster bone marrow cytogenetic assay, ciclopirox was negative for chromosome aberrations at a dosage of 5000 mg/kg body weight.

A combined oral fertility and embryofetal developmental study was conducted in rats with ciclopirox olamine. No effect on fertility or reproductive performance was noted at the highest dose tested of 3.85 mg/kg/day ciclopirox (approximately 1.2 times the maximum recommended human dose based on body surface area comparisons).

## **Pregnancy**

### **Teratogenic Effects**

#### *Pregnancy Category B*

There are no adequate or well-controlled studies in pregnant women. Therefore, ciclopirox olamine cream should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Oral embryofetal developmental studies were conducted in mice, rats, rabbits and monkeys. Ciclopirox or ciclopirox olamine was orally administered during the period of organogenesis. No maternal toxicity, embryotoxicity or teratogenicity were noted at the highest doses of 77, 125, 80 and 38.5 mg/kg/day ciclopirox in mice, rats, rabbits and monkeys, respectively (approximately 11, 37, 51 and 24 times the maximum recommended human dose based on body surface area comparisons, respectively).

Dermal embryofetal developmental studies were conducted in rats and rabbits with ciclopirox olamine dissolved in PEG 400. Ciclopirox olamine was topically administered during the period of organogenesis. No maternal toxicity, embryotoxicity or teratogenicity were noted at the highest doses of 92 mg/kg/day and 77 mg/kg/day ciclopirox in rats and rabbits, respectively (approximately 27 and 49 times the maximum recommended human dose based on body surface area comparisons, respectively).

## **Nursing Mothers**

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when ciclopirox olamine cream is administered to a nursing woman.

## **Pediatric Use**

Safety and effectiveness in pediatric patients below the age of 10 years have not been established.

## **ADVERSE REACTIONS**

In all controlled clinical studies with 514 patients using ciclopirox olamine cream and in 296 patients using the vehicle cream, the incidence of adverse reactions was low. This included pruritus at the site of application in one patient and worsening of the clinical signs and symptoms in another patient using ciclopirox cream and burning in one patient and worsening of the clinical signs and symptoms in another patient using the vehicle cream.

## **DOSAGE AND ADMINISTRATION**

Gently massage ciclopirox olamine cream into the affected and surrounding skin areas twice daily, in the morning and evening. Clinical improvement with relief of pruritus and other symptoms usually occurs within the first week of treatment. If a patient shows no clinical improvement after four weeks of treatment with ciclopirox olamine cream, the diagnosis should be redetermined. Patients with tinea versicolor usually exhibit clinical and mycological clearing after two weeks of treatment.

## **HOW SUPPLIED**

Product: 53002-9450

NDC: 53002-9450-2 15 g in a TUBE

NDC: 53002-9450-1 30 g in a TUBE

Mfd. by: Taro Pharmaceuticals Inc., Brampton, Ontario, Canada L6T 1C1

Dist. by: **Taro Pharmaceuticals U.S.A., Inc.**, Hawthorne, NY 10532

Revised: July 2013

PK-4597-1 226

## Ciclopirox Olamine 0.77% Cream

**Front of Tube:**  
NDC# 53002-9450-1  
LIST# 51672-1318-1  
**CICLOPIROX OLAMINE 0.77% CREAM**  
15 Gm Tube  
LOT# 18267-001  
EXP DATE: 08-31-2019  
Shape & Color: \_\_\_\_\_  
Markings: \_\_\_\_\_  
**Rx only**  
PATIENT NAME: \_\_\_\_\_  
**CICLOPIROX 0.77% CREAM**  
TARO  
Rx# 182660871000 ITEM# 9450 945-03  
**APPLY TO AFFECTED AREA TWICE A DAY OR AS DIRECTED.**  
CLINIC NAME GOES HERE  
THIS DRUG IS AN ANTIFUNGAL FOR EXTERNAL USE ONLY. USE AS DIRECTED & FOR THE FULL LENGTH OF TREATMENT PRESCRIBED. NOT FOR OPHTHALMIC USE.  
Date Dispensed: \_\_\_\_\_  
182660871000

**Back of Tube:**  
APP AFF AREA (SD OR LD)  
LOT# 18267-001 EXP 08-31-2019  
Rx# 182660871 000 FCA-9450  
15 Gm CICLOPIROX 0.77% CREAM  
**CICLOPIROX OLAMINE 0.77% CREAM**  
15 Gm Tube  
DISCARD BY 08-31-2019  
NDC# 53002-9450-1  
Rx# 182660871 000  
Clinic Name Here  
BILLING NDC# 51672-1318-01  
Rx# 182660871 000  
15 Gm CICLOPIROX 0.77% CREAM  
BILLING NDC# 51672-1318-01  
Rx# 182660871 000  
15 Gm CICLOPIROX 0.77% CREAM  
15 Gm CICLOPIROX 0.77% CREAM  
Rx# 182660871000 ITEM# 9450

## CICLOPIROX OLAMINE

ciclopirox olamine cream

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:53002-9450(NDC:51672-1318)
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CICLOPIROX OLAMINE (UNII: 50MD4SB4AP) (CICLOPIROX - UNII:19W019ZDRJ)	CICLOPIROX	7.7 mg in 1 g

### Inactive Ingredients

Ingredient Name	Strength
benzyl alcohol (UNII: LKG8494WBH)	
cetyl alcohol (UNII: 936JST6JCN)	
Coco Diethanolamide (UNII: 92005F972D)	
LACTIC ACID, UNSPECIFIED FORM (UNII: 33X04XA5AT)	
mineral oil (UNII: T5L8T28FGP)	
myristyl alcohol (UNII: V4203409PU)	
octyldodecanol (UNII: 461N10614Y)	

<b>polysorbate 60</b> (UNII: CAL22UVI4M)	
<b>water</b> (UNII: 059QF0K00R)	
<b>sorbitan monostearate</b> (UNII: NVZ4I0H58X)	
<b>stearyl alcohol</b> (UNII: 2KR89I4H1Y)	

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53002-9450-2	15 g in 1 TUBE; Type 0: Not a Combination Product	10/01/2018	
2	NDC:53002-9450-1	30 g in 1 TUBE; Type 0: Not a Combination Product	10/01/2018	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076790	04/12/2005	

**Labeler** - RPK Pharmaceuticals, Inc. (147096275)

### Establishment

Name	Address	ID/FEI	Business Operations
RPK Pharmaceuticals, Inc.		147096275	RELABEL(53002-9450) , REPACK(53002-9450)

Revised: 12/2020

RPK Pharmaceuticals, Inc.