

SELENIUM SULFIDE- selenium sulfide lotion
A-S Medication Solutions

SELENIUM SULFIDE LOTION, USP 2.5%
SHAKE WELL BEFORE USING.

APPLICATION INSTRUCTIONS

Keep tightly capped. SHAKE WELL BEFORE USING. Product may damage jewelry; remove jewelry before use.

For treatment of tinea versicolor:

1. Apply to affected areas and lather with a small amount of water.
2. Allow to remain on skin for 10 minutes.
3. Rinse body thoroughly.
4. Repeat this procedure once a day for 7 days.

For treatment of dandruff and seborrheic dermatitis of the scalp.

1. Massage 1 or 2 teaspoonfuls of shampoo into wet scalp.
2. Allow to remain on scalp for 2 to 3 minutes.
3. Rinse scalp thoroughly.
4. Repeat application and rinse thoroughly.
5. After treatment, wash hands well.
6. Repeat treatments as directed by physician.

WARNINGS AND PRECAUTIONS

For External Use Only. Do not use on broken skin or inflamed areas. If allergic reactions occur, discontinue use. Avoid getting shampoo in eyes or in contact with genital area as it may cause irritation and burning.

FOR EXTERNAL USE ONLY. WARNING: KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Store at controlled room temperature, (15 - 30) °C ((59 - 86) °F) [see USP].

CLINICAL PHARMACOLOGY

Selenium sulfide appears to have a cytostatic effect on cells of the epidermis and follicular epithelium, reducing corneocyte production.

INDICATIONS AND USAGE

For treatment of tinea versicolor, seborrheic dermatitis of scalp and treatment of dandruff.

CONTRAINDICATIONS

Not to be used by patients allergic to any of its ingredients.

PRECAUTIONS

General:

Not to be used when acute inflammation or exudation is present as increased absorption may occur.

Information for Patients:

See Warnings and Precautions section under Application Instructions.

Carcinogenesis:

Dermal application of 25% and 50% solutions of 2.5% selenium sulfide lotion on mice over an 88 week period, indicated no carcinogenic effects.

Pregnancy:

WHEN USED ON BODY SURFACES FOR THE TREATMENT OF TINEA VERSICOLOR, SELENIUM SULFIDE LOTION, USP 2.5% IS CLASSIFIED AS PREGNANCY CATEGORY C. Animal reproduction studies have not been conducted with selenium sulfide. It is also not known whether selenium sulfide can cause fetal harm when applied to body surfaces of a pregnant woman or can affect reproduction capacity. Under ordinary circumstances selenium sulfide should not be used for the treatment of tinea versicolor in pregnant women.

Pediatric Use:

Safety and effectiveness in infants have not been established.

ADVERSE REACTIONS

In decreasing order of severity: skin irritation; occasional reports of increase in normal hair loss; discoloration of hair (can be avoided or minimized by thorough rinsing of hair after treatment). As with other shampoos, oiliness or dryness of hair and scalp may occur.

OVERDOSAGE

Accidental Oral Ingestion:

No documented reports of serious toxicity in humans resulting from acute ingestion of selenium sulfide, however, acute toxicity studies in animals suggest that ingestion of large amounts could result in potential human toxicity. Evacuation of the stomach contents should be considered in cases of acute oral ingestion.

DOSAGE AND ADMINISTRATION

See Application Instructions.

For treatment of tinea versicolor

Apply to affected areas and lather with a small amount of water. Allow product to remain on skin for 10 minutes, then rinse the body thoroughly. Repeat procedure once a day for 7 days.

For treatment of dandruff and seborrheic dermatitis

Usually two applications each week for two weeks will afford control. After this, the lotion may be used at less frequent intervals – weekly, every two weeks, or every 3 or 4 weeks in some cases. Should not be applied more frequently than required to maintain control.

HOW SUPPLIED

Product: 50090-2166

NDC: 50090-2166-0 118 mL in a BOTTLE, PLASTIC

WARNINGS

KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

Rx Only

Product No.: 8528

A50-8528-04

REV. 01-05

Manufactured By:

Morton Grove Pharmaceuticals, Inc.

Morton Grove, IL 60053

Selenium Sulfide



SELENIUM SULFIDE

selenium sulfide lotion

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:50090-2166(NDC:60432-528)
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SELENIUM SULFIDE (UNII: Z 69D9E381Q) (SELENIUM SULFIDE - UNII:Z 69D9E381Q)	SELENIUM SULFIDE	25 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
BENTONITE (UNII: A3N5ZCN45C)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
WATER (UNII: 059QF0KO0R)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM (UNII: 3980JIH2SW)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
DISODIUM COCOAMPHODIACETATE (UNII: 18L9G3U51M)	
GLYCOL STEARATE (UNII: 0324G66D0E)	
LAURIC DIETHANOLAMIDE (UNII: I29I2VHG38)	
CAPTAN (UNII: EOL5G26Q9F)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
GLYCERYL RICINOLEATE (UNII: ZUE0CEL42O)	

Packaging

		Marketing Start	Marketing End
--	--	-----------------	---------------

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50090-2166-0	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/21/2015	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA088228	09/01/1983	

Labeler - A-S Medication Solutions (830016429)

Establishment

Name	Address	ID/FEI	Business Operations
A-S Medication Solutions		830016429	RELABEL(50090-2166)

Revised: 2/2021

A-S Medication Solutions