

**FAMOTIDINE- famotidine tablet**  
**Chain Drug Consortium, LLC**

-----  
**Dr.Reddy's Laboratories Limited**

**Active ingredient (in each tablet)**

Famotidine USP, 10 mg/20 mg

**Purpose**

Acid reducer

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings**

**Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

**Ask a doctor or pharmacist before use if you are** taking a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

## **If pregnant or breast-feeding,**

ask a health professional before use.

## **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## **Directions**

- **For Famotidine 10 mg:**
- adults and children 12 years and over:
  - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
  - to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **15 to 60 minutes before** eating food or drinking beverages that cause heartburn
  - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor
- **For Famotidine 20 mg:**
- adults and children 12 years and over:
  - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
  - to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
  - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

## **Other information**

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture and light

## **Inactive ingredients**

colloidal silicon dioxide, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, synthetic red iron oxide (only in 10 mg), talc and titanium dioxide

## **Questions or comments?**

call **1-888-375-3784**

## **Tips for Managing Heartburn**

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed

- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

## Principal Display Panel - 10 mg Carton

111580061





**ORIGINAL STRENGTH**  
**Acid Reducer**  
Famotidine Tablets  
USP, 10 mg

Drug Facts	Purpose
<b>Active ingredient (in each tablet)</b> Famotidine USP, 10 mg	Acid reducer
<b>Uses</b>	
<ul style="list-style-type: none"> <li>■ relieves heartburn associated with acid indigestion and sour stomach</li> <li>■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages</li> </ul>	
<b>Warnings</b>	
<p><b>Allergy alert:</b> Do not use if you are allergic to famotidine or other acid reducers</p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.</li> <li>■ with other acid reducers</li> </ul> <p><b>Ask a doctor before use if you have</b></p> <ul style="list-style-type: none"> <li>■ had heartburn over 3 months. This may be a sign of a more serious condition.</li> <li>■ heartburn with lightheadedness, sweating, or dizziness</li> <li>■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck, or shoulders; or lightheadedness</li> <li>■ frequent chest pain</li> <li>■ frequent wheezing, particularly with heartburn</li> <li>■ unexplained weight loss</li> <li>■ nausea or vomiting</li> <li>■ stomach pain</li> <li>■ kidney disease</li> </ul>	
<b>Drug Facts (continued)</b>	
<p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>■ your heartburn continues or worsens</li> <li>■ you need to take this product for more than 14 days</li> </ul> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>	
<b>Directions</b>	
<ul style="list-style-type: none"> <li>■ adults and children 12 years and over:               <ul style="list-style-type: none"> <li>■ to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.</li> <li>■ to prevent symptoms, swallow 1 tablet with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn</li> </ul> </li> <li>■ do not use more than 2 tablets in 24 hours</li> <li>■ children under 12 years: ask a doctor</li> </ul>	
<b>Other information</b>	
<ul style="list-style-type: none"> <li>■ read the directions and warnings before use</li> <li>■ keep the carton. It contains important information.</li> <li>■ store at 20°-25°C (68°-77°F)</li> <li>■ protect from moisture</li> </ul>	
<b>Inactive ingredients</b>	
colloidal silicon dioxide, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, synthetic red iron oxide, talc and titanium dioxide	
<b>Questions?</b> call 1-888-375-3784	

**Tips for Managing Heartburn**

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

**JUST ONE TABLET** prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods or beverages

**TAMPER EVIDENT: DO NOT USE IF FOIL SEAL UNDER CAP PRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING**

 \*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Pepcid AC®. Pepcid AC® is a registered trademark of Johnson & Johnson.

Distributed By: Pharmacy Value Alliance, LLC  
407 East Lancaster Avenue,  
Wayne, PA 19087

Made in India  
REV 07/21



150085111

Compare to the active ingredient in Pepcid AC®



**ORIGINAL STRENGTH**  
**Acid Reducer**  
Famotidine Tablets  
USP, 10 mg

Just One Tablet!  
Prevents & Relieves Heartburn Due to Acid Indigestion

**30 TABLETS**

actual size





8 40986 03722 1

LOT

EXP

## Principal Display Panel - 20 mg Carton



## FAMOTIDINE

famotidine tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:68016-800
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	10 mg

### Inactive Ingredients

Ingredient Name	Strength
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
STARCH, CORN (UNII: O8232NY3SJ)	
Polyethylene Glycol, Unspecified (UNII: 3WJQ0SDW1A)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	

## Product Characteristics

<b>Color</b>	PINK	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	6mm
<b>Flavor</b>		<b>Imprint Code</b>	C;118
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68016-800-30	1 in 1 CARTON	11/01/2021	
1		30 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077367	11/01/2021	

## FAMOTIDINE

famotidine tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:68016-801
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>FAMOTIDINE</b> (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	20 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>Polyethylene Glycol, Unspecified</b> (UNII: 3WJQ0SDW1A)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

**Product Characteristics**

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	8mm
<b>Flavor</b>		<b>Imprint Code</b>	L1
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68016-801-25	1 in 1 CARTON	11/01/2021	
1		25 in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077367	11/01/2021	

**Labeler** - Chain Drug Consortium, LLC (101668460)

Revised: 10/2024

Chain Drug Consortium, LLC