

**RIGHT REMEDIES ASPIRIN LOW DOSE 81 MG- aspirin tablet**  
**Strive Pharmaceuticals Inc**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**RIGHT REMEDIES Aspirin LOW DOSE 81 mg**

**Drug Facts**

**Active ingredient (in each tablet)**

Aspirin 81 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever

**Uses**

- for the temporary relief of minor aches and pains or as recommended by your doctor.
- Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**
- ask your doctor about other uses for 81 mg Aspirin

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** aspirin may cause a severe allergic reaction which may include:

- hives • facial swelling • asthma (wheezing) • shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if** taking a prescription drug for • gout • diabetes • arthritis

**Stop use and ask a doctor if** • an allergic reaction occurs. Seek medical help right away.

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur
  - ringing in the ears or a loss of hearing occurs

**If pregnant or breast-feeding**, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Keep out of reach of children.**

### ***Directions***

- drink a full glass of water with each dose

adults and children 12 years and over	<ul style="list-style-type: none"> <li>• take 4 to 8 tablets every 4 hours</li> <li>• do not take more than 48 tablets in 24 hours</li> </ul>
children under 12 years	<ul style="list-style-type: none"> <li>• consult a doctor</li> </ul>

### Other information

- store at 20° -25°C (68° - 77°F)
- Avoid excessive heat above 40°C (104°F)

### Inactive ingredients



colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10, FD&C yellow #6, fumed silica, methacrylic acid and ethyl acrylate co-polymer, microcrystalline cellulose, stearic acid, talc, titanium dioxide, triethyl citrate

### Questions or comments?

call **1-888-577-8033** Monday - Friday 8am - 4pm EST

### Packaging

#### PACKAGE - 300 tablets



Compare to active ingredient in  
**BAYER® Low Dose Aspirin**

**LOW DOSE**

**Aspirin**

Pain reliever  
Aspirin regimen  
with safety coating

(NSAID)  
**300**  
tablets

actual size

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**Drug Facts**

<b>Active ingredient (in each tablet)</b> Aspirin 81 mg (NSAID)*	<b>Purpose</b> Pain reliever
---	---------------------------------

\*nonsteroidal anti-inflammatory drug

**Uses**

- for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**
- ask your doctor about other uses for 81 mg Aspirin

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Drug Facts (continued under label)**

†This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Bayer® Low Dose Aspirin

Distributed by:  
Strive Pharmaceuticals Inc.  
East Brunswick, NJ 08816

REV.00-062021 LB7069216131



PEEL HERE

LOT: \_\_\_\_\_

EXP: \_\_\_\_\_

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if taking a prescription drug for**

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur
  - ringing in the ears or a loss of hearing occurs

**If pregnant or breast-feeding,** ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Directions**

- drink a full glass of water with each dose

adults and children 12 years and over	take 4 to 8 tablets every 4 hours
children under 12 years	do not take more than 48 tablets in 24 hours
	consult a doctor

**Other information**

- store at 20° -25°C (68° -77°F)
- Avoid excessive heat above 40°C (104°F)

**Inactive ingredients**

colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10, FD&C yellow #6, fumed silica, methacrylic acid and ethyl acrylate co-polymer, microcrystalline cellulose, stearic acid, talc, titanium dioxide, triethyl citrate

**Questions or comments?**

call 1-888-577-8033 Monday - Friday 8am - 4pm EST



**ADHESIVE  
DO NOT PRINT**

**PACKAGE - 365 tablets**

†compare to active ingredient in  
**BAYER® Low Dose Aspirin**

**RIGHT REMEDIES**

**LOW DOSE**

**Aspirin**

**Purpose**  
Pain reliever

is recommended by your doctor if you do not provide fast relief of pain.

are recovering from chicken pox or flu-like symptoms, if you are using this product, if you consult a doctor because of a rare but serious illness, or if you are taking other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others).

in which may include:  
fever, stomach pain, or  
stomach bleeding, which may cause severe stomach bleeding.

typical NSAIDs (aspirin, ibuprofen, naproxen, or others)

in this product

er label)  
r HealthCare LLC,

Distributed by:  
Strive Pharmaceuticals Inc.  
East Brunswick, NJ 08816

No Varnish

**Aspirin**  
Pain reliever  
aspirin regimen  
enteric safety coated

aspirin (NSAID)  
**81 mg**  
each tablet

Actual Size

365 tablets

**RIGHT KIDNEES**

**LOW DOSE**

Compare to active ingredient in  
**BAYER®** Low Dose Aspirin

**Drug Facts**  
Active ingredient (in each tablet)  
Aspirin 81 mg (NSAID)\*  
\*nonsteroidal anti-inflammatory drug

**Uses**  
for the temporary relief of minor aches and pains or a doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief. ask your doctor about other uses for Aspirin 81 mg

**Warnings**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
Allergy alert: Aspirin may cause a severe allergic reaction which may include:  
hives ■ facial swelling ■ asthma (wheezing) ■ shock  
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you  
are age 60 or older  
have had stomach ulcers or bleeding problems  
take a blood thinning (anticoagulant) or steroid drug  
take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
have 3 or more alcoholic drinks every day while using this product  
take more or for a longer time than directed

**Drug Facts (continued under label)**  
This product is not manufactured or distributed by Bayer distributor of Bayer® Low Dose Aspirin.

LOT: REV00-082021 EXP: LB7069216165

3 70692 00055 9

NDC 70692-161-65

aspirin (NSAID)  
**81 mg**  
each tablet

Actual Size

365 tablets

**Aspirin**  
Pain reliever  
aspirin regimen  
enteric safety coated

**RIGHT KIDNEES**

**LOW DOSE**

Compare to active ingredient in  
**BAYER®** Low Dose Aspirin

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

**Drug Facts**

**Active ingredient (in each tablet)** Purpose  
Aspirin 81 mg (NSAID)\* Pain reliever  
\*nonsteroidal anti-inflammatory drug

**Uses**  
for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief. ask your doctor about other uses for Aspirin 81 mg

**Warnings**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
Allergy alert: Aspirin may cause a severe allergic reaction which may include:  
hives ■ facial swelling ■ asthma (wheezing) ■ shock  
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you  
are age 60 or older  
have had stomach ulcers or bleeding problems  
take a blood thinning (anticoagulant) or steroid drug  
take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
have 3 or more alcoholic drinks every day while using this product  
take more or for a longer time than directed

**Drug Facts (continued under label)**

This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Bayer® Low Dose Aspirin.

3 70692 00055 9

Distributed by:  
Strive Pharmaceuticals Inc.  
East Brunswick, NJ 08816

**Drug Facts (continued)**

**Do not use**  
if you are allergic to aspirin or any other pain reliever/fever reducer  
if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**  
stomach bleeding warning applies to you  
you have a history of stomach problems, such as heartburn  
you have high blood pressure, heart disease, liver cirrhosis, or kidney disease  
you are taking a diuretic  
you have asthma

**Ask a doctor or pharmacist before use if you are taking a prescription drug for**  
gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if**  
an allergic reaction occurs. Seek medical help right away.  
you experience any of the following signs of stomach bleeding:  
feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better  
pain gets worse or lasts more than 10 days  
fever gets worse or lasts more than 3 days  
redness or swelling is present  
new symptoms occur  
ringing in the ears or a loss of hearing occurs

**If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Directions**  
drink a full glass of water with each dose

adults and children 12 years and over	take 4 to 8 tablets every 4 hours do not take more than 48 tablets in 24 hours
children under 12 years	consult a doctor

**Other information**  
store at room temperature between 20°-25°C (68°-77°F)  
Avoid excessive heat above 40°C (104°F)

**Inactive ingredients**  
colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10, FD&C yellow #6, fumed silica, methacrylic acid and ethyl acrylate co-polymer, microcrystalline cellulose, stearic acid, talc, titanium dioxide, triethyl citrate

**Questions or comments?**  
1-888-577-8033 Monday - Friday 8am - 4pm EST

PEEL  
HERE

LOT: No Varnish  
 EXP: REV.00-082021  
 LB7069216165

**ADHESIVE  
 DO NOT PRINT**

**PACKAGE - 500 tablets**

†compare to active ingredient in  
**BAYER®** Low Dose Aspirin

**RIGHT  
 REMEDIES**

**LOW DOSE  
 Aspirin**

**pain reliever  
 aspirin regimen  
 enteric safety coated**

aspirin (NSAID)  
**81 mg**  
 each tablet

**500  
 tablets**

actual size

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**Drug Facts**

**Active ingredient (in each tablet)**  
 Aspirin 81 mg (NSAID)\* ..... Pain reliever

**Uses**

■ for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

■ ask your doctor about other uses for 81 mg Aspirin

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** aspirin may cause a severe allergic reaction which may include:

- hives ■ facial swelling ■ asthma (wheezing) ■ shock


**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed


**Drug Facts (continued under label)**

†This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Bayer® Low Dose Aspirin

REV.00-082021 LB7069216165



3 70692 00025 1 2



PEEL  
HERE

REV.00-082021  
 LB7069216165

Distributed by:  
 Shire Pharmaceuticals Inc.  
 East Brunswick, NJ 08816

NDC 70692-161-50

each tablet

aspirin (NSAID)  
**81 mg**  
 each tablet

**500  
 tablets**

actual size

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**RIGHT  
 REMEDIES**

**LOW DOSE  
 Aspirin**

**pain reliever  
 aspirin regimen  
 enteric safety coated**

aspirin (NSAID)  
**81 mg**  
 each tablet

**500  
 tablets**

actual size

†compare to active ingredient in  
**BAYER®** Low Dose Aspirin

**Drug Facts**

**Active ingredient (in each tablet)**  
 Aspirin 81 mg (NSAID)\* ..... Pain reliever

**Uses**

■ for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

■ ask your doctor about other uses for 81 mg Aspirin

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** aspirin may cause a severe allergic reaction which may include:

- hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Drug Facts (continued)**

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if taking a prescription drug for**

- gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if** ■ an allergic reaction occurs. Seek medical help right away.

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

**If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Directions**

- drink a full glass of water with each dose

adults and children 12 years and over	<ul style="list-style-type: none"> <li>■ take 4 to 8 tablets every 4 hours</li> <li>■ do not take more than 48 tablets in 24 hours</li> </ul>
children under 12 years	<ul style="list-style-type: none"> <li>■ consult a doctor</li> </ul>

**Other information**

- store at 20° -25°C (68° -77°F)

**Drug Facts (continued under label)**

This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Bayer® Low Dose Aspirin

Distributed by:  
Strive Pharmaceuticals Inc.  
East Brunswick, NJ 08816

REV.00-062021 LB7069216150



3 70692 00025 2



PEEL HERE

■ Avoid excessive heat above 40°C (104°F)

**Inactive ingredients**  
colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10, FD&C yellow #6, fumed silica, methacrylic acid and ethyl acrylate co-polymer, microcrystalline cellulose, stearic acid, talc, titanium dioxide, triethyl citrate

**Questions or comments?**  
call 1-888-577-8033 Monday - Friday 8am - 4pm EST

**PACKAGE - 1000 tablets**

†compare to active ingredient in  
**BAYER® Low Dose Aspirin**

**RIGHT REMEDIES**

**LOW DOSE Aspirin**

pain reliever  
aspirin regimen  
enteric safety coated

aspirin(NSAID)  
**81 mg**  
each tablet

**1000 tablets**

actual size

**Drug Facts**

**Active ingredient (in each tablet)**  
Aspirin 81 mg (NSAID)\* ..... Pain reliever  
\*nonsteroidal anti-inflammatory drug

**Purpose**  
Pain reliever

**Uses**  
■ for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.  
■ ask your doctor about other uses for 81 mg Aspirin


**Warnings**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
Allergy alert: aspirin may cause a severe allergic reaction which may include:  
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ take a blood thinning (anticagulant) or steroid drug  
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
■ have 3 or more alcoholic drinks every day while using this product  
■ take more or for a longer time than directed

**Drug Facts (continued under label)**

This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Bayer® Low Dose Aspirin


Distributed by:  
Strive Pharmaceuticals Inc.  
East Brunswick, NJ 08816

REV.00-062021 LB7069216111



3 70692 12901 4

LOT: \_\_\_\_\_  
EXP: \_\_\_\_\_



PEEL HERE

NDC 70692-161-11

actual size

aspirin(NSAID)  
**81 mg**  
each tablet

**RIGHT REMEDIES**

**LOW DOSE Aspirin**

pain reliever  
aspirin regimen  
enteric safety coated

†compare to active ingredient in  
**BAYER® Low Dose Aspirin**

**Drug Facts**

**Active ingredient (in each tablet)**  
Aspirin 81 mg (NSAID)\* ..... Pain reliever  
\*nonsteroidal anti-inflammatory drug

**Purpose**  
Pain reliever

**Uses**  
■ for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.  
■ ask your doctor about other uses for 81 mg Aspirin

**Warnings**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this

**Drug Facts (continued)**

**Do not use**  
■ if you are allergic to aspirin or any other pain reliever/fever reducer  
■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**  
■ stomach bleeding warning applies to you  
■ you have a history of stomach problems, such as heartburn  
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease  
■ you are taking a diuretic  
■ you have asthma

**Ask a doctor or pharmacist before use if taking a prescription drug for**  
■ gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if** ■ an allergic reaction occurs. Seek medical help right away.  
■ you experience any of the following signs of stomach bleeding:  
■ feel faint  
■ vomit blood  
■ have bloody or black stools  
■ have stomach pain that does not get better  
■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present  
■ new symptoms occur  
■ ringing in the ears or a loss of hearing occurs

**If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Drug Facts (continued under label)**

This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Bayer® Low Dose Aspirin

REV.00-052021 LB7069216111

Distributed by:  
Strive Pharmaceuticals Inc.  
East Brunswick, NJ 08816



LOT:

EXP:



**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Directions**

■ drink a full glass of water with each dose

adults and children 12 years and over	■ take 4 to 8 tablets every 4 hours ■ do not take more than 48 tablets in 24 hours
children under 12 years	■ consult a doctor

**Other information**

- store at 20° -25°C (68° - 77°F)
- Avoid excessive heat above 40°C (104°F)

**Inactive ingredients**

colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10, FD&C yellow #6, fumed silica, methacrylic acid and ethyl acrylate co-polymer, microcrystalline cellulose, stearic acid, talc, titanium dioxide, triethyl citrate

**Questions or comments?**

call 1-888-577-8033 Monday - Friday 8am - 4pm EST

ADHESIVE

**RIGHT REMEDIES ASPIRIN LOW DOSE 81 MG**

aspirin tablet

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:70692-161
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER</b> (UNII: NX76LV5T8J)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)	

**Product Characteristics**



<b>Color</b>	yellow	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	7mm
<b>Flavor</b>		<b>Imprint Code</b>	A1
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70692-161-31	300 in 1 BOTTLE; Type 0: Not a Combination Product	06/09/2021	
2	NDC:70692-161-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	06/09/2021	
3	NDC:70692-161-11	1000 in 1 BOTTLE; Type 0: Not a Combination Product	06/09/2021	
4	NDC:70692-161-65	365 in 1 BOTTLE; Type 0: Not a Combination Product	12/20/2021	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	06/09/2021	

**Labeler** - Strive Pharmaceuticals Inc (080028013)

Revised: 12/2021

Strive Pharmaceuticals Inc