

**VITACILINA FIRST AID ANTIBIOTIC- bacitracin zinc, neomycin sulfate,  
polymyxin b sulfate ointment  
Taisho Pharmaceutical California Inc.**

-----  
**Vitacilina First Aid Antibiotic**

***Drug Facts***

***Active ingredient (in each gram)***

Bacitracin Zinc 400 units

Neomycin sulfate 3.5 mg

Polymyxin B sulfate 5,000 units

***Purpose***

First aid antibiotic

***Use***

First aid to help prevent infection in minor cuts, scrapes, and burns

***Warnings***

**For external use only.**

**Do not use**

- if you are allergic to any of the ingredients
- in the eyes
- over large areas of the body

**Ask a doctor before use if you have**

- deep or puncture wounds
- animal bites
- serious burns

**Stop use and ask a doctor if**

- you need to use longer than 1 week
- condition persists or gets worse
- rash or other allergic reaction develops

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center

**Directions**

- clean the affected area
- apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage

**Other information**

- store at 15 to 30°C (59 to 86°F )
- DO NOT USE IF SEAL ON TUBE IS BROKEN OR MISSING. **Tamper Evident:**

**Inactive ingredient**

petrolatum

**Package Labeling 28g**

The diagram illustrates the packaging for Vitacilina 28g. It features two main panels: a front panel and a back panel. The front panel is green and white, with 'VITACILINA' in large white letters on a green background. Above it, it says 'FIRST AID ANTIBIOTIC' and 'NET WT 1 OZ (28 g)'. The back panel is white with a green grid pattern and contains detailed text including 'Drug Facts', 'Warnings', 'Directions', and 'Use'. A barcode is located on the top right of the front panel. A glue flap is on the bottom right. Callouts indicate 'NO COLOR' for the top and bottom panels and 'GLUE FLAP NO COLOR / TEXT NO COATING' for the bottom flap.

**Top Panel (Back of Tube):**

- Drug Facts (continued)**
- Other information**
  - store at 15 to 30°C (59 to 86°F)
  - Tamper Evident: DO NOT USE IF SEAL ON TUBE IS BROKEN OR MISSING
- Inactive ingredient** petrolatum
- DISTRIBUTED BY**  
Taisho Pharmaceutical California Inc. 3528 Torrance Blvd, Suite #320 Torrance, CA 90503 U.S.A.
- Barcode:** 3 48875 00000 9
- Text:** PL33103B V01 NO COLOR

**Front Panel:**

- Text:** FIRST AID ANTIBIOTIC (top left and right)
- Text:** NDC 81929-003-01 (top right)
- Text:** VITACILINA® (center, large white letters on green background)
- Text:** NET WT 1 OZ (28 g) (bottom right)

**Bottom Panel (Back of Tube):**

- Text:** FIRST AID ANTIBIOTIC (top left and right)
- Text:** NDC 81929-003-01 (top right)
- Text:** VITACILINA® (center, large white letters on green background)
- Text:** NET WT 1 OZ (28 g) (bottom right)

**Bottom Panel (Front of Tube):**

<b>Drug Facts</b>		<b>Drug Facts (continued)</b>
<b>Active ingredient (in each gram)</b>	<b>Purpose</b>	▪ animal bites ▪ serious burns
Bacitracin zinc 400 units.....First aid antibiotic	.....First aid antibiotic	<b>Stop use and ask a doctor if</b> ▪ you need to use longer than 1 week
Neomycin sulfate 3.5 mg.....First aid antibiotic	.....First aid antibiotic	▪ condition persists or gets worse ▪ rash or other allergic reaction develops
Polymyxin B sulfate 5,000 units.....First aid antibiotic	.....First aid antibiotic	<b>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away: 800-222-1222.</b>
<b>Use</b>	First aid to help prevent infection in minor cuts, scrapes, and burns	<b>Directions</b>
<b>Warnings</b>	For external use only.	▪ clean the affected area
<b>Do not use</b> ▪ if you are allergic to any of the ingredients ▪ in the eyes	▪ over large areas of the body	▪ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
<b>Ask a doctor before use if you have</b> ▪ deep or puncture wounds		▪ may be covered with a sterile bandage

**Glue Flap:** GLUE FLAP NO COLOR / TEXT NO COATING

Add image transcription here...

FIRST AID ANTIBIOTIC

NDC 81929-003-01

# VITACILINA®

NET WT 1 OZ (28 g)

<b>Drug Facts</b>		<b>Drug Facts</b> (continued)	
<b>Active ingredient (in each gram)</b>	<b>Purpose</b>	<b>Stop use and ask a doctor if</b> ■ you need to use longer than 1 week ■ condition persists or gets worse ■ rash or other allergic reaction develops	
Bacitracin zinc 400 units.....	First aid antibiotic	<b>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center</b>	
Neomycin sulfate 3.5 mg.....	First aid antibiotic	<b>Directions</b> ■ clean the affected area ■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily ■ may be covered with a sterile bandage	
Polymyxin B sulfate 5,000 units....	First aid antibiotic	<b>Other information</b> ■ store at 15 to 30°C (59 to 86°F) ■ <b>Tamper Evident: DO NOT USE IF SEAL ON TUBE IS BROKEN OR MISSING.</b>	
<b>Use</b> First aid to help prevent infection in minor cuts, scrapes, and burns		<b>Inactive ingredient</b> petrolatum	
<b>Warnings</b> For external use only.		<b>DISTRIBUTED BY</b> Taisho Pharmaceutical California Inc. PL33103T 3528 Torrance Blvd, Suite #320 Torrance, CA 90503 U.S.A. V01	
<b>Do not use</b> ■ if you are allergic to any of the ingredients ■ in the eyes ■ over large areas of the body			
<b>Ask a doctor before use if you have</b> ■ deep or puncture wounds ■ animal bites ■ serious burns ▶			

Add image transcription here...

## Package Labeling 14g

NO  
COLOR

**Drug Facts** (continued)

**Other information** ■ store at 15 to 30°C (59 to 86°F)  
■ **Tamper Evident:** DO NOT USE IF FLAPS ON CARTON ARE NOT SEALED OR MISSING

**Inactive ingredient** petrolatum



PL33104B  
V01  
NO  
COLOR

DISTRIBUTED BY  
Taisho Pharmaceutical California Inc. 3528 Torrance Blvd, Suite #320 Torrance, CA 90503 U.S.A.



VITACILINA

FIRST AID ANTIBIOTIC

NDC 81929-003-02

**VITACILINA**<sup>®</sup>

NET WT 0.5 OZ (14 g)

NO  
COLOR

FIRST AID ANTIBIOTIC

NDC 81929-003-02

**VITACILINA**<sup>®</sup>

NET WT 0.5 OZ (14 g)

NO  
COLOR

GLUE FLAP  
NO COLOR / TEXT  
NO COATING

**Drug Facts**

Active ingredient (in each gram)	Purpose
Bacitracin zinc 400 units.....	First aid antibiotic
Neomycin sulfate 3.5 mg.....	First aid antibiotic
Polymyxin B sulfate 5,000 units.....	First aid antibiotic

**Use** First aid to help prevent infection in minor cuts, scrapes, and burns

**Warnings**

For external use only.

**Do not use** ■ if you are allergic to any of the ingredients  
■ in the eyes ■ over large areas of the body

**Drug Facts** (continued)

**Ask a doctor before use if you have** ■ deep or puncture wounds ■ animal bites ■ serious burns

**Stop use and ask a doctor if** ■ you need to use longer than 1 week ■ condition persists or gets worse ■ rash or other allergic reaction develops

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away: 800-222-1222.

**Directions** ■ clean the affected area ■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily  
■ may be covered with a sterile bandage

GLUE FLAP  
NO COLOR / TEXT  
NO COATING

FIRST AID ANTIBIOTIC

**VITACILINA®**

NET WT 0.5 OZ (14 g)

<b>Drug Facts</b>		<b>Drug Facts</b> (continued)
<b>Active ingredient (in each gram)</b>	<b>Purpose</b>	gets worse ■ rash or other allergic reaction develops
Bacitracin zinc 400 units.....	First aid antibiotic	<b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away: 800-222-1222.
Neomycin sulfate 3.5 mg.....	First aid antibiotic	
Polymyxin B sulfate 5,000 units..	First aid antibiotic	
<b>Use</b>	First aid to help prevent infection in minor cuts, scrapes, and burns	<b>Directions</b> ■ clean the affected area ■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area up to 3 times daily ■ may be covered with a sterile bandage
<b>Warnings</b> For external use only.		<b>Other information</b> ■ store at 15 to 30°C (59 to 86°F)
<b>Do not use</b> ■ if you are allergic to any of the ingredients ■ in the eyes ■ over large areas of the body		<b>Inactive ingredient</b> petrolatum
<b>Ask a doctor before use if you have</b> ■ deep or puncture wounds ■ animal bites ■ serious burns		<b>DISTRIBUTED BY</b> NDC 81929-003-02 Taisho Pharmaceutical California Inc. 3528 Torrance Blvd, Suite #320 Torrance, CA 90503 U.S.A. PL33104T V01
<b>Stop use and ask a doctor if</b> ■ you need to use longer than 1 week ■ condition persists or ►		

## VITACILINA FIRST AID ANTIBIOTIC

bacitracin zinc, neomycin sulfate, polymyxin b sulfate ointment

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:81929-003
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BACITRACIN ZINC</b> (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN	400 U in 1 g
<b>NEOMYCIN SULFATE</b> (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN	3.5 mg in 1 g
<b>POLYMYXIN B SULFATE</b> (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K)	POLYMYXIN B	5000 U in 1 g

### Inactive Ingredients

Ingredient Name		Strength		
PETROLATUM (UNII: 4T6H12BN9U)				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:81929-003-01	1 in 1 CARTON	07/01/2022	
1		28 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:81929-003-02	1 in 1 CARTON	07/01/2022	
2		14 g in 1 TUBE; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M004	07/01/2022		

**Labeler** - Taisho Pharmaceutical California Inc. (603827635)

Revised: 11/2023

Taisho Pharmaceutical California Inc.