

**BENZETHONIUM CHLORIDE PLUS DYCLONINE HYDROCHLORIDE- liquid
bandage liquid
Meijer Distribution Inc**

Meijer Liquid Bandage

Active ingredients

Benzethonium chloride 0.2%
Dyclonine hydrochloride 0.75%

Purpose

First aid antiseptic
Topical analgesic

Uses

- first aid to prevent bacterial contamination in minor cuts, scrapes and burns
- for the temporary relief of pain

Warnings

For external use only.

Flammable:

Keep away from fire or flame.

Do not use

- in the eyes
- over large areas of the body
- longer than 1 week unless directed by a doctor

Ask a doctor before use if you have

- deep puncture wounds
- deep cuts
- animal bites
- serious burns

When using this product

- avoid contact with eyes

Stop use and ask a doctor if

- conditions persists or gets worse
- infection occurs
- symptoms persist for than 7 days or clear up and occur again within a few days

Keep out of reach of children.

If swallowed, get medical help or call a Poison Control Center right away.

Directions

- for adults and children 2 years of age and older
- clean affected area
- apply a small amount on the area 1 to 3 times daily and let dry
- a second coating may be applied for extra protection
- children under 2 years of age: consult a doctor

Other information

- cap bottle tightly
- store between 20° to 25°C (68° to 77°F)
- may temporarily sting upon application
- to remove, apply more Liquid Bandage and quickly wipe off

Inactive ingredients

acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

Questions

Call 1-866-964-0939

Principal Display Panel

MEDICATED

liquid bandage

Salicylic Acid

For Small Cuts & Wounds

ANTISEPTIC | INVISIBLE

Topical Analgesic & Antiseptic

Seals to Protect

NET 0.3 FL OZ (9mL)



Salicylic Acid

For Small Cuts & Wounds

ANTISEPTIC | INVISIBLE

Topical Analgesic & Antiseptic
Seals to Protect

NET 0.3 FL OZ (9 mL)

USE LIQUID BANDAGE FOR:

Blisters, chapped & cracked finger tips, hangnails, paper cuts, shaving nicks & help in preventing the formation of calluses.

GREAT FOR:

Athletes, active lifestyles, outdoor sportsmen & musicians.

53-033ME

meijer

MEDICATED

liquid bandage

For Small Cuts & Wounds

NET 0.3 FL OZ (9 mL)

NO VARNISH
AREA
(WHITE ONLY)



DIST. BY MEIJER DISTRIBUTION, INC.
GRAND RAPIDS, MI 49544
www.meijer.com
MADE IN USA WITH U.S.
AND IMPORTED PARTS

PID 369389



*This product is not manufactured or distributed by Moberg Pharma North America LLC, owner of the registered trademark New-Skin®.

Care ■ Wipe top of bottle clean after each use.

Questions? Call 1-866-964-0939

Inactive ingredients: acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

Other information ■ cap bottle tightly ■ store between 20° to 25°C (68° to 77°F) ■ may temporarily sting upon application ■ to remove, apply more liquid bandage and quickly wipe off

Directions ■ for adults and children 2 years of age and older ■ clean affected area ■ apply a small amount on the area 1 to 3 times daily and let dry ■ a second coating may be applied for extra protection ■ children under 2 years of age: consult a doctor

Warnings ■ Keep out of reach of children. If swallowed, get medical help or call a Poison Control Center right away.

Stop use and ask a doctor if ■ condition persists or gets worse ■ infection occurs ■ symptoms persist for more than 7 days or clear up and occur again within a few days

When using this product ■ avoid contact with eyes ■ animal bites ■ serious burns ■ Ask a doctor before use if you have ■ deep or puncture wounds ■ deep cuts ■ Do not use ■ in the eyes ■ over large areas of the body ■ longer than 1 week unless directed by a doctor

Flammable: Keep away from fire or flame. For external use only.

Uses ■ first aid to prevent bacterial contamination in minor cuts, scrapes and burns ■ for the temporary relief of pain

Active ingredients Benzethonium chloride 0.2%.....Topical antiseptic Dicyclanil hydrochloride 0.75%.....First aid analgesic

Drug Facts

DO NOT USE IF NECKBAND IS BROKEN



BENZETHONIUM CHLORIDE PLUS DYCLONINE HYDROCHLORIDE

liquid bandage liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-132
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZETHONIUM CHLORIDE (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	0.2 mg in 10 mL
DYCLONINE HYDROCHLORIDE (UNII: ZEC193879Q) (DYCLONINE - UNII:078A24Q300)	DYCLONINE HYDROCHLORIDE	0.75 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
ACETONE (UNII: 1364PS73AF)	
AMYL ACETATE (UNII: 92Q24NH7AS)	
CASTOR OIL (UNII: D5340Y2I9G)	
ETHYL ACETATE (UNII: 76845O8NMZ)	
PYROXYLIN (UNII: KYR8BR2X6O)	
ALCOHOL (UNII: 3K9958V90M)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-	10 mL in 1 BOTTLE, WITH APPLICATOR; Type 0: Not a	01/05/2012	10/31/2025

132-10	Combination Product	01/05/2012	10/31/2025
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	01/05/2012	10/31/2025

Labeler - Meijer Distribution Inc (006959555)

Revised: 5/2024

Meijer Distribution Inc