

NASAL DECONGESTANT- pseudoephedrine hcl tablet, film coated
Publix Super Markets Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Publix Super Markets, Inc. Nasal Decongestant Drug Facts

Active ingredient (in each tablet)

Pseudoephedrine HCl 30 mg

Purpose

Nasal decongestant

Uses

- temporarily relieves sinus congestion and pressure
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

When using this product

do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

adults and children 12 years and over	<ul style="list-style-type: none">• take 2 tablets every 4 to 6 hours• do not take more than 8 tablets in 24 hours
children ages 6 to 11 years	<ul style="list-style-type: none">• take 1 tablet every 4 to 6 hours• do not take more than 4 tablets in 24 hours
children under 6 years	do not use this product in children under 6 years of age

Other information

- each tablet contains: calcium 20 mg
- store at 20°-25° C (68°-77° F)
- do not use if blister unit is broken or torn

Inactive ingredients

carnauba wax, dibasic calcium phosphate dihydrate, FD&C red no. 40 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, silicon dioxide, titanium dioxide

Principal Display Panel

NON-DROWSY – MAXIMUM STRENGTH

nasal decongestant

PSEUDOEPHEDRINE HYDROCHLORIDE 30mg

nasal & sinus congestion

sinus pressure + congestion

48 TABLETS

ACTUAL SIZE

Compare to Active Ingredient in Sudafed® Congestion

NON-DROWSY • MAXIMUM STRENGTH



nasal decongestant

PSEUDOEPHEDRINE HYDROCHLORIDE 30mg



NDC 56062-432-67

NON-DROWSY • MAXIMUM STRENGTH

nasal decongestant

PSEUDOEPHEDRINE HYDROCHLORIDE 30mg



- nasal & sinus congestion
- sinus pressure + congestion

48 TABLETS



ACTUAL SIZE

*Compare to the Active Ingredient in Sudafed® Congestion



Important: Read all product information before using. Keep this box for important information.

Drug Facts

Active Ingredient (In each tablet)	Purpose
Pseudoephedrine HCl 30 mg	Nasal decongestant

Uses

- temporarily relieves sinus congestion and pressure
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Drug Facts (continued)

Directions

adults and children 12 years and over	take 2 tablets every 4 to 6 hours do not take more than 8 tablets in 24 hours
children ages 6 to 11 years	take 1 tablet every 4 to 6 hours do not take more than 4 tablets in 24 hours
children under 6 years	do not use this product in children under 6 years of age

Other Information

- each tablet contains: calcium 20 mg
- store at 20°-25°C (68°-77°F)
- do not use if blister unit is broken or torn

Inactive Ingredients

camellia wax, dibasic calcium phosphate dihydrate, FD&C red no. 40 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyorbate 80, silicon dioxide, titanium dioxide

*This product is not manufactured or distributed by the owner of the registered trademark Sudafed®.

DISTRIBUTED BY
PUBLIX SUPER MARKETS, INC.,
3300 PUBLIX CORPORATE PARKWAY
LAKE WOOD, FL 33811
publix.com

PUBLIX GUARANTEE:
COMPLETE SATISFACTION OR YOUR MONEY BACK

43267 13 C3

OPEN OTHER END

CONVENIENT RECLOSING TAB





NASAL DECONGESTANT

pseudoephedrine hcl tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:56062-432
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	

Product Characteristics

Color	RED	Score	no score
Shape	ROUND (convex)	Size	7mm
Flavor		Imprint Code	L432
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:56062-432-67	2 in 1 CARTON	10/12/2000	
1		24 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:56062-432-80	4 in 1 CARTON	04/18/2002	
2		24 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:56062-432-62	1 in 1 CARTON	09/09/1993	
3		24 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	09/09/1993	

Labeler - Publix Super Markets Inc (006922009)

Revised: 11/2019

Publix Super Markets Inc