

WAL-ZAN 75- ranitidine tablet usp, 75mg tablet
Walgreens Company

Drug Facts

Active ingredient (in each tablet)

Ranitidine 75 mg (as ranitidine hydrochloride, USP 84 mg)

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

Pregnancy/Breastfeeding

If pregnant or breast-feeding, ask a health professional before use.

Keep Out of Reach of Children

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water
 - to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

Other information

- this product is sodium and sugar free
- Blister: do not use if individual blister unit is open or torn
Bottle: do not use if printed foil under bottle cap is open or torn
- avoid excessive heat or humidity
- store at 20°-25°C (68°-77°F)

Inactive ingredients

FD&C red #40 aluminum lake, hypromellose, iron oxide black, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide.

Question Section

Questions? call 1-888-375-3784

Read the directions, consumer information leaflet and warnings before use. Keep the carton. It contains important information.

Principal Display Panel

COATING
FREE AREA

150058072



Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts

Active ingredient (in each tablet) Purpose
Ranitidine 75 mg (as ranitidine hydrochloride, USP 84 mg) Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

- Warning alert:** Do not use if you are allergic to ranitidine or other acid reducers.
- Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have frequent chest pain

- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with dizziness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or fainting

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- pregnant or breastfeeding, ask a health professional before use.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 12 years and over:
 - to relieve symptoms, swallow 1 tablet with a glass of water
 - to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn
 - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

Other information

- this product is sodium and sugar free
- do not use if cap or printed foil under cap is open or torn
- avoid excessive heat or humidity
- store at 20°-25°C (68°-77°F)

Inactive ingredients

FD&C red #40, aluminum lake, hydroxypropyl methylcellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide.

Questions? Call 1-888-675-3784

IMPORTANT: Read the directions, consumer information booklet and warnings before use. Keep the cap on. Contains important information. **Not for use in children.** Walgreens Pharmacist Survey Study, November 2012. This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, Inc., distributor of ZANTAC 75® tablets. ZANTAC 75® is a registered trademark of Boehringer Ingelheim Pharmaceuticals, Inc.

Well at Walgreens

WALGREENS PHARMACY

NON-PRESCRIPTION STRENGTH

Wal-Zan® 75

Ranitidine Tablets, USP, 75 mg / Acid Reducer

BONUS SIZE BUY 30 & GET 15 FREE

Well at Walgreens

WALGREENS PHARMACY

NON-PRESCRIPTION STRENGTH

Wal-Zan® 75

Ranitidine Tablets, USP, 75 mg / Acid Reducer

- Prevents & relieves heartburn associated with acid indigestion & sour stomach



Actual Size

45 TABLETS (30 + 15 FREE)

Compare to Zantac 75® active ingredient*

WALGREENS PHARMACY, THE WALGREENS COMPANY, 100% SATISFACTION GUARANTEED

ITEM 379342 W10024-1114-L



COATING FREE AREA

LOT

EXP



Well at Walgreens

NON-PRESCRIPTION STRENGTH

Wal-Zan® 75

Ranitidine Tablets, USP, 75 mg / Acid Reducer

- Prevents & relieves heartburn associated with acid indigestion & sour stomach

45 TABLETS (30 + 15 FREE)



150058072

COATING
FREE AREA



Ranitidine Tablets USP, 75 mg - Container carton label



WAL-ZAN 75

ranitidine tablet usp, 75mg tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0131(NDC:55111-131)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ranitidine hydrochloride (UNII: BK76465IHM) (ranitidine - UNII:884KT10YB7)	ranitidine	75 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
magnesium stearate (UNII: 70097M6I30)	
cellulose, microcrystalline (UNII: OP1R32D61U)	

Polyethylene Glycol, Unspecified (UNII: 3WJQ0SDW1A)

ferrosoferric oxide (UNII: XM0M87F357)

titanium dioxide (UNII: 15FIX9V2JP)

Product Characteristics

Color	PINK	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	P75
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0131-30	1 in 1 CARTON	01/06/2014	
1		30 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0363-0131-80	1 in 1 CARTON	01/06/2014	
2		80 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:0363-0131-33	1 in 1 CARTON	01/06/2014	
3		45 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075294	01/06/2014	

Labeler - Walgreens Company (008965063)

Revised: 12/2019

Walgreens Company