

**ANTI-DIARRHEAL- loperamide hcl tablet, coated**  
**First Aid Direct**

-----  
**248 - Health A2Z Anti-Diarrheal**

**Active Ingredient (in each caplet)**

Loperamide HCl 2mg

**Purpose**

Anti-diarrheal

**Use**

controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reactions to Loperamide HCl.

**Heart alert:** Taking more than directed can cause serious heart problems or death.

**Do not use**

- if you have bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease

**Ask a doctor or pharmacist before use if you are**

taking antibiotics

**When using this product**

tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging.

These may be signs of a serious condition.

**If pregnant or breast feeding,**

**If pregnant or breast feeding,** ask a health professional before use.

**Keep out of reach of children**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222) .

### **Directions**

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.
- **adults and children 12 years and over:** 2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
- **children 9-11 years (60-95 lbs):** 1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
- **children 6-8 years (48-59 lbs):** 1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
- **children 2-5 years (34 to 47 lbs):** ask a doctor
- **children under 2 years (up to 33 lbs):** do not use

### **Other information**

- store at 20°-25°C (68°-77° F)
- avoid excessive heat above 40°C (104°F)
- do not use if blister unit is broken or torn


### **Inactive ingredients**

corn starch, D&C yellow #10 aluminum lake, dicalcium phosphate dihydrate, FD&C blue #1 brilliant blue lake, magnesium stearate, microcrystalline cellulose, silica gel

### **Questions**

1-800-327-2704

### **Principal Display Panel**



### Anti-Diarrheal

## ANTI-DIARRHEAL

loperamide hcl tablet, coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:53117-248
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

### Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	

### Product Characteristics

<b>Color</b>	green	<b>Score</b>	2 pieces
<b>Shape</b>	CAPSULE	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;375
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53117-248-86	2 in 1 CARTON	05/03/2005	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076497	05/03/2005	

**Labeler** - First Aid Direct (962283128)

Revised: 9/2019

First Aid Direct