


**RETIN-A- tretinoin cream**  
**RETIN-A- tretinoin gel**  
**Bausch Health US, LLC**

-----  
**RETIN-A®**  
**(tretinoin)**  
**Cream • Gel**

**For Topical Use Only**

**DESCRIPTION**

RETIN-A Gel and Cream, containing tretinoin, are used for the topical treatment of acne vulgaris. RETIN-A Gel contains tretinoin (retinoic acid, vitamin A acid) in either of two strengths, 0.025% or 0.01% by weight, in a gel vehicle of butylated hydroxytoluene, hydroxypropyl cellulose and alcohol (denatured with *tert*-butyl alcohol and brucine sulfate) 90% w/w. RETIN-A (tretinoin) Cream contains tretinoin in one of three strengths, 0.1%, 0.05%, or 0.025% by weight, in a hydrophilic cream vehicle of stearic acid, isopropyl myristate, polyoxyl 40 stearate, stearyl alcohol, xanthan gum, sorbic acid, butylated hydroxytoluene, and purified water. Chemically, tretinoin is *all-trans*-retinoic acid and has the following structure:



**CLINICAL PHARMACOLOGY**

Although the exact mode of action of tretinoin is unknown, current evidence suggests that topical tretinoin decreases cohesiveness of follicular epithelial cells with decreased microcomedo formation. Additionally, tretinoin stimulates mitotic activity and increased turnover of follicular epithelial cells causing extrusion of the comedones.

**INDICATIONS AND USAGE**

RETIN-A is indicated for topical application in the treatment of acne vulgaris. The safety and efficacy of the long-term use of this product in the treatment of other disorders have not been established.

**CONTRAINDICATIONS**

Use of the product should be discontinued if hypersensitivity to any of the ingredients is noted.

## **PRECAUTIONS**

### **General**

If a reaction suggesting sensitivity or chemical irritation occurs, use of the medication should be discontinued. Exposure to sunlight, including sunlamps, should be minimized during the use of RETIN-A, and patients with sunburn should be advised not to use the product until fully recovered because of heightened susceptibility to sunlight as a result of the use of tretinoin. Patients who may be required to have considerable sun exposure due to occupation and those with inherent sensitivity to the sun should exercise particular caution. Use of sunscreen products and protective clothing over treated areas is recommended when exposure cannot be avoided. Weather extremes, such as wind or cold, also may be irritating to patients under treatment with tretinoin.

RETIN-A (tretinoin) acne treatment should be kept away from the eyes, the mouth, angles of the nose, and mucous membranes. Topical use may induce severe local erythema and peeling at the site of application. If the degree of local irritation warrants, patients should be directed to use the medication less frequently, discontinue use temporarily, or discontinue use altogether. Tretinoin has been reported to cause severe irritation on eczematous skin and should be used with utmost caution in patients with this condition.

### **Drug Interactions**

Concomitant topical medication, medicated or abrasive soaps and cleansers, soaps and cosmetics that have a strong drying effect, and products with high concentrations of alcohol, astringents, spices or lime should be used with caution because of possible interaction with tretinoin. Particular caution should be exercised in using preparations containing sulfur, resorcinol, or salicylic acid with RETIN-A. It also is advisable to “rest” a patient’s skin until the effects of such preparations subside before use of RETIN-A is begun.

### **Carcinogenesis, Mutagenesis, Impairment to Fertility**

In a 91-week dermal study in which CD-1 mice were administered 0.017% and 0.035% formulations of tretinoin, cutaneous squamous cell carcinomas and papillomas in the treatment area were observed in some female mice. A dose-related incidence of liver tumors in male mice was observed at those same doses. The maximum systemic doses associated with the administered 0.017% and 0.035% formulations are 0.5 and 1.0 mg/kg/day, respectively. These doses are two and four times the maximum human systemic dose, when adjusted for total body surface area. The biological significance of these findings is not clear because they occurred at doses that exceeded the dermal maximally tolerated dose (MTD) of tretinoin and because they were within the background natural occurrence rate for these tumors in this strain of mice. There was no evidence of carcinogenic potential when 0.025 mg/kg/day of tretinoin was administered topically to mice (0.1 times the maximum human systemic dose, adjusted for total body surface area). For purposes of comparisons of the animal exposure to systemic human exposure, the maximum human systemic dose is defined as 1 gram of 0.1% RETIN-A applied daily to a 50 kg person (0.02 mg tretinoin/kg body weight).

Studies in hairless albino mice suggest that concurrent exposure to tretinoin may enhance the tumorigenic potential of carcinogenic doses of UVB and UVA light from a

solar simulator. This effect has been confirmed in a later study in pigmented mice, and dark pigmentation did not overcome the enhancement of photocarcinogenesis by 0.05% tretinoin. Although the significance of these studies to humans is not clear, patients should minimize exposure to sunlight or artificial ultraviolet irradiation sources.

The mutagenic potential of tretinoin was evaluated in the Ames assay and in the in vivo mouse micronucleus assay, both of which were negative.

In dermal Segment I fertility studies of another tretinoin formulation in rats, slight (not statistically significant) decreases in sperm count and motility were seen at 0.5 mg/kg/day (4 times the maximum human systemic dose adjusted for total body surface area) and slight (not statistically significant) increases in the number and percent of nonviable embryos in females treated with 0.25 mg/kg/day (2 times the maximum human systemic dose adjusted for total body surface area) and above were observed. A dermal Segment III study with RETIN-A has not been performed in any species. In oral Segment I and Segment III studies in rats with tretinoin, decreased survival of neonates and growth retardation were observed at doses in excess of 2 mg/kg/day (16 times the human topical dose adjusted for total body surface area).

## **Pregnancy**

### Teratogenic Effects

Oral tretinoin has been shown to be teratogenic in rats, mice, hamsters, and subhuman primates. It was teratogenic and fetotoxic in Wistar rats when given orally or topically in doses greater than 1 mg/kg/day (8 times the maximum human systemic dose adjusted for total body surface area). However, variations in teratogenic doses among various strains of rats have been reported. In the cynomolgus monkey, which metabolically is closer to humans for tretinoin than the other species examined, fetal malformations were reported at doses of 10 mg/kg/day or greater, but none were observed at 5 mg/kg/day (83 times the maximum human systemic dose adjusted for total body surface area), although increased skeletal variations were observed at all doses. A dose-related increase in embryoletality and abortion was reported. Similar results have also been reported in pigtail macaques.

Topical tretinoin in animal teratogenicity tests has generated equivocal results. There is evidence for teratogenicity (shortened or kinked tail) of topical tretinoin in Wistar rats at doses greater than 1 mg/kg/day (8 times the maximum human systemic dose adjusted for total body surface area). Anomalies (humerus: short 13%, bent 6%, os parietale incompletely ossified 14%) have also been reported when 10 mg/kg/day was topically applied.

There are other reports in New Zealand White rabbits administered doses of greater than 0.2 mg/kg/day (3.3 times the maximum human systemic dose adjusted for total body surface area) of an increased incidence of domed head and hydrocephaly, typical of retinoid-induced fetal malformations in this species.

In contrast, several well-controlled animals studies have shown that dermally applied tretinoin may be fetotoxic, but not overly teratogenic in rats and rabbits at doses of 1.0 and 0.5 mg/kg/day, respectively (8 times the maximum human systemic dose adjusted for total body surface area in both species).

With widespread use of any drug, a small number of birth defect reports associated temporally with the administration of the drug would be expected by chance alone.

Thirty human cases of temporally associated congenital malformations have been reported during two decades of clinical use of RETIN-A. Although no definite pattern of teratogenicity and no causal association have been established from these cases, five of the reports describe the rare birth defect category holoprosencephaly (defects associated with incomplete midline development of the forebrain). The significance of these spontaneous reports in terms of risk to the fetus is not known.

### Nonteratogenic Effects

Topical tretinoin has been shown to be fetotoxic in rabbits when administered 0.5 mg/kg/day (8 times the maximum human systemic dose adjusted for total body surface area). Oral tretinoin has been shown to be fetotoxic, resulting in skeletal variations and increased intrauterine death in rats when administered 2.5 mg/kg/day (20 times the maximum human systemic dose adjusted for total body surface area).

There are, however, no adequate and well-controlled studies in pregnant women. Tretinoin should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

### **Nursing Mothers**

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when RETIN-A is administered to a nursing woman.

### **Pediatric Use**

Safety and effectiveness in pediatric patients below the age of 12 have not been established.

### **Geriatric Use**

Safety and effectiveness in a geriatric population have not been established. Clinical studies of RETIN-A did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger patients.

**GELS ARE FLAMMABLE.** Note: Keep away from heat and flame. Keep tube tightly closed.

### **ADVERSE REACTIONS**

The skin of certain sensitive individuals may become excessively red, edematous, blistered, or crusted. If these effects occur, the medication should either be discontinued until the integrity of the skin is restored, or the medication should be adjusted to a level the patient can tolerate. True contact allergy to topical tretinoin is rarely encountered. Temporary hyper- or hypopigmentation has been reported with repeated application of RETIN-A. Some individuals have been reported to have heightened susceptibility to sunlight while under treatment with RETIN-A. To date, all adverse effects of RETIN-A have been reversible upon discontinuance of therapy (see **DOSAGE AND ADMINISTRATION**).

**To report SUSPECTED ADVERSE REACTIONS, contact Bausch Health US, LLC at 1-800-321-4576 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

## OVERDOSAGE

If medication is applied excessively, no more rapid or better results will be obtained and marked redness, peeling, or discomfort may occur. Oral ingestion of the drug may lead to the same side effects as those associated with excessive oral intake of Vitamin A.

## DOSAGE AND ADMINISTRATION

RETIN-A Gel or Cream should be applied once a day, before retiring, to the skin where acne lesions appear, using enough to cover the entire affected area lightly. Gel: Excessive application results in “pilling” of the gel, which minimizes the likelihood of overapplication by the patient.

Application may cause a transitory feeling of warmth or slight stinging. In cases where it has been necessary to temporarily discontinue therapy or to reduce the frequency of application, therapy may be resumed or frequency of application increased when the patients become able to tolerate the treatment.

Alterations of vehicle, drug concentration, or dose frequency should be closely monitored by careful observation of the clinical therapeutic response and skin tolerance.

During the early weeks of therapy, an *apparent* exacerbation of inflammatory lesions may occur. This is due to the action of the medication on deep, previously unseen lesions and should not be considered a reason to discontinue therapy.

Therapeutic results should be noticed after 2 to 3 weeks but more than 6 weeks of therapy may be required before definite beneficial effects are seen.

Once the acne lesions have responded satisfactorily, it may be possible to maintain the improvement with less frequent applications, or other dosage forms.

Patients treated with RETIN-A (tretinoin) acne treatment may use cosmetics, but the area to be treated should be cleansed thoroughly before the medication is applied(see **PRECAUTIONS**).

## HOW SUPPLIED

RETIN-A (tretinoin) is supplied as:

RETIN-A Cream			RETIN-A Gel		
NDC Code	RETIN-A Strength/ Form	RETIN-A Qty.	NDC Code	RETIN-A Strength/ Form	RETIN-A Qty.
0187-5160-20	0.025% Cream	20 g	0187-5172-15	0.01% Gel	15 g
0187-5160-45	0.025% Cream	45 g	0187-5172-45	0.01% Gel	45 g
0187-5162-20	0.05% Cream	20 g	0187-5170-15	0.025% Gel	15 g
0187-5162-45	0.05% Cream	45 g	0187-5170-	0.025% Gel	45 g

0187-5164-20	0.1% Cream	20 g
0187-5164-45	0.1% Cream	45 g

---

**Storage Conditions:**

RETIN-A Gel, 0.025% and 0.01%: store below 86°F. RETIN-A Cream, 0.1%, 0.05%, and 0.025%: store below 80°F.

**Manufactured for:**

Bausch Health US, LLC  
Bridgewater, NJ 08807 USA

**Manufactured by:**

Bausch Health Companies Inc.  
Laval, Quebec H7L 4A8, Canada

Retin-A is a trademark of Bausch Health Companies Inc. or its affiliates.

© 2019 Bausch Health Companies Inc. or its affiliates

9440905

Revised: 09/2019

**PATIENT INSTRUCTIONS****RETIN-A®  
(tretinoin)****Cream • Gel****For Topical Use Only****PATIENT INSTRUCTIONS****Acne Treatment****IMPORTANT****Read Directions Carefully Before Using**

**THIS LEAFLET TELLS YOU ABOUT RETIN-A (TRETINOIN) ACNE TREATMENT AS PRESCRIBED BY YOUR PHYSICIAN. THIS PRODUCT IS TO BE USED ONLY ACCORDING TO YOUR DOCTOR'S INSTRUCTIONS, AND IT SHOULD NOT BE APPLIED TO OTHER AREAS OF THE BODY OR TO OTHER GROWTHS OR LESIONS. THE LONG-TERM SAFETY AND EFFECTIVENESS OF THIS PRODUCT IN OTHER DISORDERS HAVE NOT BEEN EVALUATED. IF YOU HAVE ANY QUESTIONS, BE SURE TO ASK YOUR DOCTOR.**

**WARNINGS**

**RETIN-A GELS ARE FLAMMABLE. AVOID FIRE, FLAME OR SMOKING DURING USE.** Keep out of reach of children. Keep tube tightly closed. Do not expose to heat or store at temperatures above 120°F (49°C).

## PRECAUTIONS

**The effects of the sun on your skin.** As you know, overexposure to natural sunlight or the artificial sunlight of a sunlamp can cause sunburn. Overexposure to the sun over many years may cause premature aging of the skin and even skin cancer. The chance of these effects occurring will vary depending on skin type, the climate and the care taken to avoid overexposure to the sun. Therapy with RETIN-A may make your skin more susceptible to sunburn and other adverse effects of the sun, so unprotected exposure to natural or artificial sunlight should be minimized.

**Laboratory findings.** *When laboratory mice are exposed to artificial sunlight, they often develop skin tumors. These sunlight-induced tumors may appear more quickly and in greater number if the mouse is also topically treated with the active ingredient in RETIN-A, tretinoin. In some studies, under different conditions, however, when mice treated with tretinoin were exposed to artificial sunlight, the incidence and rate of development of skin tumors was reduced. There is no evidence to date that tretinoin alone will cause the development of skin tumors in either laboratory animals or humans. However, investigations in this area are continuing.*

**Use caution in the sun.** When outside, even on hazy days, areas treated with RETIN-A should be protected. An effective sunscreen should be used any time you are outside (consult your physician for a recommendation of an SPF level which will provide you with the necessary high level of protection). For extended sun exposure, protective clothing, like a hat, should be worn. Do not use artificial sunlamps while you are using RETIN-A. If you do become sunburned, stop your therapy with RETIN-A until your skin has recovered.

**Avoid excessive exposure to wind or cold.** Extremes of climate tend to dry or burn normal skin. Skin treated with RETIN-A may be more vulnerable to these extremes. Your physician can recommend ways to manage your acne treatment under such conditions.

**Possible problems.** The skin of certain sensitive individuals may become excessively red, swollen, blistered or crusted. If you are experiencing severe or persistent irritation, discontinue the use of RETIN-A and consult your physician.

There have been reports that, in some patients, areas treated with RETIN-A developed a temporary increase or decrease in the amount of skin pigment (color) present. The pigment in these areas returned to normal either when the skin was allowed to adjust to RETIN-A or therapy was discontinued.

**Use other medications only on your physician's advice.** Only your physician knows which other medications may be helpful during treatment and will recommend them to you if necessary. Follow the physician's instructions carefully. In addition, you should avoid preparations that may dry or irritate your skin. These preparations may include certain astringents, toiletries containing alcohol, spices or lime, or certain medicated soaps, shampoos and hair permanent solutions. Do not allow anyone else to use this medication.

Do not use other medications with RETIN-A which are not recommended by your doctor. The medications you have used in the past might cause unnecessary redness or peeling.

**If you are pregnant, think you are pregnant or are nursing an infant:** No

studies have been conducted in humans to establish the safety of RETIN-A in pregnant women. If you are pregnant, think you are pregnant, or are nursing a baby, consult your physician before using this medication.


### **AND WHILE YOU'RE ON RETIN-A THERAPY**

*Use a mild, non-medicated soap. Avoid frequent washings and harsh scrubbing.* Acne isn't caused by dirt, so no matter how hard you scrub, you can't wash it away. Washing too frequently or scrubbing too roughly may at times actually make your acne worse. Wash your skin gently with a mild, bland soap. Two or three times a day should be sufficient. Pat skin dry with a towel. Let the face dry 20 to 30 minutes before applying RETIN-A. Remember, excessive irritation such as rubbing, too much washing, use of other medications not suggested by your physician, etc., may worsen your acne.

### **HOW TO USE RETIN-A (TRETINOIN)**

To get the best results with RETIN-A therapy, it is necessary to use it properly. Forget about the instructions given for other products and the advice of friends. Just stick to the special plan your doctor has laid out for you and be patient. Remember, when RETIN-A is *used properly*, many users see improvement by 12 weeks. **AGAIN, FOLLOW INSTRUCTIONS - BE PATIENT - DON'T START AND STOP THERAPY ON YOUR OWN - IF YOU HAVE QUESTIONS, ASK YOUR DOCTOR.**

To help you use the medication correctly, keep these simple instructions in mind.



- Apply RETIN-A once daily before bedtime, or as directed by your physician. Your physician may advise, especially if your skin is sensitive, that you start your therapy by applying RETIN-A every other night. First, wash with a mild soap and dry your skin gently. **WAIT 20 TO 30 MINUTES BEFORE APPLYING MEDICATION;** it is important for skin to be completely dry in order to minimize possible irritation.
- It is better not to use more than the amount suggested by your physician or to apply more frequently than instructed. Too much may irritate the skin, waste medication and won't give faster or better results.



- Keep the medication away from the corners of the nose, mouth, eyes and open wounds. *Spread away from these areas when applying.*
- *Cream:* Squeeze about a half inch or less of medication onto the fingertip. While that should be enough for your whole face, after you have some experience with the medication, you may find you need slightly more or less to do the job. The medication should become invisible almost immediately. If it is still visible, you are using too much. Cover the affected area lightly with RETIN-A (tretinoin) Cream by first dabbing it on your forehead, chin and both cheeks, then spreading it over the entire affected area. Smooth gently into the skin.
- *Gel:* Squeeze about a half inch or less of medication onto the fingertip. While that should be enough for your whole face, after you have some experience with the medication, you may find you need slightly more or less to do the job. The medication should become invisible almost immediately. If it is still visible, or if dry flaking occurs from the gel **within a minute or so**, you are using too much. Cover the affected area lightly with RETIN-A (tretinoin) Gel by first dabbing it on your forehead, chin and both cheeks, then spreading it over the entire affected area. Smooth gently into the skin.
- It is recommended that you apply a moisturizer or a moisturizer with sunscreen that will not aggravate your acne (noncomedogenic) every morning after you wash.

## WHAT TO EXPECT WITH YOUR NEW TREATMENT

RETIN-A works deep inside your skin and this takes time. You cannot make RETIN-A work any faster by applying more than one dose each day, but an excess amount of RETIN-A may irritate your skin. Be patient.

There may be some discomfort or peeling during the early days of treatment. Some patients also notice that their skin begins to take on a blush.

These reactions do not happen to everyone. If they do, it is just your skin adjusting to RETIN-A and this usually subsides within 2 to 4 weeks. These reactions can usually be minimized by following instructions carefully. Should the effects become excessively troublesome, consult your doctor.

BY 3 TO 6 WEEKS, some patients notice an appearance of new blemishes (papules and pustules). At this stage it is *important to continue* using RETIN-A.

If RETIN-A is going to have a beneficial effect for you, you should notice a continued improvement in your appearance after 6 to 12 weeks of therapy. Don't be discouraged if you see no immediate improvement. Don't stop treatment at the first signs of improvement.

Once your acne is under control you should continue regular application of RETIN-A until your physician instructs otherwise.

## IF YOU HAVE QUESTIONS

All questions of a medical nature should be taken up with your doctor. For more information about RETIN-A (tretinoin), call our toll-free number: 800-321-4576. Call between 9:00 a.m. and 3:00 p.m. Eastern Time, Monday through Friday.

**Storage Conditions:** RETIN-A Gel, 0.025% and 0.01%: store below 86°F. RETIN-A Cream, 0.1%, 0.05%, and 0.025%: store below 80°F.

**Manufactured for:**

Bausch Health US, LLC  
Bridgewater, NJ 08807 USA

**Manufactured by:**

Bausch Health Companies Inc.  
Laval, Quebec H7L 4A8, Canada

Retin-A is a trademark of Bausch Health Companies Inc. or its affiliates.

© 2019 Bausch Health Companies Inc. or its affiliates

9440905

Revised: 09/2019

**PRINCIPAL DISPLAY PANEL - 0.025% 45 g Cream Carton**

**NDC** 0187-5160-45

**Retin-A® Cream**  
**(tretinoin) 0.025%**

**Each gram contains:** Tretinoin 0.025%, stearic acid, isopropyl myristate, polyoxyl 40 stearate, stearyl alcohol, xanthan gum, sorbic acid, butylated hydroxytoluene, and purified water.

**For Topical Use Only**


**Rx only**

**Usual Dosage: See package insert.**

**Store below 80°F.**

**WARNING: Keep out of reach of children.**

**Net Wt. 45 g**



## PRINCIPAL DISPLAY PANEL - 0.05% 45 g Cream Carton

NDC 0187-5162-45

### Retin-A<sup>®</sup> Cream (tretinoin) 0.05%

**Each gram contains:** Tretinoin 0.05%, stearic acid, isopropyl myristate, polyoxyl 40 stearate, stearyl alcohol, xanthan gum, sorbic acid, butylated hydroxytoluene, and purified water.

#### For Topical Use Only

**Rx only**

**Usual Dosage: See package insert.**


**Store below 80°F.**

**WARNING: Keep out of reach of children.**

**Net Wt. 45 g**

9394604

50104835E



## PRINCIPAL DISPLAY PANEL - 0.1% 45 g Cream Carton

NDC 0187-5164-45

**Retin-A<sup>®</sup> Cream  
(tretinoin) 0.1%**

**Each gram contains:** Tretinoin 0.1%, stearic acid, isopropyl myristate, polyoxyl 40 stearate, stearyl alcohol, xanthan gum, sorbic acid, butylated hydroxytoluene, and purified water.

**For Topical Use Only**

**Rx only**

**Usual Dosage: See package insert.**

**Store below 80°F.**

**WARNING: Keep out of reach of children.**

**Net Wt. 45 g**



## PRINCIPAL DISPLAY PANEL - 0.01% 45 g Gel Carton

NDC 0187-5172-45

**Retin-A® Gel**  
**(tretinoin) 0.01%**

**Each gram contains:** Tretinoin 0.01%, butylated hydroxytoluene, hydroxypropyl cellulose and alcohol, (denatured with *tert*-butyl alcohol and brucine sulfate) 90% w/w.

**For Topical Use Only**


**Rx only**

**Usual Dosage: See package insert.**

**Store below 86°F.**

**WARNING: Keep out of reach of children.**

**Net Wt. 45 g**



## PRINCIPAL DISPLAY PANEL - 0.025% 45 g Gel Carton

NDC 0187-5170-45

**Retin-A® Gel  
(tretinoin) 0.025%**

**Each gram contains:** Tretinoin 0.025%, butylated hydroxytoluene, hydroxypropyl cellulose and alcohol (denatured with *tert*-butyl alcohol and brucine sulfate) 90% w/w.

**For Topical Use Only**

**Rx only**


**Usual Dosage:** See package insert.

**Store below 86°F.**

**WARNING:** Keep out of reach of children.

**Net Wt. 45 g**





## RETIN-A

tretinoin cream

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0187-5160
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>Tretinoin</b> (UNII: 5688UTC01R) (Tretinoin - UNII:5688UTC01R)	Tretinoin	0.25 mg in 1 g

### Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>Stearic Acid</b> (UNII: 4ELV7Z65AP)	
<b>Isopropyl Myristate</b> (UNII: 0RE8K4LNJS)	
<b>Polyoxyl 40 Stearate</b> (UNII: 13A4J4NH9I)	
<b>Stearyl Alcohol</b> (UNII: 2KR89I4H1Y)	
<b>Xanthan Gum</b> (UNII: TTV12P4NEE)	
<b>Sorbic Acid</b> (UNII: X045VJ989B)	

<b>Butylated Hydroxytoluene</b> (UNII: 1P9D0Z171K)				
<b>Water</b> (UNII: 059QF0KO0R)				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0187-5160-20	1 in 1 CARTON	10/30/1988	
1		20 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:0187-5160-45	1 in 1 CARTON	10/30/1988	
2		45 g in 1 TUBE; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
<b>Marketing Category</b>		<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
NDA		NDA019049	10/30/1988	

## RETIN-A

tretinoin cream

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0187-5162
Route of Administration	TOPICAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
Tretinoin (UNII: 5688UTC01R) (Tretinoin - UNII:5688UTC01R)		Tretinoin	0.5 mg in 1 g
Inactive Ingredients			
Ingredient Name			Strength
Stearic Acid (UNII: 4ELV7Z65AP)			
Isopropyl Myristate (UNII: 0RE8K4LNJS)			
Polyoxyl 40 Stearate (UNII: 13A4J4NH9I)			
Stearyl Alcohol (UNII: 2KR89I4H1Y)			
Xanthan Gum (UNII: TTV12P4NEE)			
Sorbic Acid (UNII: X045WJ989B)			
Butylated Hydroxytoluene (UNII: 1P9D0Z171K)			
Water (UNII: 059QF0KO0R)			



Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0187-5162-20	1 in 1 CARTON	07/30/1974	
1		20 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:0187-5162-45	1 in 1 CARTON	07/30/1974	
2		45 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA017522	07/30/1974	

RETIN-A			
tretinoin cream			

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0187-5164
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
Tretinoin (UNII: 5688UTC01R) (Tretinoin - UNII:5688UTC01R)	Tretinoin	1 mg in 1 g

Inactive Ingredients	
Ingredient Name	Strength
Stearic Acid (UNII: 4ELV7Z65AP)	
Isopropyl Myristate (UNII: 0RE8K4LNJS)	
Polyoxyl 40 Stearate (UNII: 13A4J4NH9I)	
Stearyl Alcohol (UNII: 2KR89I4H1Y)	
Xanthan Gum (UNII: TTV12P4NEE)	
Sorbic Acid (UNII: X045WJ989B)	
Butylated Hydroxytoluene (UNII: 1P9D0Z171K)	
Water (UNII: 059QF0KO0R)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0187-5164-20	1 in 1 CARTON	04/30/1973	

1		20 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:0187-5164-45	1 in 1 CARTON	04/30/1973	
2		45 g in 1 TUBE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA017340	04/30/1973	

### RETIN-A

tretinoin gel

#### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0187-5172
Route of Administration	TOPICAL		

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Tretinoin</b> (UNII: 5688UTC01R) (Tretinoin - UNII:5688UTC01R)	Tretinoin	0.1 mg in 1 g

#### Inactive Ingredients

Ingredient Name	Strength
<b>Butylated Hydroxytoluene</b> (UNII: 1P9D0Z171K)	
<b>Alcohol</b> (UNII: 3K9958V90M)	
<b>TERT-BUTYL ALCOHOL</b> (UNII: MD83SFE959)	
<b>BRUCINE SULFATE HEPTAHYDRATE</b> (UNII: O900CDH328)	
<b>HYDROXYPROPYL CELLULOSE, UNSPECIFIED</b> (UNII: 9XZ8H6N6OH)	

#### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0187-5172-15	1 in 1 CARTON	07/30/1979	
1		15 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:0187-5172-45	1 in 1 CARTON	07/30/1979	
2		45 g in 1 TUBE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA017955	07/30/1979	

## RETIN-A

tretinoin gel

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0187-5170
Route of Administration	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Tretinoin</b> (UNII: 5688UTC01R) (Tretinoin - UNII:5688UTC01R)	Tretinoin	0.25 mg in 1 g

### Inactive Ingredients

Ingredient Name	Strength
<b>Butylated Hydroxytoluene</b> (UNII: 1P9D0Z171K)	
<b>Alcohol</b> (UNII: 3K9958V90M)	
<b>TERT-BUTYL ALCOHOL</b> (UNII: MD83SFE959)	
<b>BRUCINE SULFATE HEPTAHYDRATE</b> (UNII: O900CDH328)	
<b>HYDROXYPROPYL CELLULOSE, UNSPECIFIED</b> (UNII: 9XZ8H6N6OH)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0187-5170-15	1 in 1 CARTON	07/30/1975	
1		15 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:0187-5170-45	1 in 1 CARTON	07/30/1975	
2		45 g in 1 TUBE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA017579	07/30/1975	

**Labeler -** Bausch Health US, LLC (831922468)

## Establishment

Name	Address	ID/FEI	Business Operations
Bausch Health Companies Inc.		245141858	MANUFACTURE(0187-5160, 0187-5162, 0187-5164, 0187-5172, 0187-5170)

Revised: 9/2019

Bausch Health US, LLC