

# **CALCIUM ACETATE- calcium acetate tablet**

## **Chartwell RX LLC**

-----

### **HIGHLIGHTS OF PRESCRIBING INFORMATION**

**These highlights do not include all the information needed to use CALCIUM ACETATE TABLETS safely and effectively. See full prescribing information for CALCIUM ACETATE TABLETS.**

### **CALCIUM ACETATE tablets, for oral use**

**Initial U.S. Approval: 1990**

#### **INDICATIONS AND USAGE**

- Calcium acetate is a phosphate binder indicated for the reduction of serum phosphorus in patients with end stage renal disease. ( 1)

#### **DOSAGE AND ADMINISTRATION**

- Starting dose is 2 tablets with each meal. ( 2)
- Titrate the dose every 2 to 3 weeks until acceptable serum phosphorus level is reached. Most patients require 3 to 4 tablets with each meal. ( 2)

#### **DOSAGE FORMS AND STRENGTHS**

- Tablets: 667 mg calcium acetate per tablet. ( 3)

#### **CONTRAINDICATIONS**

- Hypercalcemia. ( 4)

#### **WARNINGS AND PRECAUTIONS**

- Treat mild hypercalcemia by reducing or interrupting calcium acetate tablets and Vitamin D. Severe hypercalcemia may require hemodialysis and discontinuation of calcium acetate tablets. ( 5.1)
- Hypercalcemia may aggravate digitalis toxicity. ( 5.2)

#### **ADVERSE REACTIONS**

- The most common (> 10%) adverse reactions are hypercalcemia, nausea and vomiting. ( 6.1)
- In clinical studies, patients have occasionally experienced nausea during calcium acetate therapy. ( 6)

**To report SUSPECTED ADVERSE REACTIONS, contact Chartwell RX, LLC., at 845-232-1683 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch)**

#### **DRUG INTERACTIONS**

- Calcium acetate may decrease the bioavailability of tetracyclines or fluoroquinolones. ( 7)
- When clinically significant drug interactions are expected, administer the drug at least one hour before or at least three hours after calcium acetate or consider monitoring blood levels of the drug. ( 7)

**See 17 for PATIENT COUNSELING INFORMATION.**

**Revised: 5/2019**

---

## **FULL PRESCRIBING INFORMATION: CONTENTS\***

### **1 INDICATIONS AND USAGE**

## **2 DOSAGE AND ADMINISTRATION**

## **3 DOSAGE FORMS AND STRENGTHS**

## **4 CONTRAINDICATIONS**

## **5 WARNINGS AND PRECAUTIONS**

5.1 Hypercalcemia

5.2 Concomitant Use with Medications

## **6 ADVERSE REACTIONS**

6.1 Clinical Trial Experience

6.2 Postmarketing Experience

## **7 DRUG INTERACTIONS**

7.1 Ciprofloxacin

## **8 USE IN SPECIFIC POPULATIONS**

8.1 Pregnancy

8.2 Labor and Delivery

8.3 Nursing Mothers

8.4 Pediatric Use

8.5 Geriatric Use

## **10 OVERDOSAGE**

## **11 DESCRIPTION**

## **12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.2 Pharmacodynamics

## **13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment and Fertility

## **14 CLINICAL STUDIES**

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

## **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

Calcium acetate is a phosphate binder indicated to reduce serum phosphorus in patients with end stage renal disease (ESRD).

### **2 DOSAGE AND ADMINISTRATION**

The recommended initial dose of calcium acetate for the adult dialysis patient is 2 tablets with each meal. Increase the dose gradually to lower serum phosphorus levels to the target range, as long as hypercalcemia does not develop. Most patients require 3 to 4 tablets with each meal.

### **3 DOSAGE FORMS AND STRENGTHS**

Tablets: 667 mg calcium acetate per tablet.

## **4 CONTRAINDICATIONS**

Patients with hypercalcemia.

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Hypercalcemia**

Patients with end stage renal disease may develop hypercalcemia when treated with calcium, including calcium acetate . Avoid the use of calcium supplements, including calcium based nonprescription antacids, concurrently with calcium acetate.

An overdose of calcium acetate may lead to progressive hypercalcemia, which may require emergency measures. Therefore, early in the treatment phase during the dosage adjustment period, monitor serum calcium levels twice weekly. Should hypercalcemia develop, reduce the calcium acetate dosage, or discontinue the treatment, depending on the severity of hypercalcemia.

More severe hypercalcemia (Ca >12 mg/dL) is associated with confusion, delirium, stupor and coma. Severe hypercalcemia can be treated by acute hemodialysis and discontinuing calcium acetate therapy.

Mild hypercalcemia (10.5 to 11.9 mg/dL) may be asymptomatic or manifest as constipation, anorexia, nausea, and vomiting. Mild hypercalcemia is usually controlled by reducing the calcium acetate dose or temporarily discontinuing therapy. Decreasing or discontinuing Vitamin D therapy is recommended as well.

Chronic hypercalcemia may lead to vascular calcification and other soft-tissue calcification. Radiographic evaluation of suspected anatomical regions may be helpful in early detection of soft tissue calcification. The long-term effect of calcium acetate on the progression of vascular or soft tissue calcification has not been determined.

Hypercalcemia (>11 mg/dL) was reported in 16% of patients in a 3-month study of solid dose formulation of calcium acetate; all cases resolved upon lowering the dose or discontinuing treatment.

Maintain the serum calcium-phosphorus (Ca × P) product below 55 mg <sup>2</sup>/dL <sup>2</sup>.

### **5.2 Concomitant Use with Medications**

Hypercalcemia may aggravate digitalis toxicity.

## **6 ADVERSE REACTIONS**

Hypercalcemia is discussed elsewhere [*see Warnings and Precautions (5.1)*]

### **6.1 Clinical Trial Experience**

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

In clinical studies, calcium acetate has been generally well tolerated.

Calcium acetate was studied in a 3-month, open-label, non-randomized study of 98 enrolled ESRD hemodialysis patients and an alternate liquid formulation of calcium acetate was studied in a two week double-blind, placebo-controlled, cross-over study with 69 enrolled ESRD hemodialysis patients. Adverse reactions (>2% on treatment) from these trials are presented in Table 1.

**Table 1: Adverse Reactions in Patients with End-Stage renal Disease undergoing Hemodialysis**

Preferred Term	Total adverse reactions reported for calcium acetate n=167 n (%)	3-mo, open-label study of calcium acetate n=98 n (%)	Double-blind, placebo-controlled, cross-over study of liquid calcium acetate n=69	
			Calcium acetate n (%)	Placebo n (%)
Nausea	6 (3.6)	6 (6.1)	0 (0.0)	0 (0.0)
Vomiting	4 (2.4)	4 (4.1)	0 (0.0)	0 (0.0)
Hypercalcemia	21 (12.6)	16 (16.3)	5 (7.2)	0 (0.0)

Mild hypercalcemia may be asymptomatic or manifest itself as constipation, anorexia, nausea, and vomiting. More severe hypercalcemia is associated with confusion, delirium, stupor, and coma. Decreasing dialysate calcium concentration could reduce the incidence and severity of calcium acetate -induced hypercalcemia. Isolated cases of pruritus have been reported, which may represent allergic reactions.

## 6.2 Postmarketing Experience

Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to estimate their frequency or to establish a causal relationship to drug exposure.

The following additional adverse reactions have been identified during post-approval of calcium acetate: dizziness, edema, and weakness.

## 7 DRUG INTERACTIONS

The drug interaction of calcium acetate is characterized by the potential of calcium to bind to drugs with anionic functions (e.g., carboxyl, and hydroxyl groups). Calcium acetate may decrease the bioavailability of tetracyclines or fluoroquinolones via this mechanism.

There are no empirical data on avoiding drug interactions between calcium acetate and most concomitant drugs. When administering an oral medication with calcium acetate where a reduction in the bioavailability of that medication would have a clinically significant effect on its safety or efficacy, administer the drug one hour before or three hours after calcium acetate. Monitor blood levels of the concomitant drugs that have a narrow therapeutic range.

Patients taking anti-arrhythmic medications for the control of arrhythmias and anti-seizure medications for the control of seizure disorders were excluded from the clinical trials with all forms of calcium acetate.

### **7.1 Ciprofloxacin**

In a study of 15 healthy subjects, a co-administered single dose of 4 calcium acetate tablets, approximately 2.7g, decreased the bioavailability of ciprofloxacin by approximately 50%.

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

#### **Pregnancy Category C**

Calcium acetate tablets contain calcium acetate. Animal reproduction studies have not been conducted with calcium acetate, and there are no adequate and well controlled studies of calcium acetate use in pregnant women. Patients with end stage renal disease may develop hypercalcemia with calcium acetate treatment [*see Warnings and Precautions (5.1)*]. Maintenance of normal serum calcium levels is important for maternal and fetal well being. Hypercalcemia during pregnancy may increase the risk for maternal and neonatal complications such as stillbirth, preterm delivery, and neonatal hypocalcemia and hypoparathyroidism. Calcium acetate treatment, as recommended, is not expected to harm a fetus if maternal calcium levels are properly monitored during and following treatment.

### **8.2 Labor and Delivery**

The effects of calcium acetate on labor and delivery are unknown.

### **8.3 Nursing Mothers**

Calcium acetate tablets contain calcium acetate and is excreted in human milk. Human milk feeding by a mother receiving calcium acetate is not expected to harm an infant, provided maternal serum calcium levels are appropriately monitored.

### **8.4 Pediatric Use**

Safety and effectiveness in pediatric patients have not been established.

### **8.5 Geriatric Use**


Clinical studies of calcium acetate did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other clinical experience has not identified differences in responses between elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

## 10 OVERDOSAGE

Administration of calcium acetate in excess of the appropriate daily dosage may result in hypercalcemia [see *Warnings and Precautions* (5.1)].

## 11 DESCRIPTION

Calcium acetate tablets act as a phosphate binder. Its chemical name is calcium acetate. Its molecular formula is  $C_4H_6CaO_4$ , and its molecular weight is 158.17. Its structural formula is:



Each tablet contains 667 mg calcium acetate, USP (anhydrous;  $Ca(CH_3COO)_2$ ; MW=158.17 grams) equal to 169 mg (8.45 mEq) calcium. Inactive Ingredients: polyethylene glycol 8000, sodium lauryl sulfate, crospovidone and sodium stearyl fumarate.

Calcium acetate tablets are administered orally for the control of hyperphosphatemia in end stage renal failure.

## 12 CLINICAL PHARMACOLOGY

Patients with ESRD retain phosphorus and can develop hyperphosphatemia. High serum phosphorus can precipitate serum calcium resulting in ectopic calcification. Hyperphosphatemia also plays a role in the development of secondary hyperparathyroidism in patients with ESRD.

### 12.1 Mechanism of Action

Calcium acetate, when taken with meals, combines with dietary phosphate to form an insoluble calcium phosphate complex, which is excreted in the feces, resulting in decreased serum phosphorus concentration.

### 12.2 Pharmacodynamics

Orally administered calcium acetate from pharmaceutical dosage forms is systemically absorbed up to approximately 40% under fasting conditions and up to approximately 30% under nonfasting conditions. This range represents data from both healthy subjects and renal dialysis patients under various conditions.

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment and Fertility

No carcinogenicity, mutagenicity, or fertility studies have been conducted with calcium

acetate.

## 14 CLINICAL STUDIES

Effectiveness of calcium acetate in decreasing serum phosphorus has been demonstrated in two studies of the calcium acetate solid oral dosage form.

Ninety-one patients with end-stage renal disease who were undergoing hemodialysis and were hyperphosphatemic (serum phosphorus >5.5 mg/dL) following a 1-week phosphate binder washout period contributed efficacy data to an open-label, non-randomized study.

The patients received calcium acetate 667 mg tablets at each meal for a period of 12 weeks. The initial starting dose was 2 tablets per meal for 3 meals a day, and the dose was adjusted as necessary to control serum phosphorus levels. The average final dose after 12 weeks of treatment was 3.4 tablets per meal. Although there was a decrease in serum phosphorus, in the absence of a control group the true magnitude of effect is uncertain.

The data presented in Table 2 demonstrate the efficacy of calcium acetate in the treatment of hyperphosphatemia in end-stage renal disease patients. The effects on serum calcium levels are also presented.

**Table 2: Average Serum Phosphorous and Calcium Levels at Pre-Study, Interim, and Study Completion Time points**

Parameter	Pre-Study	Week 4 *	Week 8	Week 12	p-value †
Phosphorus (mg/dL) ‡	7.4 ± 0.17	5.9 ± 0.16	5.6 ± 0.17	5.2 ± 0.17	≤0.01
Calcium (mg/dL) ‡	8.9 ± 0.09	9.5 ± 0.10	9.7 ± 0.10	9.7 ± 0.10	≤0.01

\* Ninety-one patients completed at least 6 weeks of the study.

† ANOVA of difference in values at pre-study and study completion.

‡ Values expressed as mean ± SE.

There was a 30% decrease in serum phosphorus levels during the 12 week study period ( $p < 0.01$ ).

Two-thirds of the decline occurred in the first month of the study. Serum calcium increased 9% during the study mostly in the first month of the study.

Treatment with the phosphate binder was discontinued for patients from the open-label study, and those patients whose serum phosphorus exceeded 5.5 mg/dL were eligible for entry into a double-blind, placebo-controlled, cross-over study. Patients were randomized to receive calcium acetate or placebo, and each continued to receive the same number of tablets as had been individually established during the previous study. Following 2 weeks of treatment, patients switched to the alternative therapy for an additional 2 weeks.

The phosphate binding effect of calcium acetate is shown in the Table 3.

**Table 3: Serum Phosphorous and Calcium Levels at Study Initiation and After Completion of Each Treatment Arm**

Parameter	Pre-Study	Post-Treatment		p-value *
		Calcium Acetate	Placebo	
Phosphorus (mg/dL) †	7.3 ± 0.18	5.9 ± 0.24	7.8 ± 0.22	<0.01
Calcium (mg/dL) †	8.9 ± 0.11	9.5 ± 0.13	8.8 ± 0.12	<0.01

\* ANOVA of calcium acetate vs. placebo after 2 weeks of treatment.

† Values expressed as mean ± SEM

Overall, 2 weeks of treatment with calcium acetate statistically significantly ( $p < 0.01$ ) decreased serum phosphorus by a mean of 19% and increased serum calcium by a statistically significant ( $p < 0.01$ ) but clinically unimportant mean of 7%.

## 16 HOW SUPPLIED/STORAGE AND HANDLING

Each tablet for oral administration is debossed "CE" over "2" on one side and plain on the other side. Each white round tablet contains 667 mg calcium acetate (anhydrous;  $\text{Ca}(\text{CH}_3\text{COO})_2$ ; MW=158.17 grams) equal to 169 mg (8.45 mEq) calcium.

NDC 62135-192-  
22 Bottles of 200

**STORAGE:** Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

## 17 PATIENT COUNSELING INFORMATION

Inform patients to take calcium acetate with meals, adhere to their prescribed diets, and avoid the use of calcium supplements including nonprescription antacids. Inform the patients about the symptoms of hypercalcemia [see *Warnings and Precautions (5.1)* and *Adverse Reactions (6.1)*].

Advise patients who are taking an oral medication where reduction in the bioavailability of that medication would have clinically significant effect on its safety or efficacy to take the drug one hour before or three hours after calcium acetate.

Manufactured by: Chartwell Pharmaceuticals, LLC.  
Congers, NY 10920

Manufactured for: Chartwell RX, LLC.

Made in USA

L70338

Rev: 05/2019

## PRINCIPAL DISPLAY PANEL - 667 mg Tablet Bottle Label

NDC 62135-192-22



Calcium Acetate  
Tablets, USP

667 mg\*

Rx Only

200 Tablets

Chartwell Rx

NDC 62135-**192**-22

**Calcium Acetate**  
Tablets, USP

667 mg\*

Rx Only  
200 Tablets



Chartwell Rx

\*Each tablet contains 667 mg calcium acetate, USP equivalent to 169 mg calcium.

**STORAGE:**  
Store at 20° to 25°C (68° to 77°F)  
[see USP Controlled Room Temperature].

**KEEP THIS AND ALL MEDICATIONS  
OUT OF REACH OF CHILDREN**


**DIRECTIONS:**  
SWALLOW TABLETS. DO NOT CHEW.  
Take as directed by your physician.

Manufactured by: Chartwell Pharmaceuticals, LLC.  
Congers, NY 10920  
Manufactured for: Chartwell RX, LLC.

GTIN 00362135192228

L70339 REV. 01 02/19

  
N 3 62135 19222 8

 Made in U.S.A.

UNVARNISHED  
AREA

## CALCIUM ACETATE

calcium acetate tablet

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:62135-192
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CALCIUM ACETATE (UNII: Y882YXF34X) (CALCIUM CATION - UNII:2M83C4R6ZB)	CALCIUM ACETATE	667 mg

### Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	CE2
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62135-192-22	200 in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2019	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202420	07/01/2019	

**Labeler** - Chartwell RX LLC (079394054)

## Establishment

Name	Address	ID/FEI	Business Operations
Chartwell Pharmaceuticals Congers, LLC.		118673447	analysis(62135-192) , label(62135-192) , manufacture(62135-192) , pack(62135-192)

Revised: 12/2022

Chartwell RX LLC