

SODIUM CHLORIDE- sodium chloride injection, solution
Baxter Healthcare Corporation

Sodium Chloride Injection, USP
in VIAFLO Plastic Container

DESCRIPTION

Sodium Chloride Injection, USP is a sterile, nonpyrogenic solution for fluid and electrolyte replenishment in single dose containers for intravenous administration. It contains no antimicrobial agents. The nominal pH is 5.0 (4.5 to 7.0). Composition, osmolarity, and ionic concentration are shown below:

0.9% Sodium Chloride Injection, USP contains 9 g/L Sodium Chloride, USP (NaCl) with an osmolarity of 308 mOsmol/L (calc). It contains 154 mEq/L sodium and 154 mEq/L chloride.

VIAFLO is a flexible plastic container fabricated from a multilayer sheeting (PL-2442) composed of Polypropylene (PP), Polyamide (PA) and Polyethylene (PE). The amount of water that can permeate from inside the container into the overwrap is insufficient to affect the solution significantly. Two different administration connectors are available with VIAFLO containers. The VIAFLO driplless access container (DAC) will not drip once the spike is removed. The non-DAC VIAFLO will drip once the spike is removed from the administration port.

VIAFLO is not made with natural rubber latex, DEHP, or PVC.

CLINICAL PHARMACOLOGY

Sodium Chloride Injection, USP has value as a source of water and electrolytes. It is capable of inducing diuresis depending on the clinical condition of the patient.

INDICATIONS AND USAGE

Sodium Chloride Injection, USP is indicated as a source of water and electrolytes.

0.9% Sodium Chloride Injection, USP is also indicated for use as a priming solution in hemodialysis procedures.

CONTRAINDICATIONS

None known.

WARNINGS

Hypersensitivity

Hypersensitivity and infusion reactions, including hypotension, pyrexia, tremor, chills,

urticaria, rash, and pruritus have been reported with 0.9% Sodium Chloride Injection, USP.

Stop the infusion immediately if signs or symptoms of a hypersensitivity reaction develop, such as tachycardia, chest pain, dyspnea and flushing. Appropriate therapeutic countermeasures must be instituted as clinically indicated.

Electrolyte Imbalances

Fluid Overload

Depending on the volume and rate of infusion, and the patient's underlying clinical condition, the intravenous administration of Sodium Chloride Injection, USP can cause fluid disturbances such as overhydration/hypervolemia and congested states, including pulmonary congestion and edema.

Avoid 0.9% Sodium Chloride Injection, USP in patients with or at risk for fluid and/or solute overloading. If use cannot be avoided, monitor fluid balance, electrolyte concentrations, and acid base balance, as needed and especially during prolonged use.

Hyponatremia

Sodium Chloride Injection, USP may cause hyponatremia. Hyponatremia can lead to acute hyponatremic encephalopathy characterized by headache, nausea, seizures, lethargy, and vomiting. Patients with brain edema are at particular risk of severe, irreversible and life-threatening brain injury.

The risk of hospital-acquired hyponatremia is increased in patients with cardiac or pulmonary failure, and in patients with non-osmotic vasopressin release (including SIADH) treated with high volume of Sodium Chloride Injection, USP.

The risk for hyponatremia is increased in pediatric patients, elderly patients, postoperative patients, those with psychogenic polydipsia, and in patients treated with medications that increase the risk of hyponatremia (such as diuretics, certain antiepileptic and psychotropic medications). See **DRUG INTERACTIONS**.

Patients at increased risk for developing complications of hyponatremia such as hyponatremic encephalopathy, include pediatric patients, women (in particular premenopausal women), patients with hypoxemia, and patients with underlying central nervous system disease. Avoid Sodium Chloride Injection, USP in patients with or at risk for hyponatremia. If use cannot be avoided, monitor serum sodium concentrations.

Rapid correction of hyponatremia is potentially dangerous with risk of serious neurologic complications. Brain adaptations reducing risk of cerebral edema make the brain vulnerable to injury when chronic hyponatremia is too rapidly corrected, which is known as osmotic demyelination syndrome (ODS). To avoid complications, monitor serum sodium and chloride concentrations, fluid status, acid-base balance, and signs of neurologic complications.

Hypernatremia

Hypernatremia may occur with Sodium Chloride Injection, USP. Conditions that may increase the risk of hypernatremia, fluid overload and edema (central and peripheral), include patients with: primary hyperaldosteronism; secondary hyperaldosteronism associated with, for example, hypertension, congestive heart failure, liver disease (including cirrhosis), renal disease (including renal artery stenosis, nephrosclerosis); and

pre-eclampsia.

Certain medications, such as corticosteroids or corticotropin, may also increase risk of sodium and fluid retention, see **DRUG INTERACTIONS**.

Avoid Sodium Chloride Injection, USP in patients with, or at risk for, hypernatremia. If use cannot be avoided, monitor serum sodium concentrations.

Rapid correction of hypernatremia is potentially dangerous with risk of serious neurologic complications. Excessively rapid correction of hypernatremia is also associated with a risk for serious neurologic complications such as osmotic demyelination syndrome (ODS) with risk of seizures and cerebral edema.

PRECAUTIONS

Patients with Severe Renal Impairment

Administration of Sodium Chloride Injection, USP in patients with or at risk of severe renal impairment, may result in hypernatremia and/or fluid overload (see **WARNINGS**). Avoid Sodium Chloride Injection, USP in patients with severe renal impairment or conditions that may cause sodium and/or potassium retention, fluid overload, or edema. If use cannot be avoided, monitor patients with severe renal impairment for development of these adverse reactions.

Drug Interactions

Other Products that Affect Fluid and/or Electrolyte Balance

Administration of Sodium Chloride Injection, USP to patients treated concomitantly with drugs associated with sodium and fluid retention may increase the risk of hypernatremia and volume overload. Avoid use of Sodium Chloride Injection, USP in patients receiving such products, such as corticosteroids or corticotropin. If use cannot be avoided, monitor serum electrolytes, fluid balance and acid base balance.

Lithium

Renal sodium and lithium clearance may be decreased during administration of Sodium Chloride Injection, USP. Monitor serum lithium concentration during concomitant use.

Renal sodium and lithium clearance may be increased during administration of 0.9% Sodium Chloride Injection, USP. Monitor serum lithium concentrations during concomitant use.

Other Drugs that Increase the Risk of Hyponatremia

Administration of Sodium Chloride Injection, USP in patients treated concomitantly with medications associated with hyponatremia may increase the risk of developing hyponatremia.

Avoid use of Sodium Chloride Injection, USP in patients receiving products, such as diuretics, and certain antiepileptic and psychotropic medications. Drugs that increase the vasopressin effect reduce renal electrolyte free water excretion and may also increase the risk of hyponatremia following treatment with intravenous fluids. If use cannot be avoided, monitor serum sodium concentrations.

Pregnancy

There are no adequate and well controlled studies with Sodium Chloride Injection, USP in pregnant women and animal reproduction studies have not been conducted with this drug. Therefore, it is not known whether Sodium Chloride Injection, USP can cause fetal harm when administered to a pregnant woman. Sodium Chloride Injection, USP should be given during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nursing Mothers

It is not known whether this drug is present in human milk. Because many drugs are present in human milk, caution should be exercised when Sodium Chloride Injection, USP is administered to a nursing woman.

Pediatric Use

The use of Sodium Chloride Injection, USP in pediatric patients is based on clinical practice. (See **DOSE AND ADMINISTRATION**).

Closely monitor plasma electrolyte concentrations in pediatric patients who may have impaired ability to regulate fluids and electrolytes. In very low birth weight infants, excessive or rapid administration of Sodium Chloride Injection, USP may result in increased serum osmolality and risk of intracerebral hemorrhage.

Children (including neonates and older children) are at increased risk of developing hyponatremia as well as for developing hyponatremic encephalopathy.

Geriatric Use

Geriatric patients are at increased risk of developing electrolyte imbalances. Sodium Chloride Injection, USP is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Therefore, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy. Consider monitoring renal function in elderly patients.

ADVERSE REACTIONS

Post-Marketing Adverse Reactions

The following adverse reactions have been identified during post approval use of Sodium Chloride Injection, USP. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

The following adverse reactions have been reported in the post-marketing experience during use of 0.9% Sodium Chloride Injection, USP and include the following:

General disorders and administration site conditions: Infusion site erythema, injection site streaking, burning sensation, and infusion site urticaria

Hypersensitivity reactions: Hypotension, pyrexia, tremor, chills, urticaria, rash, and

pruritus.

Metabolism and nutrition disorders: Hyponatremia, hyponatremia, hyperchloremic metabolic acidosis.

Nervous System Disorders: Hyponatremic encephalopathy

If an adverse reaction does occur, discontinue the infusion, evaluate the patient, institute appropriate therapeutic countermeasures and save the remainder of the fluid for examination if deemed necessary.

OVERDOSAGE

Excessive administration of:

- 0.9% Sodium Chloride Injection, USP may lead to hypernatremia. Hypernatremia can lead to CNS manifestations, including seizures, coma, cerebral edema and death.
- Sodium Chloride Injection, USP can cause fluid overload (which can lead to pulmonary and/or peripheral edema). See **WARNINGS and ADVERSE REACTIONS**.

When assessing an overdose, any additives in the solution must also be considered. The effects of an overdose may require immediate medical attention and treatment.

Interventions include discontinuation of Sodium Chloride Injection, USP administration, dose reduction, and other measures as indicated for the specific clinical constellation (e.g., monitoring of fluid balance, electrolyte concentrations and acid base balance).

DOSAGE AND ADMINISTRATION

Important Administration Instructions

- Sodium Chloride Injection, USP is intended for intravenous administration using sterile equipment.
- Do not connect flexible plastic containers in series in order to avoid air embolism due to possible residual air contained in the primary container.
- Set the vent to the closed position on a vented intravenous administration set to prevent air embolism.
- Use a dedicated line without any connections to avoid air embolism.
- Do not pressurize intravenous solutions contained in flexible plastic containers to increase flow rates in order to avoid air embolism due to incomplete evacuation of residual air in the container.
- Prior to infusion, visually inspect the solution for particulate matter and discoloration. The solution should be clear and there should be no precipitates. Do not administer unless solution is clear, and container is undamaged.

Dosing Information

The choice of product, dosage, volume, rate, and duration of administration is dependent upon the age, weight and clinical condition of the patient and concomitant therapy, and administration should be determined by a physician experienced in

intravenous fluid therapy.

Introduction of Additives

Additives may be incompatible.

Evaluate all additions to the plastic container for compatibility and stability of the resulting preparation. Consult with a pharmacist, if available.

If, in the informed judgment of the physician, it is deemed advisable to introduce additives, use aseptic technique. Mix thoroughly when additives have been introduced. After addition, if there is discoloration and/or the appearance of precipitates, insoluble complexes or crystals, do not use. Do not store solutions containing additives. Discard any unused portion.

HOW SUPPLIED

The available sizes of each 0.9% Sodium Chloride Injection, USP in VIAFLO dripless access containers (DAC) and non-DAC plastic containers are shown below. VIAFLO DAC will not drip once the spike is removed. VIAFLO non-DAC will drip once the spike is removed from the administration port:

Code	Size (mL)	Product	NDC
UE1306G	50	Non-DAC	0338-9657-75
UE1306DG	50	DAC	0338-9659-75
UE1307G	100	Non-DAC	0338-9661-60
UE1307DG	100	DAC	0338-9663-60
UE1322	250	Non-DAC	0338-9543-01
UE1322D	250	DAC	0338-9543-02
UE1323	500	Non-DAC	0338-9543-03
UE1323D	500	DAC	0338-9543-04
UE1324	1000	Non-DAC	0338-9543-05
UE1324D	1000	DAC	0338-9543-06

Exposure of pharmaceutical products to heat should be minimized. Avoid excessive heat. It is recommended the product be stored at room temperature (25°C/77°F); brief exposure up to 40°C/104°F does not adversely affect the product.

DIRECTIONS FOR USE OF VIAFLO PLASTIC CONTAINER

For Information on Risk of Air Embolism – see **DOSAGE AND ADMINISTRATION**.

To Open

Tear overwrap down side at slit and remove solution container. Visually inspect the container. If the outlet port protector is damaged, detached, or not present, discard container as solution path sterility may be impaired. Some opacity of the plastic due to moisture absorption during the sterilization process may be observed. This is normal and does not affect the solution quality or safety. The opacity will diminish gradually. Check for minute leaks by squeezing inner bag firmly. If leaks are found, discard solution as sterility may be impaired. If supplemental medication is desired, follow directions below.

1. Remove the container from the over wrap just before use.
2. Check for minute leaks by squeezing inner bag firmly. If leaks are found, discard solution, as sterility may be impaired.
3. Check solution for clarity and absence of foreign matter. If solution is not clear or contains foreign matter, discard the solution.

Preparation for Administration

1. Suspend container from eyelet support.
2. Remove plastic protector from outlet port at bottom of container.
 - Grip the small wing on the neck of the port with one hand,
 - grip the large wing on the cap with the other hand and twist,
 - the cap will pop off.
3. Use an aseptic method to set up the infusion.
4. Attach administration set. Refer to directions accompanying set for connection, priming of the set and administration of the solution.

To Add Medication

Additives may be incompatible.

To add medication before solution administration

1. Prepare medication site.
2. Using syringe with 19 to 22 gauge needle, puncture resealable medication port and

inject.

3. Mix solution and medication thoroughly. For high density medication such as potassium chloride, tap ports gently while ports are upright and mix thoroughly.

To add medication during solution administration

1. Close clamp on the set.
2. Prepare medication site.
3. Using syringe with 19 to 22 gauge needle, puncture resealable medication port and inject.
4. Remove container from IV pole and/or turn to an upright position.
5. Evacuate both ports by tapping them gently while container is in the upright position.
6. Mix solution and medication thoroughly.
7. Return container to in-use position and continue administration.

Baxter Healthcare Corporation

Deerfield, IL 60015 USA

Made in Spain

SA-30-02-879

Rev. November 2023

Baxter and Viaflo are trademarks of Baxter International Inc.

Customers can contact Baxter Medical Information at 1-800-933-0303 for more information on the containers.

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL

Baxter

NDC 0338-9543-01

UE1322



250 mL

0.9% Sodium Chloride Injection USP

50

pH 5.0 (4.5 – 7.0) Sterile non pyrogenic
Osmolarity 308 mOsm/L (calc) Single dose container

50

Each 100 mL contains 900 mg Sodium Chloride USP mEq/L
Sodium 154 Chloride 154

100

Additives may be incompatible Consult with pharmacist if available
when introducing additives Use aseptic technique Mix thoroughly
Do not store Dosage intravenously as directed by a physician
See directions Cautions Squeeze and inspect inner bags which
maintains product sterility Discard if leaks are found Must not be
used in series connections Do not use unless solution is clear

100

150

Rx Only Store unit in moisture barrier overwrap at room temperature
(25°C/77°F) until ready to use Avoid excessive heat See insert
VIAFLO is not made with natural rubber latex, DEHP, or PVC
Baxter Viaflo and PL-2442 are trademarks of Baxter
International Inc For product information
1-800-933-0303 Baxter Healthcare Corporation
Deerfield IL 60015 USA Made in Spain

150



UN-35-03-047

200



(01)00303389543015

200



Lot

**Expiry
Container Label**

Baxter logo

NDC 0338-9543-01
UE1322

Viaflo logo

250 mL

0.9% Sodium Chloride Injection USP

pH 5.0 (4.5 – 7.0)
Osmolarity 308 mOsm/L (calc)

Sterile non pyrogenic
Single dose container

Each 100 mL contains 900 mg Sodium Chloride USP mEq/L
Sodium 154 Chloride 154

Additives may be incompatible Consult with pharmacist if available
when introducing additives Use aseptic technique Mix thoroughly

Do not store Dosage intravenously as directed by a physician
See directions Cautions Squeeze and inspect inner bags which
maintains product sterility Discard if leaks are found Must not be
used in series connections Do not use unless solution is clear

Rx Only Store unit in moisture barrier overwrap at room temperature
(25°C/77°F) until ready to use Avoid excessive heat See insert

VIAFLO is not made with natural rubber latex, DEHP, or PVC
Baxter Viaflo and PL-2442 are trademarks of Baxter
International Inc For product information
1-800-933-0303 **Baxter Healthcare Corporation**
Deerfield IL 60015 USA Made in Spain

BAR CODE
(01) 00303389543015

Lot
Expiry

UN-35-03-047

50
100
150
200

Baxter

NDC 0338-9657-75 **UE1306G**
50 mL

0.9% Sodium Chloride Injection USP

pH 5.0 (4.5 – 7.0) Sterile non pyrogenic
Osmolarity 308 mOsm/L (calc) Single dose container
Each 50 mL contains 450 mg Sodium Chloride USP mEq/50mL
Sodium 8 Chloride 8

Read Package Leaflet for full Information

Additives may be incompatible Dosage intravenously
as directed by a physician Cautions Must not be
used in series connections Do not use unless solution
is clear Rx Only Store unit in moisture barrier
overwrap at room temperature (25°C/77°F) until ready
to use Avoid excessive heat See insert
VIAFLO is not made with natural rubber latex, DEHP,
or PVC


Baxter Viaflo and PL-2442 are trademarks of Baxter
International Inc

For product information 1-800-933-0303

Baxter Healthcare Corporation
Deerfield IL 60015 USA Made in Spain

Lot

UN-35-04-346 ①
Expiry



Container Label

Baxter Logo

NDC 0338-9657-75

UE1306G

Viaflo Logo

50 mL

0.9% Sodium Chloride Injection USP

pH 5.0 (4.5 – 7.0)

Sterile non pyrogenic

Osmolarity 308 mOsm/L (calc)

Single dose container

Each 50 mL contains 450 mg Sodium Chloride USP mEq/50mL

Sodium 8 Chloride 8

Read Package Leaflet for full Information

Additives may be incompatible Dosage intravenously
as directed by a physician Cautions Must not be
used in series connections Do not use unless solution
is clear Rx Only Store unit in moisture barrier
overwrap at room temperature (25°C/77°F) until ready
to use Avoid excessive heat See insert
VIAFLO is not made with natural rubber latex, DEHP,
or PVC

Baxter Viaflo and PL-2442 are trademarks of Baxter
International Inc

For product information 1-800-933-0303

07 Symbol

Baxter Healthcare Corporation

Deerfield IL 60015 USA Made in Spain

UN-35-04-346 **1** Symbol

Lot Expiry

Barcode

(01)00303389657750

Baxter

NDC 0338-9659-75

UE1306DG



50 mL

0.9% Sodium Chloride Injection USP

pH 5.0 (4.5 – 7.0)

Osmolarity 308 mOsm/L (calc)

Sterile non pyrogenic

Single dose container

Each 50 mL contains 450 mg Sodium Chloride USP mEq/50mL

Sodium 8 Chloride 8

Read Package Leaflet for full Information

Additives may be incompatible Dosage intravenously as directed by a physician Cautions Must not be used in series connections Do not use unless solution is clear Rx Only Store unit in moisture barrier overwrap at room temperature (25°C/77°F) until ready to use Avoid excessive heat See insert

VIAFLO is not made with natural rubber latex, DEHP, or PVC

Baxter Viaflo and PL-2442 are trademarks of Baxter International Inc

For product information 1-800-933-0303

Baxter Healthcare Corporation

Deerfield IL 60015 USA Made in Spain

Lot



UN-35-04-147 ①
Expiry

Container Label

Baxter Logo

NDC 0338-9659-75

UE1306DG

Viaflo Logo

50 mL

0.9% Sodium Chloride Injection USP

pH 5.0 (4.5 – 7.0)

Sterile non pyrogenic

Osmolarity 308 mOsm/L (calc)

Single dose container

Each 50 mL contains 450 mg Sodium Chloride USP mEq/50mL

Sodium 8 Chloride 8

Read Package Leaflet for full Information

Additives may be incompatible Dosage intravenously as directed by a physician Cautions Must not be used in series connections Do not use unless solution is clear Rx Only Store unit in moisture barrier overwrap at room temperature (25°C/77°F) until ready to use Avoid excessive heat See insert

VIAFLO is not made with natural rubber latex, DEHP, or PVC

Baxter Viaflo and PL-2442 are trademarks of Baxter International Inc

For product information 1-800-933-0303

07 Symbol

Baxter Healthcare Corporation

Deerfield IL 60015 USA Made in Spain

UN-35-04-147 **1** Symbol


Lot Expiry

Barcode

(01)00303389659754

Baxter

NDC 0338-9661-60 UE1307G

 **100 mL**

0.9% Sodium Chloride Injection USP

pH 5.0 (4.5 – 7.0) Sterile non pyrogenic
Osmolarity 308 mOsm/L (calc) Single dose container

Each 100 mL contains 900 mg Sodium Chloride USP mEq/100mL
Sodium 15 Chloride 15

Read Package Leaflet for full Information

Additives may be incompatible Dosage intravenously as directed by a physician Cautions Must not be used in series connections Do not use unless solution is clear Rx Only Store unit in moisture barrier overwrap at room temperature (25°C/77°F) until ready to use Avoid excessive heat See insert


VIAFLO is not made with natural rubber latex, DEHP, or PVC

Baxter Viaflo and PL-2442 are trademarks of Baxter International Inc


For product information 1-800-933-0303

Baxter Healthcare Corporation
Deerfield IL 60015 USA Made in Spain

Lot

 UN-35-04-347 ①
Expiry

Container Label



Baxter Logo

NDC 0338-9661-60

UE1307G

Viaflo Logo
100 mL

**0.9% Sodium Chloride
Injection USP**

pH 5.0 (4.5 – 7.0)
Sterile non pyrogenic
Osmolarity 308 mOsm/L (calc)
Single dose container

Each 100 mL contains 900 mg Sodium Chloride USP mEq/100mL
Sodium 15 Chloride 15

Read Package Leaflet for full Information

Additives may be incompatible Dosage intravenously
as directed by a physician Cautions Must not be
used in series connections Do not use unless solution
is clear Rx Only Store unit in moisture barrier
overwrap at room temperature (25°C/77°F) until ready
to use Avoid excessive heat See insert
VIAFLO is not made with natural rubber latex, DEHP,
or PVC

Baxter Viaflo and PL-2442 are trademarks of Baxter
International Inc

For product information 1-800-933-0303

07 Symbol

Baxter Healthcare Corporation
Deerfield IL 60015 USA Made in Spain

UN-35-04-347 **1** Symbol

Lot Expiry

Barcode
(01)00303389661603

Baxter

NDC 0338-9663-60

UE1307DG



100 mL

**0.9% Sodium Chloride
Injection USP**

pH 5.0 (4.5 – 7.0)

Osmolarity 308 mOsm/L (calc)

Sterile non pyrogenic

Single dose container

Each 100 mL contains 900 mg Sodium Chloride USP mEq/100mL
Sodium 15 Chloride 15

Read Package Leaflet for full Information

Additives may be incompatible Dosage intravenously
as directed by a physician Cautions Must not be
used in series connections Do not use unless solution
is clear Rx Only Store unit in moisture barrier
overwrap at room temperature (25°C/77°F) until ready
to use Avoid excessive heat See insert

VIAFLO is not made with natural rubber latex, DEHP,
or PVC

Baxter Viaflo and PL-2442 are trademarks of Baxter
International Inc

For product information 1-800-933-0303

Baxter Healthcare Corporation

Deerfield IL 60015 USA Made in Spain

Lot



UN-35-04-148 ①

Expiry

Container Label

Baxter Logo

NDC 0338-9663-60

UE1307DG

Viaflo Logo

100 mL

**0.9% Sodium Chloride
Injection USP**

pH 5.0 (4.5 – 7.0)

Sterile non pyrogenic

Osmolarity 308 mOsm/L (calc)

Single dose container

Each 100 mL contains 900 mg Sodium Chloride USP mEq/100mL
Sodium 15 Chloride 15

Read Package Leaflet for full Information

Additives may be incompatible Dosage intravenously
as directed by a physician Cautions Must not be
used in series connections Do not use unless solution

is clear Rx Only Store unit in moisture barrier overwrap at room temperature (25°C/77°F) until ready to use Avoid excessive heat See insert VIAFLO is not made with natural rubber latex, DEHP, or PVC

Baxter Viaflo and PL-2442 are trademarks of Baxter International Inc

For product information 1-800-933-0303

07 Symbol

Baxter Healthcare Corporation
Deerfield IL 60015 USA Made in Spain

UN-35-04-148 **1** Symbol

Lot Expiry

Barcode
(01)00303389663607

UE1306G 75 x 50 mL (≈ 6 kg)

0.9% Sodium Chloride Injection, USP

Formula per 50 mL
Sodium Chloride 450 mg
Water for Injection
Osmolarity 308 mOsm/L (calc)
mmol per 50 mL (approx.)
Sodium 8
Chloride 8
pH 5.0 (4.5 – 7.0)

IV administration.
Read package insert before use.
Keep out of the reach and sight of children.
Do not connect partially used drugs.

Baxter Healthcare Corporation
Deerfield IL 60015 USA
Made in Spain



P333E744
??-0001

Lot: ???????
Exp: YYYY/MM



**UE1306G
75 x 50 mL
(≈ 6 kg)**

0.9% Sodium Chloride Injection, USP



Lot: ???????
Exp: YYYY/MM

(01)50303389663607(10)???????

Carton Label

UE1306G 75 x 50 mL (≈ 6 kg)

0.9% Sodium Chloride Injection, USP

Formula per 50 mL
Sodium Chloride 450 mg
Water for Injection

Osmolarity 308 mOsm/L (calc)

mmol per 50 mL (approx..)

Sodium 8

Chloride 8

pH 5.0 (4.5 – 7.0)

IV administration.

Read package insert before use.

Keep out of the reach and sight of children.

Do not connect partially used drugs.

Baxter Healthcare Corporation

Deerfield IL 60015 USA

Made in Spain

Viaflo Logo

Lot: ????????

Exp: YYYY/MM

P333E744

?? - 00001

Barcode

(01)50303389657755(17)YYMMOO(10)?????????

UE1306G 75 x 50 mL (≈ 6 kg)

0.9% Sodium

Chloride

Injection, USP

Viaflo Logo

2D barcode

Lot:

?????????

Exp:

YYYY/MM

(01)50303389657755(17)YYMMOO(10)?????????

UE1306DG 75 x 50 mL (≈ 6 kg)

0.9% Sodium Chloride Injection, USP

Formula per 50 mL
Sodium Chloride 450 mg
Water for Injection
Osmolarity 308 mOsm/L (calc)
mmol per 50 mL (approx.)
Sodium 8
Chloride 8
pH 5.0 (4.5 – 7.0)
IV administration.
Read package insert before use.
Keep out of the reach and sight of children.
Do not connect partially used drugs.

Baxter Healthcare Corporation
Deerfield IL 60015 USA
Made in Spain



P333E748
?? - 0001

Lot: ????????
Exp: YYYY/MM



(01)5030389659759(17)YYMMDD(10)???????

UE1306DG
75 x 50 mL
(≈ 6 kg)

0.9% Sodium Chloride Injection, USP



Lot:
????????
Exp:
YYYY/MM

(01)5030389659759(17)YYMMDD(10)???????

Carton Label

UE1306DG 75 x 50 mL (≈ 6 kg)

0.9% Sodium Chloride Injection, USP

Formula per 50 mL
Sodium Chloride 450 mg
Water for Injection
Osmolarity 308 mOsm/L (calc)
mmol per 50 mL (approx..)
Sodium 8
Chloride 8
pH 5.0 (4.5 – 7.0)

IV administration.
Read package insert before use.
Keep out of the reach and sight of children.
Do not connect partially used drugs.

Baxter Healthcare Corporation
Deerfield IL 60015 USA

Made in Spain

Viaflo Logo

**Lot: ????????
Exp: YYYY/MM**

P333E748
?? - 0001

Barcode
(01)50303389659759(17)YYMMOO(10)????????

UE1306DG
75 x 50 mL (≈ 6 kg)

**0.9% Sodium
Chloride
Injection, USP**

Viaflo Logo

2D barcode

Lot:
????????

Exp:
YYYY/MM

(01)50303389659759(17)YYMMOO(10)????????

UE1307G 60 x 100 mL (≈ 8 kg)

**0.9% Sodium Chloride
Injection, USP**

Formula per 100 mL
Sodium Chloride 900 mg
Water for Injection
Osmolarity 308 mOsm/L (calc)
mmol per 100 mL (approx.)
Sodium 15
Chloride 15
pH 5.0 (4.5 – 7.0)
IV administration.

Read package insert before use.
Keep out of the reach and sight of children.
Do not connect partially used drugs.

Baxter Healthcare Corporation
Deerfield IL 60015 USA
Made in Spain



P338E745
?-0001

Lot: ????????

Exp: YYYY/MM



(01)50303389651608(17)YYMMOO(10)????????

UE1307G
60 x 100 mL
(≈ 8 kg)

**0.9% Sodium
Chloride
Injection, USP**



Lot:
????????

Exp:
YYYY/MM

(01)50303389651608(17)YYMMOO(10)????????

Carton Label

UE1307G 60 x 100 mL (≈ 8 kg)

**0.9% Sodium Chloride
Injection, USP**

Formula per 100 mL
Sodium Chloride 900 mg
Water for Injection
Osmolarity 308 mOsm/L (calc)

mmol per 100 mL (approx..)

Sodium 15

Chloride 15

pH 5.0 (4.5 – 7.0)

IV administration.

Read package insert before use.

Keep out of the reach and sight of children.

Do not connect partially used drugs.

Baxter Healthcare Corporation

Deerfield IL 60015 USA

Made in Spain

Viaflo Logo

Lot: ????????

Exp: YYYY/MM

P333E745

?? - 00001

Barcode

(01)503033896616608(17)YYMMOO(10)?????????

UE1307G

60 x 100 mL (≈ 8 kg)

0.9% Sodium

Chloride

Injection, USP

Viaflo Logo

2D barcode

Lot:

?????????

Exp:

YYYY/MM

(01)503033896616608(17)YYMMOO(10)?????????

UE1307DG 60 x 100 mL (≈ 8 kg)

0.9% Sodium Chloride Injection, USP

Formula per 100 mL
Sodium Chloride 900 mg
Water for Injection
Osmolarity 308 mOsm/L (calc)
mmol per 100 mL (approx.)
Sodium 15
Chloride 15
pH 5.0 (4.5 – 7.0)

IV administration.
Read package insert before use.
Keep out of the reach and sight of children.
Do not connect partially used drugs.

Baxter Healthcare Corporation
Deerfield IL 60015 USA
Made in Spain



P333E749
?? - 0001

Lot: ????????
Exp: YYYY/MM



(01)50303389663602(17)YYMM00(10)???????

UE1307DG
60 x 100 mL
(≈ 8 kg)

0.9% Sodium Chloride Injection, USP



Lot:
????????
Exp:
YYYY/MM

(01)50303389663602(17)YYMM00(10)???????

Carton Label

UE1307DG 60 x 100 mL (≈ 8 kg)

0.9% Sodium Chloride Injection, USP

Formula per 100 mL
Sodium Chloride 900 mg
Water for Injection
Osmolarity 308 mOsm/L (calc)
mmol per 100 mL (approx..)
Sodium 15
Chloride 15
pH 5.0 (4.5 – 7.0)

IV administration.
Read package insert before use.
Keep out of the reach and sight of children.
Do not connect partially used drugs.

Baxter Healthcare Corporation
Deerfield IL 60015 USA

Made in Spain

Viaflo Logo

Lot: ????????
Exp: YYYY/MM

P333E749
?? - 0001

Barcode

(01)50303389663602(17)YYMMOO(10)?????????

UE1307G

60 x 100 mL (≈ 8 kg)

**0.9% Sodium
Chloride
Injection, USP**

Viaflo Logo

2D barcode

Lot:

?????????

Exp:

YYYY/MM

(01)50303389663602(17)YYMMOO(10)?????????

SODIUM CHLORIDE				
sodium chloride injection, solution				
Product Information				
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0338-9543	
Route of Administration	INTRAVENOUS			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)		SODIUM CHLORIDE	9 g in 100 mL	
Inactive Ingredients				
Ingredient Name		Strength		
WATER (UNII: 059QF0K00R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-9543-01	250 mL in 1 BAG; Type 0: Not a Combination Product	12/09/1970	
2	NDC:0338-9543-02	250 mL in 1 BAG; Type 0: Not a Combination Product	12/09/1970	
3	NDC:0338-9543-03	500 mL in 1 BAG; Type 0: Not a Combination Product	12/09/1970	
4	NDC:0338-9543-04	500 mL in 1 BAG; Type 0: Not a Combination Product	12/09/1970	
5	NDC:0338-9543-	1000 mL in 1 BAG; Type 0: Not a Combination	12/09/1970	

5	05	Product	12/09/1970	
6	NDC:0338-9543-06	1000 mL in 1 BAG; Type 0: Not a Combination Product	12/09/1970	
7	NDC:0338-9543-12	1000 mL in 1 BAG; Type 0: Not a Combination Product	12/09/1970	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA016677	12/09/1970	

SODIUM CHLORIDE

sodium chloride injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0338-9657
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	9 g in 1000 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-9657-75	1000 mL in 1 BAG; Type 0: Not a Combination Product	08/09/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA016677	12/09/1970	

SODIUM CHLORIDE

sodium chloride injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0338-9659
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	9 g in 1000 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-9659-75	1000 mL in 1 BAG; Type 0: Not a Combination Product	08/09/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA016677	12/09/1970	

SODIUM CHLORIDE

sodium chloride injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0338-9661
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	9 g in 1000 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-9661-60	1000 mL in 1 BAG; Type 0: Not a Combination Product	08/09/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA016677	12/09/1970	

SODIUM CHLORIDE

sodium chloride injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0338-9663
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	9 g in 1000 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-9663-60	1000 mL in 1 BAG; Type 0: Not a Combination Product	08/09/2024	

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
-----------	---------------------------------	-----------------	---------------

Category	Citation	Date	Date
NDA	NDA016677	12/09/1970	

Labeler - Baxter Healthcare Corporation (005083209)

Establishment

Name	Address	ID/FEI	Business Operations
Bieffe Medital SA		464755693	ANALYSIS(0338-9543, 0338-9657, 0338-9659, 0338-9661, 0338-9663) , MANUFACTURE(0338-9543, 0338-9657, 0338-9659, 0338-9661, 0338-9663) , LABEL(0338-9543, 0338-9657, 0338-9659, 0338-9661, 0338-9663) , PACK(0338-9543, 0338-9657, 0338-9659, 0338-9661, 0338-9663) , STERILIZE(0338-9543, 0338-9657, 0338-9659, 0338-9661, 0338-9663)

Establishment

Name	Address	ID/FEI	Business Operations
Baxter Healthcare S.A.		988899845	ANALYSIS(0338-9543, 0338-9657, 0338-9659, 0338-9661, 0338-9663)

Revised: 8/2024

Baxter Healthcare Corporation