

**MOTION SICKNESS RELIEF- dimenhydrinate tablet**  
**L.N.K. International, Inc.**

-----  
**Quality Plus 44-198**

***Active ingredient (in each tablet)***

Dimenhydrinate 50 mg

***Purpose***

Antiemetic

***Uses***

for prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

***Warnings***

**Do not use**

for children under 2 years of age unless directed by a doctor.

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

**Ask a doctor or pharmacist before use if you are**

taking sedatives or tranquilizers.

**When using this product**

- marked drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- to prevent motion sickness, the first dose should be taken one-half to one hour before starting activity

adults and children 12 years and over	1 to 2 tablets every 4-6 hours; do not exceed 8 tablets in 24 hours, or as directed by a doctor
children 6 to under 12 years	½ to 1 tablet every 6-8 hours; do not exceed 3 tablets in 24 hours, or as directed by a doctor
children 2 to under 6 years	½ tablet every 6-8 hours; do not exceed 1½ tablets in 24 hours, or as directed by a doctor

**Other information**

- **each tablet contains:** calcium 35 mg
- see end flap for expiration date and lot number
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)
- protect from moisture

**Inactive ingredients**

croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid

**Questions or comments?**

**1-800-426-9391**

**Principal display panel**

**QUALITY  
+PLUS**

NDC 50844-199-02

\*Compare to active ingredient in  
Dramamine® Original Formula

**ORIGINAL FORMULA**

**Motion Sickness Relief**

Dimenhydrinate 50 mg • Antiemetic

PREVENTS NAUSEA,  
VOMITING & DIZZINESS  
FOR CHILDREN & ADULTS

**12  
Tablets**

ACTUAL  
SIZE

\*This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Dramamine® Original Formula.

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

50844 REV0518B19802

Distributed by **LNK INTERNATIONAL, INC.**  
60 Arkay Drive  
Hauppauge, NY 11788  
USA

No Print/No Varnish  
Lot & Exp Date

3 50844 19902 7

P-1603-198-02-R  
REV0518B19802

**QUALITY PLUS**

**ORIGINAL FORMULA**

# Motion Sickness Relief

**Dimenhydrinate 50 mg • Antiemetic**

**12 Tablets**



**PREVENTS NAUSEA,  
VOMITING & DIZZINESS  
FOR CHILDREN & ADULTS**

**ACTUAL SIZE**



NDC 50844-199-02

\*Compare to active ingredient in Dramamine® Original Formula

**Drug Facts** (continued)

**Warnings**

Do not use for children under 2 years of age unless directed by a doctor.

nausea ■ vomiting ■ dizziness

**Drug Facts** (continued)

**Drug Facts**

**KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION**

**Active ingredient (in each tablet)**  
Dimenhydrinate 50 mg - Antiemetic

**Purpose**  
Antiemetic

**Uses**  
for prevention and treatment of these symptoms associated with motion sickness:

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

50844 REV0518B19802  
Distributed by LNK INTERNATIONAL, INC.  
60 Arkay Drive  
Hauppauge, NY 11788  
USA

**Drug Facts** (continued)

**Directions**

to prevent motion sickness, the first dose should be taken one-half to one hour before starting activity.

adults and children 12 years and over: 1 to 2 tablets every 4-6 hours, do not exceed 8 tablets in 24 hours, or as directed by a doctor.

children 6 to under 12 years: ½ to 1 tablet every 6-8 hours, do not exceed 3 tablets in 24 hours, or as directed by a doctor.

children 2 to under 6 years: ½ tablet every 6-8 hours, do not exceed 1½ tablets in 24 hours, or as directed by a doctor.

**Other information**

each tablet contains: calcium 35 mg

see end flap for expiration date and lot number

**Drug Facts** (continued)

**Warnings**

Ask a doctor before use if you have a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland or tranquilizers.

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

**When using this product**

marked drowsiness may occur ■ avoid alcoholic beverages ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Drug Facts** (continued)

**Warnings**

TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ protect from moisture

**Questions or comments?** 1-800-426-9391

**Drug Facts** (continued)

**Warnings**

This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Dramamine® Original Formula.

**Drug Facts** (continued)

**Warnings**

Do not use for children under 2 years of age unless directed by a doctor.

nausea ■ vomiting ■ dizziness

**Drug Facts** (continued)

**Quality Plus 44-198**

**MOTION SICKNESS RELIEF**

dimenhydrinate tablet

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50844-199
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

<b>DIMENHYDRINATE</b> (UNII: JB937PER5C) (DIPHENHYDRAMINE - UNII:8GTS82S83M, 8-CHLOROTHEOPHYLLINE - UNII:GE2UA340FM)	<b>DIMENHYDRINATE</b>	50 mg
--	-----------------------	-------

### Inactive Ingredients

Ingredient Name	Strength
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>DIBASIC CALCIUM PHOSPHATE DIHYDRATE</b> (UNII: O7TSZ97GEP)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	44;198
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-199-02	2 in 1 CARTON	12/01/1992	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M009	12/01/1992	

**Labeler** - L.N.K. International, Inc. (038154464)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(50844-199)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(50844-199) , pack(50844-199)

### Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

Revised: 6/2024

L.N.K. International, Inc.