

**4386 FIRST AID KIT- 4386 first aid**  
**Honeywell Safety Products USA, Inc.**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click [here](#).*

-----  
**0498-4386: First Aid Kit (PVP wipes, NaCl irr, EW, BZK wipe, alcohol wipe, PAWS- Z346100)**

**Eyewash**  
***Active ingredient***

Sterile Water 99%

**Eyewash**  
***Purpose***

Eyewash

**Eyewash**  
***Uses***

- for flushing the eye to remove loose foreign material, air pollutants or chlorinated water

**Eyewash**  
***Warnings***

**For external use only**

- Obtain immediate medical treatment for all open wounds in or near eyes.
- To avoid contamination, do not touch tip of container to any surface.
- Do not reuse.
- Once opened, discard.

**Do not use**

- if solution changes color or becomes cloudy
- if you have open wounds in or near the eyes, get medical help right away.

**Stop use and ask a doctor if**

- you experience eye pain
- changes in vision
- continued redness or irritation of the eye
- condition worsens or persists

**Keep out of reach of children**

If swallowed, get medical help or contact a Poison Control Center right away.

## **Eyewash**

### ***Directions***

- remove contacts before using
- twist top to remove
- flush the affected area as needed
- control rate of flow by pressure on the bottle
- if necessary, continue flushing with emergency eyewash or shower

## **Eyewash**

### ***Inactive ingredients***

sodium chloride, sodium phosphate dibasic, sodium phosphate monobasic

## **Eyewash**

### ***Questions***

Call 1-800-430-5490 Honeywell Safety Products USA, Inc. Smithfield, RI 02917

## **Isotonic Solution for Irrigation.**

For Irrigation Only.

Not for Injection.

## **Description**

### **NaCL Irrigation**

Each 100 mL contains:

Sodium Chloride USP 0.9 g; Water for Injection USP qs

pH adjusted with Hydrochloric Acid NF

pH: 5.0 (4.5–7.0) Calculated Osmolarity: 310 mOsmol/liter

Concentration of Electrolytes (mEq/liter): Sodium 154; Chloride 154

0.9% Sodium Chloride Irrigation USP is sterile, nonpyrogenic, isotonic and contains no bacteriostatic or antimicrobial agents.

The formula of the active ingredient is:

Ingredient Molecular Formula Molecular Weight

Sodium Chloride USP NaCl 58.44

The plastic container is a copolymer of ethylene and propylene formulated and developed for parenteral drugs. The copolymer contains no plasticizers and exhibits virtually no leachability. The plastic container is also virtually impermeable to vapor transmission and, therefore, requires no overwrap to maintain the proper drug

concentration. The safety of the plastic container has been confirmed by biological evaluation procedures. The material passes Class VI testing as specified in the U.S. Pharmacopeia for Biological Tests — Plastic Containers. These tests have shown that the container is nontoxic and biologically inert.

The PIC™ Container is PVC-free and DEHP-free.

Ingredient Molecular Formula Molecular Weight  
Sodium Chloride USP NaCl 58.44

The plastic container is a copolymer of ethylene and propylene formulated and developed for parenteral drugs. The copolymer contains no plasticizers and exhibits virtually no leachability. The plastic container is also virtually impermeable to vapor transmission and, therefore, requires no overwrap to maintain the proper drug concentration. The safety of the plastic container has been confirmed by biological evaluation procedures. The material passes Class VI testing as specified in the U.S. Pharmacopeia for Biological Tests — Plastic Containers. These tests have shown that the container is nontoxic and biologically inert.

The PIC™ Container is PVC-free and DEHP-free.

## **Clinical Pharmacology**

### **NaCL Irrigant**

0.9% Sodium Chloride Irrigation USP is utilized for a variety of clinical indications such as sterile irrigation of body cavities, tissues or wounds, indwelling urethral catheters, surgical drainage tubes, and for washing, rinsing or soaking surgical dressings, instruments and laboratory specimens. It also serves as a diluent or vehicle for drugs used for irrigation or other pharmaceutical preparations.

0.9% Sodium Chloride Irrigation USP provides an isotonic saline irrigation identical in composition with 0.9% Sodium Chloride Injection USP (normal saline).

Physiological irrigation solutions are considered generally compatible with living tissues and organs.

Sodium, the major cation of the extracellular fluid, functions primarily in the control of water distribution, fluid balance, and osmotic pressure of body fluids. Sodium is also associated with chloride and bicarbonate in the regulation of the acid-base equilibrium of body fluid.

Chloride, the major extracellular anion, closely follows the metabolism of sodium, and changes in the acid-base balance of the body are reflected by changes in the chloride concentration.

## **Indication and Usage**

### **NaCl Irrigant**

0.9% Sodium Chloride Irrigation USP is indicated for all general irrigation, washing, rinsing and dilution purposes which permit use of a sterile, nonpyrogenic electrolyte solution.

## **Contraindications**

### **NaCl Irrigant**

0.9% Sodium Chloride Irrigation USP is not for injection by usual parenteral routes.

An electrolyte solution should not be used for irrigation during electrosurgical procedures.

## **Warnings**

### **NaCl Irrigant**

FOR IRRIGATION ONLY. NOT FOR INJECTION.

Irrigating fluids have been demonstrated to enter the systemic circulation in relatively large volumes; thus, irrigation solutions must be regarded as systemic drugs. Absorption of large amounts can cause fluid and/or solute overload resulting in dilution of serum electrolyte concentrations, overhydration, congested states or pulmonary edema.

The risk of dilutional states is inversely proportional to the electrolyte concentrations of the administered parenteral solutions. The risk of solute overload causing congested states with peripheral and pulmonary edema is directly proportional to the electrolyte concentration.

Do not warm above 150°F (66°C).

After opening container, its contents should be used promptly to minimize the possibility of bacterial growth or pyrogen formation.

Discard unused portion of irrigating solution since it contains no preservatives.

## **Precautions**

### **NaCl Irrigant**

#### **General**

Use aseptic technique when preparing and administering sterile irrigation solutions.

Use only if solution is clear and container and seal are intact.

Do not use for irrigation that may result in absorption of large amounts of fluid into the blood.

Caution should be observed when the solution is used for continuous irrigation or allowed to "dwell" inside body cavities because of possible absorption into the blood stream and the production of circulatory overload.

When used for irrigation via appropriate irrigation equipment, the administration set should be attached promptly. Unused portions should be discarded and a fresh container of appropriate size used for the start up of each cycle or repeat procedure. For repeated irrigations of urethral catheters, a separate container should be used for each patient.

## **Laboratory Tests**

Clinical evaluation and periodic laboratory determinations are necessary to monitor changes in fluid balance, electrolyte concentrations, and acid-base balance after prolonged irrigation, when fluid absorption is suspected, or whenever the condition of the patient warrants such evaluation.

## **Drug Interactions**

Some additives may be incompatible. Consult with pharmacist. When introducing additives, use aseptic technique. Mix thoroughly.

Do not store.

## **Carcinogenesis, Mutagenesis, Impairment of Fertility**

Studies with 0.9% Sodium Chloride Irrigation USP have not been performed to evaluate carcinogenic potential, mutagenic potential, or effects on fertility.

## **Pregnancy**

### **Teratogenic Effects**

Animal reproduction studies have not been conducted with 0.9% Sodium Chloride Irrigation USP. It is also not known whether 0.9% Sodium Chloride Irrigation USP can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. 0.9% Sodium Chloride Irrigation USP should be given to a pregnant woman only if clearly needed.

## **Labor and Delivery**

Safety and effectiveness of 0.9% Sodium Chloride Irrigation USP during labor and delivery have not been established. Caution should be exercised, and the fluid balance, glucose and electrolyte concentrations, and acid-base balance, of both mother and fetus should be evaluated periodically or whenever warranted by the condition of the patient or fetus.

## **Nursing Mothers**

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when 0.9% Sodium Chloride Irrigation USP is administered to a nursing woman.

## **Pediatric Use**

The safety and effectiveness of 0.9% Sodium Chloride Irrigation USP in pediatric patients have not been established. Its limited use in pediatric patients has been inadequate to fully define proper dosage and limitations for use.

## **Geriatric Use**

Clinical studies of 0.9% Sodium Chloride Irrigation USP did not include a sufficient number of patients age 65 years and over to determine whether they respond differently from younger subjects. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater

frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function. Frequent laboratory determinations and clinical evaluations are recommended to monitor changes in blood glucose, electrolyte concentrations, and renal function.

## **Adverse Reactions**

Possible adverse effects arising from the irrigation of body cavities, tissues, or indwelling catheters and tubes can be minimized when proper procedures are followed. Displaced catheters or drainage tubes can lead to irrigation or infiltration of unintended structures or cavities. Excessive volume or pressure during irrigation of closed cavities may cause undue distension or disruption of tissues. Accidental contamination from careless technique may transmit infection.

If an adverse reaction does occur, discontinue administration of the irrigant, evaluate the patient, institute appropriate therapeutic countermeasures, and save the remainder of the fluid for examination if deemed necessary.

## **Overdosage**

In the event of overhydration or solute overload, reevaluate the patient's condition, and institute appropriate corrective treatment. Intravascular volume overload may respond to hemodialysis. See WARNINGS, PRECAUTIONS, and ADVERSE REACTIONS.

## **Dosage and Administration**

As required for irrigation.

When used as a diluent, or vehicle for other drugs, the drug manufacturer's recommendations should be followed.

Some additives may be incompatible. Consult with pharmacist. When introducing additives, use aseptic techniques. Mix thoroughly. Do not store.

Solutions should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permits.

## **How Supplied**

0.9% Sodium Chloride Irrigation USP is supplied sterile and nonpyrogenic in PIC™ (Plastic Irrigation Container). The 1000 mL and 500 mL containers are packaged 16 per case, the 2000 mL containers are packaged 8 per case, and the 4000 mL containers are packaged 4 per case.

0.9% Sodium Chloride Irrigation USP

NDC Cat. No. REF SIZE

0264-2201-00 R5200-01 1000 mL

0264-2201-10 R5201-01 500 mL

0264-2201-50 R5205-01 2000 mL

0264-2201-70 R5207 ,,,,4000 mL

Exposure of pharmaceutical products to heat should be minimized. Avoid excessive heat. Protect from freezing. It is recommended that the product be stored at room temperature (25°C); however, brief exposure up to 40°C does not adversely affect the product.

Do not warm above 150°F (66°C).

### **SPL Unclassified Section**

Rx only

Revised: March 2009

PIC is a trademark of B. Braun Medical Inc.

### **DIRECTIONS FOR USE OF PIC™ (PLASTIC IRRIGATION CONTAINER)**

Not for injection.

Aseptic technique is required.

Caution – Before use, perform the following checks:

(a) Read the label. Ensure solution is the one ordered and is within the expiration date.

(b) Invert container and inspect the solution in good light for cloudiness, haze, or particulate matter; check the container for leakage or damage. Any container which is suspect should not be used.

Use only if solution is clear and container and seal are intact

Single unit container. Discard unused portion.

Outer Closure Removal – Grasp the container with one hand and turn the breakaway ring counterclockwise with the other hand until slight resistance is felt. Then, twisting the container in the opposite direction, turn the breakaway ring sharply until the entire outer cap is loose and can be lifted off.

Figure 1

Connect the administration set through the sterile set port according to set instructions or remove screw cap and pour.

Figure 2

Do not warm above 150°F (66°C) to assure minimal bottle distortion. Keep bottles upright.

## **SPL Unclassified Section**

### **B. Braun Medical Inc.**

Irvine, CA 92614-5895 USA

Made in USA

Y36-002-699

### **PVP Wipe**

#### ***Active ingredient***

Povidone-iodine 10% (equivalent to 1% titratable iodine)

### **PVP Wipe**

#### ***Purpose***

First aid antiseptic

### **PVP Wipe**

#### ***Uses***

- first aid antiseptic to help prevent infection in minor cuts, scrapes and burns

### **PVP Wipe**

#### ***Warnings***

**For external use only.**

#### **Ask a doctor before use if you have**

- deep or puncture wounds
- animal bites
- serious burns

#### **Stop use and ask a doctor if**

- condition worsens or persists for more than 72 hours
- irritation and redness develops

#### **Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

### **PVP Wipe**

#### ***Directions***

- clean the affected area
- apply 1 to 3 times daily
- may be covered with a sterile bandage
- if bandaged, let dry first



- discard wipe after single use

## **PVP Wipe**

### ***Other information***

- do not use on individuals who are allergic or sensitive to iodine
- store at controlled temperature 59-86°F (15-30°C)
- do not use if pouch is open or torn

## **PVP Wipes**

### ***Inactive ingredients***

nonoxynol 9, water

## **PVP Wipe**

### ***Questions***

1-800-430-5490

## **PAWS**

### ***Active ingredient***

Ethyl alcohol 66.5%

## **PAWS**

### ***Purpose***

Antiseptic

## **PAWS**

### ***Uses***

- for handwashing to decrease bacteria on skin whenever soap and water is not readily available

## **PAWS**

### ***Warnings***

**For external use only**

**Flammable: keep away from fire or flame**

**Do not use in the eyes.**

If this happens, rinse thoroughly with water.

**Stop use and ask a doctor if**

- irritation or redness develop and persists for more than 72 hours

### **Keep out of reach of children**

If swallowed, get medical help or contact a Poison Control Center right away

### **PAWS**

#### ***Directions***

- wet hands and wrists thoroughly for 15 seconds and allow to air dry
- always reseal after use
- children under 6 years of age should be supervised when using this product

### **PAWS**

#### ***Inactive ingredients***

aloe vera, fragrance, purified water, triethanolamine

### **PAWS**

#### ***Questions***

1-800-430-5490

### **Alcohol Wipe**

#### ***Active ingredient***

Isopropyl alcohol 70%

### **Alcohol Wipe**

#### ***Purpose***

First aid antiseptic

### **Alcohol Wipe**

#### ***Uses***

first aid to help prevent infection in minor cuts, scrapes, and burns

### **Alcohol Wipe**

#### ***Warnings***

**For external use only**

**Flammable, keep away from fire and flame**

#### **Do not use**

- in or near eyes
- over large areas of the body

**Ask a doctor before use if you have**

- deep or puncture wounds
- animal bites
- serious burns

**When using this product**

- do not use longer than 1 week unless directed by a doctor

**Stop use and consult a doctor if**

- condition persists or gets worse

**Keep out of the reach of children**

If swallowed, get medical help or contact a Poison Control center right away

**Alcohol Wipe*****Directions***

- clean the affected area
- may be covered with a sterile bandage
- apply wipe to affected area 1 to 3 times daily
- discard wipe after single use

**Alcohol Wipe*****Directions***

- clean the affected area
- may be covered with a sterile bandage
- apply wipe to affected area 1 to 3 times daily
- discard wipe after single use

**Alcohol Wipe*****Other information***

- store at room temperature 15 ° to 25 ° C (59 ° to 77 ° F)
- do not use if packet is torn or opened

**Alcohol wipe*****Inactive ingredient***

water

**Alcohol Wipe*****Questions***

1-800-430-5490

## **Z346100 Kit Contents**

1 ALCOHOL PREP PADS 10P  
1 PVP IODINE WIPES 10 PER  
1 NITRILE GLOVES 2PR BBP  
1 O/H TAPE ADHESIVE TRI-CUT  
1 BK GZ 4.5"X4.1YD6PLY RL ST MSO  
1 FIRST AID GUIDE ASHI  
1 EMERGENCY SURVIVAL BLANKET  
2 GAUZE CLEAN-WRAP BDGE N/S 2"  
1 BLOODSTOPPER  
2 ABD COMBINE PAD 5" X 9"  
2 ABD PADS 8"X10" STERILE  
1 SOD. CHLORIDE 0.9% 500ML EA  
1 4OZ BFS EYEWASH TRILINGUAL BOTTLE  
1 EMPTY BAG RED 8X8X6  
1 LBL STOCK 6-3/8"X4"  
1 LBL STOCK 4"X2-7/8"  
1 LBL STOCK 3"x1-7/8"  
1 BANDAGE PACK FOR KIT  
1 ZIP LOCK BAG FOR KIT #3  
1 SELF-ADH WRAP 3 X 5 YDS NORTH REV E  
1 WATER-JEL BURN DRESSING 4 X 4  
1 CORRUGATED 24PK 01-0810 RSC  
1 TRI BNDG NON WOVEN 40"X40"X56"  
1 COLD PACK UNIT 4"X6" BULK  
1 CPR MSK,WPS,GLVS 1

## **Eyewash**

### ***Principal Display Panel***

Honeywell

TAMPER-EVIDENT CAP.  
TAPA CON SELLO DE SEGURIDAD.  
BOUCHON INDICATEUR D'INFRACTION.

eyesaline®

LAVAOJOS  
EYESALINE

Solución  
Isotónico Estéril

EYESALINE  
EYEWASH

Sterile  
Isotonic Solution

LAVAGE  
OCULAIRE  
EYESALINE

La Solution  
Isotonique Stérile

16 fl. oz. (473 mL)

NPN: 80057528



**Drug Facts** (for USA only)

Active ingredient	Purpose
Sterile water 99%	Eyewash
<b>Uses</b> for flushing the eye to remove loose foreign material, air pollutants, or chlorinated water.	
<b>Warnings</b> <b>For external use only</b> - Obtain immediate medical treatment for all open wounds in or near the eyes. To avoid contamination, do not touch tip of container to any surface. Do not reuse. Once opened, discard. <b>Do not use</b> • if solution changes color or becomes cloudy • if you have open wounds in or near the eyes, get medical help right away <b>Stop use and consult a doctor if:</b> • you experience eye pain • changes in vision • continued redness or irritation of the eye • condition worsens or persists <b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.	
<b>Directions</b> • remove contacts before using • twist top to remove • flush the affected area as needed • control rate of flow by pressure on the bottle • if necessary, continue flushing with emergency eyewash or shower	
<b>Inactive ingredients</b> sodium chloride, sodium phosphate dibasic, sodium phosphate monobasic	
<b>Questions?</b> Call 1-800-430-5490 Honeywell Safety Products USA, Inc. Smithfield, RI. 02917	

LABEL 432-004510 Rev. J REORDER / NUEVO PEDIDO / RÉAPPROVISIONNEMENT 032-000454-0000

space for lot code and supplier part number

PEEL / PELAR / PELER

**Datos de medicamento** (Para EE.UU. solamente)

Ingrediente Activo	Propósito
Agua estéril 99%	Lavaojos
<b>Usos</b> para el lavado de ojo para quitar las partículas sueltas y extrañas, los contaminantes aeros, o agua de cloruro.	
<b>Advertencias</b> <b>Para el uso externo sólo</b> - Obtenga tratamiento médico inmediato para todas las heridas abiertas en o cerca de los ojos. Para evitar la contaminación, no toque la punta del envase con ninguna superficie. No vuelva a usar. Vez abierto, descarte. <b>No se use</b> • si la solución se enturbia o cambia de color • si tiene heridas abiertas en o cerca del ojo, obtenga ayuda médica de inmediato <b>Deje de usar y consulte a un médico si:</b> • experimenta dolor de ojo • cambio de visión • rojez continuo o irritación del ojo • la condición empeora o persiste <b>Manténgase fuera del alcance de los niños.</b> En caso de ingestión accidental, obtenga atención médica o llame a un Centro de control de envenenamiento inmediatamente. <b>Instrucciones</b> • quítense los lentes de contacto antes de usar la solución • tuerza la tapa para quitar • enjuague el área afectada según se necesite • controle el chorro haciendo presión en la botella • si es necesario, sigue enjuagado con un lavaojos o ducha de emergencia <b>Ingredientes inactivos</b> cloruro de sodio, fosfato de sodio dibásico, fosfato de sodio monobásico	
<b>¿Preguntas?</b> Llame al 1-800-430-5490 Honeywell Safety Products USA, Inc. Smithfield, RI. 02917	

**Information**

<b>Usages</b> Pour le rinçage des yeux afin d'enlever un corps étranger, des polluants atmosphériques ou de l'eau chlorée.
<b>Advertissements</b> <b>Pour usage externe seulement</b> - Obtenir immédiatement des soins médicaux pour toutes les plaies ouvertes dans ou près des yeux. Pour éviter toute contamination, ne pas toucher la pointe du récipient à n'importe quelle surface. Ne pas réutiliser. Une fois ouvert, jetez-les. <b>Ne pas utiliser</b> • si la solution a changé de couleur ou si elle est brouillée • si vous avez des plaies ouvertes aux yeux ou à proximité, consultez immédiatement un médecin <b>Cesser d'utiliser la solution et consulter un médecin</b> • vous ressentez une douleur oculaire • si votre vision change • rougeur ou irritation persistante des yeux • condition empire ou persiste <b>Garder hors de la portée des enfants.</b> En cas d'ingestion, communiquer immédiatement avec un médecin ou avec un centre antipoison. <b>Mode d'emploi</b> • enlever les verres de contact avant l'utilisation • dévisser le bouchon pour l'enlever • rincer la zone touchée selon les besoins • ajuster le débit d'écoulement de la solution en augmentant ou en réduisant la pression exercée sur le contenant • si nécessaire, continuer de rincer avec une solution de rinçage oculaire d'urgence ou une douche <b>Ingédients</b> eau stérile, chlorure de sodium, phosphate dibasique de sodium, phosphate monobasique de sodium
<b>Des questions?</b> Faites le 1-800-430-5490 Honeywell Safety Products USA, Inc. Smithfield, RI. 02917

Principal Display Panel 500 ml Container

# 0.9% Sodium Chloride Irrigation USP

Lot

Isotonic Solution for Irrigation

Exp.

REF R5201-01  
NDC 0264-2201-10

**500 mL**  
PIC™ Container

Y37-002-347

Each 100 mL contains:  
Sodium Chloride USP 0.9 g  
Water for Injection USP qs  
pH adjusted with Hydrochloric  
Acid NF

pH: 5.0 (4.5–7.0)  
Calc. Osmolarity: 310 mOsmol/liter

Electrolytes (mEq/liter):  
Sodium 154 Chloride 154

Sterile, nonpyrogenic. Single unit  
container. Discard unused portion.

**Not for Injection.** Use only if solution  
is clear and container and seal are  
intact.

**Warning:** Do not warm above  
150°F (66°C).

**Recommended Storage:**  
Room temperature (25°C). Avoid  
excessive heat. Protect from freezing.  
See Package Insert.

Rx only  
PVC-free and DEHP-free

**B | BRAUN**



B. Braun Medical Inc.  
Irvine, CA 92614-5895 USA  
Made in USA

PIC is a trademark of  
B. Braun Medical Inc.

0.9% Sodium Chloride Irrigation USP



**PVP Wipe**  
**Principal Display Panel**



**PAWS**  
**Principal Display Panel**



- Kills 99.99% of Germs!
- Contains 66.5% Ethyl Alcohol
- Enriched with Aloe Vera

**Fresh Scent**

<b>Drug Facts</b>	
<b>Active ingredients</b>	<b>Purpose</b>
Ethyl Alcohol 66.5%.....	Antiseptic
<b>Uses</b>	
■ for handwashing to decrease bacteria on skin whenever soap and water is not readily available	
<b>Warnings</b>	
<b>For external use only. Flammable, keep away from fire or flame</b>	
<b>Do not use</b> in the eyes. If this happens, rinse thoroughly with water.	
<b>Stop use and ask a doctor if</b>	
■ irritation or redness develop and persists for more than 72 hours	
<b>Keep out of reach of children</b> If swallowed get medical help or contact a Poison Control Center right away	
<b>Directions</b>	
■ wet hands and wrists thoroughly for 15 seconds and allow to air dry	
■ always reseal after use	
■ children under 6 years of age should be supervised when using this product	
<b>Inactive ingredients</b>	
aloe vera, fragrance, purified water, triethanolamine	




**SAFETEC OF AMERICA, Inc.**  
887 Kensington Ave. Buffalo, NY 14215  
800-456-7077 www.safetec.com

Reorder no. 34409  
275 Gal. (1,041 L)

**Alcohol Wipe**  
**Principal Display Panel**





**4386 Kit Label**  
**Z346100**



**APPROVED**  
By Rodrigo Rosas Atilano at 2:54 pm, Mar 11, 2019

Distributed by Honeywell Safety Products USA, Inc. Smithfield, RI 02917

## 4386 FIRST AID KIT

4386 first aid kit

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0498-4386
---------------------	----------------	---------------------------	---------------

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-4386-01	1 in 1 KIT	09/13/2018	

### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	2 PACKET	0.0038 L
Part 2	10 POUCH	3 mL
Part 3	1 BOTTLE	118 mL
Part 4	10 POUCH	4 mL
Part 5	1 CONTAINER	500 mL

# Part 1 of 5

## PAWS

ethyl alcohol liquid

### Product Information

Item Code (Source)	NDC:0498-3111
Route of Administration	TOPICAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALCOHOL (UNII: 3K9958V90M) (ALCOHOL - UNII:3K9958V90M)	ALCOHOL	665 mL in 1 L

### Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
TROLAMINE (UNII: 9O3K93S3TK)	
ALOE VERA LEAF (UNII: ZY81Z83H0X)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		0.0019 L in 1 PACKET; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/21/2017	

# Part 2 of 5

## PVP IODINE WIPE

povidone-iodine 10% swab

### Product Information

**Item Code (Source)** NDC:0498-0121

**Route of Administration** TOPICAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>POVIDONE-IODINE</b> (UNII: 85H0HZU99M) (IODINE - UNII:9679TC07X4)	IODINE	10 mg in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>NONOXYNOL-9</b> (UNII: 48Q180SH9T)	
<b>WATER</b> (UNII: 059QF0KO0R)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0121-00	0.3 mL in 1 POUCH; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/18/2018	

## Part 3 of 5

## EYESALINE EMERGENCY EYEWASH

purified water liquid

### Product Information

**Item Code (Source)** NDC:0498-0100

**Route of Administration** OPHTHALMIC

Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
WATER (UNII: 059QF0KO0R) (WATER - UNII:059QF0KO0R)			WATER	98.6 mL in 100 mL
Inactive Ingredients				
Ingredient Name				Strength
SODIUM PHOSPHATE, DIBASIC (UNII: GR686LBA74)				
SODIUM PHOSPHATE, MONOBASIC, MONOHYDRATE (UNII: 593YOG76RN)				
SODIUM CHLORIDE (UNII: 451W47IQ8X)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0100-02	118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M018	12/18/2018	
Part 4 of 5				
ALCOHOL WIPE				
isopropyl alcohol swab				
Product Information				
Item Code (Source)		NDC:0498-0143		
Route of Administration		TOPICAL		
Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)			ISOPROPYL ALCOHOL	0.7 mL in 1 mL
Inactive Ingredients				
Ingredient Name			Strength	
WATER (UNII: 059QF0KO0R)				

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0143-04	0.4 mL in 1 POUCH; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/18/2018	

## Part 5 of 5

## SODIUM CHLORIDE

sodium chloride irrigant

## Product Information

Item Code (Source)	NDC:0264-2201
Route of Administration	IRRIGATION

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	0.9 g in 100 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>HYDROCHLORIC ACID</b> (UNII: QTT17582CB)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		500 mL in 1 CONTAINER; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/14/2009	
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/13/2018	

**Labeler** - Honeywell Safety Products USA, Inc. (118768815)