

**CRANE SAFETY IBUPROFEN- ibuprofen tablet, coated**  
**Crane Safety, LLC**

-----  
**Crane Safety Ibuprofen**

***Drug Facts***

***Active ingredient***

Ibuprofen 200 mg (\*NSAID)

\*nonsteroidal antiinflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

**Temporarily relieves minor aches and pains associated with**

- headache ■ toothache ■ backache ■ menstrual cramps
- common cold ■ muscular aches ■ minor arthritis pain

**Temporarily reduces fever**

***Warnings***

**Allergy alert:**

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives ■ skin reddening ■ asthma (wheezing) ■ facial swelling ■ rash ■ shock ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Heart attack or stroke warning:**

NSAIDS, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

### **Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred

speech ■ leg swelling

- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- redness or swelling is present in the painful area
- any new or unexpected symptoms occur

**If pregnant or breast feeding,**

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- do not use more than directed
- the smallest effective dose should be used
- do not take longer than 10 days, unless directed by a doctor (see Warnings)

**Adults and children:(12 years and older)**

Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used.

Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

**Children under 12 years:**

Do not give to children under 12 years of age.

***Other information***

- read all product information before using
- store at 68-77°F (20-25°C)
- avoid excessive heat 40°C (above 104°F)
- tamper evident sealed packets
- do not use any opened or torn packets

***Inactive ingredients***

carnauba wax\*, corn starch, hypromellose\*, iron oxide red, lactose\*, magnesium stearate\*, microcrystalline cellulose\*, polydextrose\*, polyethylene glycol, polyvinyl

alcohol\*, povidone K30\*, silicon dioxide, sodium starch glycolate, stearic acid, talc\*, titanium dioxide

\*may contain

## Questions or comments?

877.506.4291

## Crane Safety Ibuprofen Label

IBUPROFEN

Pain Reliever/Fever Reducer (NSAID)

CRANE Safety®

Pull To Open

This Package Is For Households Without Young Children.

Ached, Fever

Ibuprofen (NSAID) 200 mg

100 Tablets

Tamper Evident

Sealed Packets

<p><b>Drug Facts</b></p> <p><b>Active ingredient (in each tablet)</b> Purpose Ibuprofen 200 mg (NSAID)*.....Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug</p> <p><b>Uses</b> Temporarily relieves minor aches and pains associated with: ■ headache ■ toothache ■ backache ■ menstrual cramps ■ common cold ■ muscular aches ■ minor arthritis pain Temporarily reduces fever.</p> <p><b>Warnings</b> <b>Allergy alert:</b> Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ skin redness/itching ■ asthma (wheezing) ■ facial swelling ■ rash ■ shock ■ blisters If an allergic reaction occurs, stop use and seek medical help right away.</p> <p><b>Stomach bleeding warning:</b> This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)</p>	<p><b>Drug Facts (continued)</b></p> <p>■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed</p> <p><b>Heart attack and stroke warning:</b> NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p><b>Do not use</b> ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery</p> <p><b>Ask a doctor before use if</b> ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ stomach bleeding warning applies to you ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke ■ you are taking a diuretic</p> <p><b>Ask a doctor or pharmacist before use if you are</b> ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug (continued on opposite panel)</p>	<p><b>IBUPROFEN</b></p> <p>PAIN RELIEVER/FEVER REDUCER (NSAID)</p> <p><b>CRANE Safety</b></p> <p>Pull To Open</p> <p>THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.</p> <p>ACHES, FEVER IBUPROFEN (NSAID) 200 MG</p> <p>100 Tablets TAMPER EVIDENT SEALED PACKETS</p>	<p><b>Drug Facts (continued)</b></p> <p><b>When using this product</b> ■ take with food or milk if stomach upset occurs</p> <p><b>Stop use and ask a doctor if</b> ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ weakness in one part or side of body ■ slurred speech ■ fog swelling ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ redness or swelling is present in the painful area ■ any new or unexpected symptoms occur</p> <p><b>If pregnant or breast-feeding,</b> ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b> ■ do not use more than directed ■ the smallest effective dose should be used ■ do not take longer than 10 days, unless directed by a doctor (see Warnings)</p> <p><b>Adults and children (12 years and older)</b> Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.</p> <p><b>Children under 12 years:</b> Do not give to children under 12 years of age.</p> <p><b>Other information</b> ■ read all product information before using ■ store at 68°F-77°F (20°-25°C) ■ avoid excessive heat 104°F (above 40°C) ■ tamper evident sealed packets ■ do not use any opened or torn packets (continued on opposite panel)</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Inactive ingredients</b> carnauba wax*, corn starch, hydroxypropyl methylcellulose, iron oxide red, lactose*, magnesium stearate*, microcrystalline cellulose*, polydextrose*, polyethylene glycol, polyvinyl alcohol*, povidone (K-30)*, silicon dioxide, sodium starch glycolate, stearic acid, talc*, titanium dioxide *may contain</p> <p><b>Questions or comments?</b> 877.506.4291</p> <p><b>CRANE Safety</b></p> <p>Reorder CS90833</p> <p>Distributed by <b>Crane Safety</b> Houston, TX 77073 877.506.4291 www.cranesafetyco.com</p> <p>0 73408 90833 9</p>
--	---	---	---	--	---

## Crane Safety Ibuprofen Label

IBUPROFEN

Pain Reliever/Fever Reducer (NSAID)

CRANE Safety®

Pull To Open

This Package Is For Households Without Young Children.

Ached, Fever

Ibuprofen (NSAID) 200 mg

100 Tablets  
Tamper Evident  
Sealed Packets

<p><b>Drug Facts</b></p> <p><b>Active ingredient (in each tablet)</b> Purpose Ibuprofen 200 mg (NSAID) Pain reliever/fever reducer nonsteroidal anti-inflammatory drug</p> <p><b>Uses</b> Temporarily relieves minor aches and pains associated with: ■ headache ■ toothache ■ backache ■ menstrual cramps ■ common cold ■ muscular aches ■ minor arthritis pain Temporarily reduces fever.</p> <p><b>Warnings</b> <b>Allergy alert:</b> Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ skin reddening ■ asthma (wheezing) ■ facial swelling ■ rash ■ shock ■ blisters If an allergic reaction occurs, stop use and seek medical help right away. <b>Stomach bleeding warning:</b> This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)</p>	<p><b>Drug Facts (continued)</b></p> <p>■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed</p> <p><b>Heart attack and stroke warning:</b> NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p><b>Do not use</b> ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery</p> <p><b>Ask a doctor before use if</b> ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke ■ you are taking a diuretic</p> <p><b>Ask a doctor or pharmacist before use if you are</b> ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug. (continued on opposite panel)</p>	<p><b>IBUPROFEN</b></p> <p>PAIN RELIEVER/FEVER REDUCER (NSAID)</p> <p><b>CRANE Safety</b></p> <p>Pull to Open</p> <p>THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.</p> <p>ACHES, FEVER IBUPROFEN (NSAID) 200 MG</p> <p>100 Tablets TAMPER EVIDENT SEALED PACKETS</p>	<p><b>Drug Facts (continued)</b></p> <p><b>When using this product</b> ■ take with food or milk if stomach upset occurs</p> <p><b>Stop use and ask a doctor if</b> ■ you experience any of the following signs of stomach bleeding: ■ black, tarry ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ big swelling ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ redness or swelling is present in the painful area ■ any new or unexpected symptoms occur</p> <p><b>If pregnant or breast-feeding,</b> ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b> ■ do not use more than directed ■ the smallest effective dose should be used ■ do not take longer than 10 days, unless directed by a doctor (see Warnings)</p> <p><b>Adults and children (12 years and older)</b> Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.</p> <p><b>Children under 12 years:</b> Do not give to children under 12 years of age.</p> <p><b>Other information</b> ■ read all product information before using ■ store at 68-77°F (20-25°C) ■ avoid excessive heat 104°F (above 40°C) ■ tamper evident sealed packets ■ do not use any opened or torn packets (continued on opposite panel)</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Inactive ingredients</b> carnauba wax, corn starch, hypromellose, iron oxide red, lactose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, polyvinyl alcohol, povidone K30, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide *may contain</p> <p><b>Questions or comments?</b> 877.506.4291</p> <p><b>CRANE Safety</b></p> <p>Reorder C590833 Distributed by Crane Safety Houston, TX 77073 877.506.4291 www.cranesafetyco.com</p>
---	--	---	--	---	--

# CRANE SAFETY IBUPROFEN

ibuprofen tablet, coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:73408-709
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>IBUPROFEN</b> (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	

## Product Characteristics

<b>Color</b>	red (Reddish Brown)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	G;2
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73408-709-33	50 in 1 BOX	01/17/2020	12/31/2023
1		2 in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079174	01/17/2020	12/31/2023

## CRANE SAFETY IBUPROFEN

ibuprofen tablet, coated

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73408-609
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

### Inactive Ingredients

Ingredient Name	Strength
POLYDEXTROSE (UNII: VH2XOU12IE)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6130)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STARCH, CORN (UNII: O8232NY3SJ)	

### Product Characteristics

<b>Color</b>	brown	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;291
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73408-609-33	50 in 1 BOX	01/17/2020	12/31/2023
1		2 in 1 PACKET; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	01/17/2020	12/31/2023

**Labeler** - Crane Safety, LLC (080998015)

**Registrant** - Unifirst First Aid Corporation (832947092)

Revised: 3/2022

Crane Safety, LLC