

**CRANE SAFETY BISMUTH- bismuth subsalicylate tablet, chewable**  
**Crane Safety LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Crane Safety Bismuth**

***Drug Facts***

***Active ingredient (in each tablet)***

Bismuth Subsalicylate 262 mg

(each tablet contains 102 mg salicylate)

***Purpose***

Upset stomach reliever/antidiarrheal

***Uses***

**Temporarily relieves**

- traveler's diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn
- indigestion
- nausea
- gas
- belching
- fullness

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome a rare but serious illness.

**Allergy alert:** Contains salicylate. Do not take if you are:

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use if you have**

- bloody or black stool
- an ulcer
- a bleeding problem

**Ask a doctor before use if you have**

- fever
- mucus in the stool

**Ask a doctor or pharmacist if you are taking any drug for**

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

**Stop use and ask a doctor if**

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

**When using this product**

a temporary and harmless darkening of the tongue and/or stool may occur. Stool darkening should not be confused with melena.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep this and all drugs out of reach of children.**

In case of accidental overdose, contact a physician or poison control center immediately.

***Directions***

- **do not use more than directed**
- **chew or crush tablets completely before swallowing**
- **do not swallow tablet whole**
- use until diarrhea stops but not more than 2 days
- drink plenty of clear fluids to help prevent dehydration, which may accompany diarrhea
- do not exceed 16 tablets in 24 hours

**Adults and children: (12 years and older)** Chew 2 tablets every 1/2 to 1 hour or 4 tablets every hour as needed

**Children under 12 years:** Do not give to children under 12 years of age.

***Other information***

- phenylketonurics: contains phenylalanine 1.1mg per tablet
- each tablet contains 73mg elemental calcium
- store at room temperature 59°-86°F (15°-30°C)
- tamper-evident sealed packets
- do not use any opened or torn packets

***Inactive ingredients***

acacia gum, aspartame, calcium carbonate, D&C red #27, dextrans, flavoring, magnesium stearate, maltodextrin, microcrystalline cellulose, silicon dioxide

***Questions or comments?***

877.506.4291

**Crane Safety Bismuth Label**

Chewable

Bismuth Tabs

Upset Stomach/Diarrhea Relief

CRANE Safety®

Bismuth Subsalicylate 262 MG

Chew or Crush Tablets Completely Before Swallowing.

Do Not Swallow Tablets Whole.

36 Tablets

Tamper Evident

### Drug Facts

**Active ingredient (in each tablet)** **Purpose**  
Bismuth Subsalicylate 262 mg .....Upset stomach reliever/antidiarrheal (each tablet contains 102 mg salicylate)

**Uses**  
Temporarily relieves ■ travelers' diarrhea ■ diarrhea  
■ upset stomach due to overindulgence in food and drink, including:  
■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome a rare but serious illness.

**Allergy alert:** Contains salicylate. Do not take if you are: ■ allergic to salicylates (including aspirin) ■ taking other salicylate products

**Do not use if you have**  
■ bloody or black stool ■ an ulcer ■ a bleeding problem

**Ask a doctor before use if you have** ■ fever ■ mucus in the stool

**Ask a doctor or pharmacist if you are taking any drug for**  
■ anticoagulation (thinning of the blood) ■ diabetes ■ gout ■ arthritis

**Stop use and ask a doctor if** ■ symptoms get worse ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days

**When using this product** a temporary and harmless darkening of the tongue and/or stool may occur. Stool darkening should not be confused with melena.

**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep this and all drugs out of reach of children.** In case of accidental overdose, contact a physician or poison control center immediately. ▶

### Drug Facts (continued)

**Directions**  
■ do not use more than directed  
■ chew or crush tablets completely before swallowing  
■ do not swallow tablets whole  
■ use until diarrhea stops but not more than 2 days  
■ drink plenty of clear fluids to help prevent dehydration, which may accompany diarrhea  
■ do not exceed 16 tablets in 24 hours

<b>Adults and children: (12 years and older)</b>	Chew 2 tablets every 1/2 to 1 hour or 4 tablets every hour as needed
<b>Children under 12 years:</b>	Do not give to children under 12 years of age.

**Other information**  
■ phenylketonurics: contains phenylalanine 1.1 mg per tablet  
■ each tablet contains 73 mg elemental calcium  
■ store at room temperature 59°-86°F (15°-30°C)  
■ tamper-evident sealed packets  
■ do not use any opened or torn packets  
*(continued on opposite panel)*

# CHEWABLE BISMUTH TABS

UPSET STOMACH / DIARRHEA RELIEF

BISMUTH SUBSALICYLATE 262 MG

CHEW OR CRUSH TABLETS COMPLETELY BEFORE SWALLOWING. DO NOT SWALLOW TABLETS WHOLE.

36 Tablets

TAMPER EVIDENT

### Drug Facts (continued)

**Inactive ingredients**  
acacia gum, aspartame, calcium carbonate, D&C red #27, dextrates, flavoring, magnesium stearate, maltodextrin, microcrystalline cellulose, silicon dioxide

**Questions or comments?**  
877.506.4291

Reorder CS41220

Distributed by  
**Crane Safety**  
Houston, TX 77073  
877.506.4291  
www.cranesafetyco.com

0 73408 41220 1

## CRANE SAFETY BISMUTH

bismuth subsalicylate tablet, chewable

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:73408-406
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	262 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>D&amp;C RED NO. 27</b> (UNII: 2LRS185U6K)	
<b>CALCIUM CARBONATE</b> (UNII: H0G9379FGK)	
<b>DEXTRATES</b> (UNII: G263MI44RU)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>ACACIA</b> (UNII: 5C5403N26O)	
<b>ASPARTAME</b> (UNII: Z0H242BBR1)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	

### Product Characteristics

<b>Color</b>	pink	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	16mm

<b>Flavor</b>		<b>Imprint Code</b>	RH;046	
<b>Contains</b>				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73408-406-20	6 in 1 BOX	01/17/2020	11/29/2024
1		6 in 1 CELLO PACK; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part335	01/17/2020	11/29/2024	

**Labeler** - Crane Safety LLC (080998015)

**Registrant** - Unifirst First Aid Corporation (832947092)

Revised: 6/2022

Crane Safety LLC