

CRANE SAFETY XS PAIN RELIEF- acetaminophen, aspirin, salicylamide and caffeine tablet

Crane Safety LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Crane Safety Extra Strength Pain Relief

Drug Facts

Active ingredients (in each tablet)

Acetaminophen 110 mg

Aspirin (NSAID)* 162 mg

Caffeine 32.4 mg

Salicylamide (NSAID)* 152 mg

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Pain reliever/fever reducer

Pain reliever

Pain reliever aid

Uses

temporarily relieves minor aches and pains associated with

- headache
- muscular aches
- minor arthritis pain
- back pain
- colds
- toothache
- menstrual cramps

Temporarily reduces fever.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these

symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin and Salicylamide may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin or allergic reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours, which is maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains nonsteroidal anti-inflammatory drugs (NSAIDs), which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for pain for more than 10 days unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor
- with any other pain relievers/ fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems (such as heartburn, upset stomach or stomach pain)
- you have heart disease, liver cirrhosis, or kidney disease

- you have asthma or high blood pressure
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- anticoagulation
- diabetes
- gout
- arthritis

Ask a doctor or pharmacist if you have

- liver disease

When using this product do not exceed recommended dose

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- symptoms do not improve
- pain or fever gets worse
- ringing in the ears or loss of hearing occurs
- new symptoms occur

If pregnant or breast-feeding, ask a health care professional before use. It is especially important not to use aspirin or salicylamide during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Overdose warning:

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**

Adults and children: (12 years and older)

Take 2 tablets every 4 hours as needed with a full glass of water. Do not exceed 8 tablets in 24 hours or as directed by a doctor.

Children under 12 years:

Ask a doctor

Other information

- store at room temperature 59°-86°F (15°-30°C)

- avoid excessive heat and humidity
- tamper evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

FD&C yellow #6, magnesium stearate, microcrystalline cellulose, povidone, starch, stearic acid

Questions or comments 877-506.4291

Crane Safety Extra Strength Pain Relief Label

Extra Strength

Pain Relief

Fast Acting Advanced Formula

CRANE Safety®

Pull to Open

Aches, Fever

Acetaminophen 110 MG/

Aspirin (NSAID) 162 MG

Pain Reliever

Salicylamide (NSAID) 152 MG

Pain Reliever Aid

Caffeine 32.4 MG

This Package is For Households Without Young Children.

100 Tablets

Tamper Evident

Sealed Packets

EXTRA STRENGTH PAIN RELIEF

FAST ACTING ADVANCED FORMULA



Pull to Open

ACHES, FEVER
ACETAMINOPHEN 110 MG/
ASPIRIN (NSAID) 162 MG
PAIN RELIEVER
SALICYLAMIDE (NSAID) 152 MG
PAIN RELIEVER AID
CAFFEINE 32.4 MG

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

100 TABLETS

TAMPER EVIDENT SEALED PACKETS

Retain carton for complete product information

Drug Facts

Active ingredient (in each tablet) **Purposes**
Acetaminophen 110 mg Pain reliever/fever reducer
Aspirin (NSAID) 162 mg Pain reliever/fever reducer
Caffeine 32.4 mg Adjuvant
Salicylamide (NSAID) 152 mg Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses
temporarily relieves minor aches and pains associated with
■ headaches ■ muscular aches ■ minor arthritis pain
■ back pain ■ colds ■ toothache ■ menstrual cramps
temporarily reduces fever.

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin and Salicylamide may cause a severe allergic reaction which may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin or allergic reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 4,000 mg of acetaminophen in 24 hours, which is the maximum daily amount

■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains nonsteroidal anti-inflammatory drugs (NSAIDs), which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug

Drug Facts (continued)

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ for pain for more than 10 days unless directed by a doctor
■ for more than 3 days for fever unless directed by a doctor
■ with any other pain reliever/fever reducer

Ask a doctor before use if
■ stomach bleeding warning applies to you
■ you have a history of stomach problems (such as heartburn, upset stomach or stomach pain)
■ you have heart disease, liver cirrhosis, or kidney disease
■ you have asthma or high blood pressure
■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for
■ anticoagulation ■ diabetes ■ gout ■ arthritis
Ask a doctor or pharmacist before use if you have
■ liver disease

When using this product do not exceed recommended dose
Stop use and ask a doctor if
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ symptoms do not improve
■ pain or fever gets worse
■ ringing in the ears or loss of hearing occurs
■ new symptoms occur

Drug Facts (continued)

If pregnant or breast-feeding, ask a health care professional before use. It is especially important not to use aspirin or salicylamide during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ do not use more than directed

Adults and children: (12 years and older) Take 2 tablets every 4 hours as needed with a full glass of water. Do not exceed 8 tablets in 24 hours or as directed by a doctor.

Children under 12 years: Ask a doctor.

Other information

■ store at room temperature 59°-86°F (15°-30°C)
■ avoid excessive heat and humidity
■ tamper evident sealed packets
■ do not use any opened or torn packets

Inactive ingredients

FD&C yellow #6, magnesium stearate, microcrystalline cellulose, povidone, starch, stearic acid

Questions or comments? 877.506.4291



0 73408 91133 9

Reorder CS91133

Distributed by
CRANE Safety
Houston, TX 77073
877.506.4291
www.cranesafetyco.com

CRANE SAFETY XS PAIN RELIEF

acetaminophen, aspirin, salicylamide and caffeine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73408-911
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SALICYLAMIDE (UNII: EM8BM710ZC) (SALICYLAMIDE - UNII:EM8BM710ZC)	SALICYLAMIDE	152 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	110 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	32.4 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	162 mg

Inactive Ingredients

Ingredient Name	Strength
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
MAGNESIUM STEARATE (UNII: 70097M6130)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	orange	Score	no score
Shape	ROUND	Size	11mm

Flavor		Imprint Code	FR;2	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73408-911-33	50 in 1 BOX	01/17/2020	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph not final	part343	01/17/2020		

Labeler - Crane Safety LLC (080998015)

Registrant - Unifirst First Aid Corporation (832947092)

Revised: 3/2022

Crane Safety LLC