

**BASIC CARE DAYTIME SEVERE COLD AND FLU- acetaminophen,  
dextromethorphan hydrobromide, guaifenesin, phenylephrine  
hydrochloride solution  
Amazon.com Services LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Amazon Daytime Severe Cold & Flu Drug Facts**

**Active ingredients (in each 15 mL)**

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

**Purpose**

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

**Uses**

temporarily relieves common cold/flu symptoms:

- sinus congestion and pressure
- nasal congestion
- minor aches and pains
- headache
- sore throat
- fever
- cough due to minor throat and bronchial irritation
- reduces swelling of nasal passages
- promotes nasal and/or sinus drainage
- temporarily restores freer breathing through the nose
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- diabetes
- thyroid disease
- cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

### **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

### **When using this product**

**do not use more than directed**

## Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

## If pregnant or breast-feeding,

ask a health professional before use.

## Keep out of reach of children.

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

|                                 |                   |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 4 hrs |
| children 6 to under 12 yrs      | 15 mL every 4 hrs |
| children 4 to under 6 yrs       | ask a doctor      |
| children under 4 yrs            | do not use        |

## Other information

- **each 15 mL contains:** sodium 15 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

## Inactive ingredients

anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C green #3, FD&C red #40, FD&C yellow #6, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum

## Questions?

**1-800-719-9260**

## Package/Label Principal Display Panel

amazon

basic care

Multi-Symptom Relief

Compare to Vicks<sup>®</sup> DayQuil<sup>®</sup> Severe Honey Flavor active ingredients

Non-Drowsy

Daytime Severe Cold & Flu

Acetaminophen, Phenylephrine HCl, Dextromethorphan HBr, Guaifenesin

Pain Reliever, Fever Reducer, Cough Suppressant, Expectorant, Nasal Decongestant

Max Strength

Honey Flavor

12 FL OZ (355 mL)

amazon

basic+care

Multi-Symptom Relief

Compare to Vicks® DayGull® Severe Honey Flavor active ingredients\*

NDC 72288-515-40

Non-Drowsy

# Daytime Severe Cold & Flu

Acetaminophen, Phenylephrine HCl, Dextromethorphan HBr, Guaifenesin

Pain Reliever, Fever Reducer, Cough Suppressant, Expectorant, Nasal Decongestant

Max Strength

Honey Flavor

12 FL OZ (355 mL)

: 51U40 BN F1

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

DISTRIBUTED BY:  
AMAZON.COM SERVICES LLC  
410 TERRY AVENUE N., SEATTLE, WA 98109  
© 2022 AMAZON.COM, INC. OR IT'S AFFILIATES. ALL RIGHTS RESERVED.

**PARENTS:**

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

GLUTEN FREE

### Drug Facts

**Active ingredients (in each 15 mL)**

| Active ingredients (in each 15 mL) | Purpose                     |
|------------------------------------|-----------------------------|
| Acetaminophen 325 mg               | Pain reliever/fever reducer |
| Dextromethorphan HBr 10 mg         | Cough suppressant           |
| Guaifenesin 200 mg                 | Expectorant                 |
| Phenylephrine HCl 5 mg             | Nasal decongestant          |

**Uses** temporarily relieves common cold/flu symptoms: ■ sinus congestion and pressure ■ nasal congestion ■ minor aches and pains ■ headache ■ sore throat ■ fever ■ cough due to minor throat and bronchial irritation ■ reduces swelling of nasal passages ■ promotes nasal and/or sinus drainage ■ temporarily restores freer breathing through the nose ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away

PEEL BACK AT CORNER FOR MORE INFORMATION

: 51U40 BN B1



### Drug Facts (continued)

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

ADHESIVE AREA • NO VARNISH • NO TYPE

### Drug Facts (continued)

**Stop use and ask a doctor if** ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults) ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.


If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ take only as directed – see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

|                                 |                   |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 4 hrs |
| children 6 to under 12 yrs      | 15 mL every 4 hrs |
| children 4 to under 6 yrs       | ask a doctor      |

ADHESIVE AREA • NO VARNISH • NO TYPE



## BASIC CARE DAYTIME SEVERE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution

### Product Information

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:72288-515 |
| <b>Route of Administration</b> | ORAL           |                           |               |

### Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength             | Strength        |
|--|-------------------------------|-----------------|
| <b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)                    | ACETAMINOPHEN                 | 325 mg in 15 mL |
| <b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 10 mg in 15 mL  |
| <b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)                        | GUAIFENESIN                   | 200 mg in 15 mL |
| <b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)      | PHENYLEPHRINE HYDROCHLORIDE   | 5 mg in 15 mL   |

### Inactive Ingredients

| Ingredient Name  | Strength |
|--|----------|
| <b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)            |          |
| <b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)            |          |
| <b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)                 |          |
| <b>FD&amp;C GREEN NO. 3</b> (UNII: 3P3ONR6O1S)             |          |
| <b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)              |          |
| <b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)            |          |
| <b>GLYCERIN</b> (UNII: PDC6A3C0OX)                         |          |
| <b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A) |          |
| <b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)                 |          |
| <b>WATER</b> (UNII: 059QF0KO0R)                            |          |

|  |  |
|--|--|
| <b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)                 |  |
| <b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)                  |  |
| <b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)                  |  |
| <b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR) |  |
| <b>SORBITOL</b> (UNII: 506T60A25R)                         |  |
| <b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)                        |  |
| <b>XANTHAN GUM</b> (UNII: TTV12P4NEE)                      |  |

### Product Characteristics

|                 |                |                     |  |
|-----------------|----------------|---------------------|--|
| <b>Color</b>    | YELLOW         | <b>Score</b>        |  |
| <b>Shape</b>    |                | <b>Size</b>         |  |
| <b>Flavor</b>   | HONEY, MENTHOL | <b>Imprint Code</b> |  |
| <b>Contains</b> |                |                     |  |

### Packaging

| # | Item Code        | Package Description                                   | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:72288-515-40 | 355 mL in 1 BOTTLE; Type 0: Not a Combination Product | 04/22/2022           |                    |

### Marketing Information

| Marketing Category  | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part341                                  | 04/22/2022           |                    |

**Labeler** - Amazon.com Services LLC (128990418)

Revised: 4/2022

Amazon.com Services LLC