

**KROGER REGULAR STRENGTH PINK BISMUTH- bismuth  
subsalicylate suspension  
KROGER COMPANY**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Kroger Regular Strength Pink Bismuth Subsalicylate 525 mg Drug Facts**

**Active ingredient (in each 30 mL dose)**

Bismuth subsalicylate 525 mg

**Purposes**

Upset stomach reliever and antidiarrheal

**Uses**

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn
- indigestion
- nausea
- gas
- belching
- fullness

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea or vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use if you have**

- an ulcer
- a bleeding problem

- bloody or black stool

### **Ask a doctor before use if you have**

- fever
- mucus in the stool

### **Ask a doctor or pharmacist before use if you are**

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

### **When using this product**

a temporary, but harmless, darkening of the stool and/or tongue may occur

### **Stop use and ask a doctor if**

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away at (1-800-222-1222).

### **Directions**

- shake well before use
- only use dose cup provided
- **adults and children 12 years and over:** 30 mL 1 (dose) every 1/2 hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
- 30 mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (240 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- **children under 12 years:** ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

### **Other information**

- **each 30 mL contains:** sodium 10 mg

- **salicylate 261 mg**
- low sodium
- sugar free
- store at room temperature
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

### **Inactive ingredients**

carboxymethylcellulose sodium, D&C Red No. 22, D&C Red No. 28, flavor, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum

### **Questions or comments?**

**1-866-467-2748**

**DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN.**

### **Principal Display Panel**

**\*COMPARE TO the active ingredient in PEPTO-BISMOL® REGULAR STRENGTH**

**\*See side panel**

Regular Strength

### **Stomach Relief**

Bismuth Subsalicylate 525 mg

Upset Stomach Reliever/Antidiarrheal

### **Original Flavor**

### **5 SYMPTOM RELIEF**

#### **RELIEF:**

- Heartburn
- Indigestion
- Nausea
- Upset Stomach
- Diarrhea

**\*This product is not manufactured or distributed by Procter & Gamble, distributor of Pepto-Bismol® Regular Strength.**

**Package Label 236 mL**

**Drug Facts (continued)**

**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.


**Directions** ■ shake well before use ■ only use dose cup provided ■ **adults and children 12 years and over:** ■ 30 mL (1 dose) every ½ hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea ■ 30 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) ■ do not exceed 8 doses (240 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ **children under 12 years:** ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information** ■ each 30 mL contains: sodium 10 mg ■ salicylate 261 mg ■ low sodium ■ sugar free ■ store at room temperature ■ protect from freezing ■ avoid excessive heat (more than 104°F or 40°C)

**Inactive ingredients** carboxymethylcellulose sodium, D&C Red No. 22, D&C Red No. 28, flavor, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum

**Questions or comments?** 1-800-632-6900

COMPARE TO the active ingredient in  
**PEPTO-BISMOL® REGULAR STRENGTH** \*See side panel



NDC 30142-608-08

**Regular Strength**

# Stomach Relief

**Bismuth Subsalicylate 525 mg Upset Stomach Reliever/Antidiarrheal**

**5 SYMPTOM RELIEF:**  
Heartburn,  
Indigestion, Nausea,  
Upset Stomach, Diarrhea

**8 FL OZ (236 mL)**



**Drug Facts**

**Active ingredient (in each 30 mL dose)**  
Bismuth subsalicylate 525 mg .....Upset stomach reliever and antidiarrheal

**Purposes**  
**Uses** relieves ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including:  
■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Contains salicylate. Do not take if you are  
■ allergic to salicylates (including aspirin)  
■ taking other salicylate products

**Do not use if you have** ■ an ulcer ■ a bleeding problem ■ bloody or black stool

**Ask a doctor before use if you have** ■ fever ■ mucus in the stool

**Ask a doctor or pharmacist before use if you are taking any drug for** ■ anticoagulation (thinning the blood) ■ diabetes ■ gout ■ arthritis

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur.

**Stop use and ask a doctor if** ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days

\*Pepto-Bismol® Regular Strength is a registered trademark of Procter & Gamble, Cincinnati, OH 45202. Procter & Gamble is not affiliated with The Kroger Co. or this product.

LOT: UNVARNISHED AREA EXP: 70608KRQLR

DIST. BY THE KROGER CO. CINCINNATI, OHIO 45202  
QUALITY GUARANTEE 800-632-6900 | www.kroger.com

0 41260 01007 1

## Package Label 354 mL

**Drug Facts (continued)**

**Directions** ■ shake well before use ■ only use dose cup provided ■ **adults and children 12 years and over:** ■ 30 mL (1 dose) every ½ hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea ■ 30 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) ■ do not exceed 8 doses (240 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ **children under 12 years:** ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information** ■ each 30 mL contains: sodium 10 mg ■ salicylate 261 mg ■ low sodium ■ sugar free ■ store at room temperature ■ protect from freezing ■ avoid excessive heat (more than 104°F or 40°C)

**Inactive ingredients** carboxymethylcellulose sodium, D&C Red No. 22, D&C Red No. 28, flavor, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum

**Questions or comments?** 1-800-632-6900

COMPARE TO the active ingredient in  
**PEPTO-BISMOL® Regular Strength** \*See side panel



NDC 30142-608-12

**Regular Strength**

# Stomach Relief

**Bismuth Subsalicylate 525 mg Upset Stomach Reliever/Antidiarrheal**

**5 SYMPTOM RELIEF:**  
Heartburn,  
Indigestion, Nausea,  
Upset Stomach, Diarrhea

**12 FL OZ (354 mL)**



**Drug Facts**

**Active ingredient (in each 30 mL dose)**  
Bismuth subsalicylate 525 mg .....Upset stomach reliever and antidiarrheal

**Purposes**  
**Uses** relieves ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including:  
■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Contains salicylate. Do not take if you are  
■ allergic to salicylates (including aspirin)  
■ taking other salicylate products

**Do not use if you have** ■ an ulcer ■ a bleeding problem ■ bloody or black stool

**Ask a doctor before use if you have** ■ fever ■ mucus in the stool

**Ask a doctor or pharmacist before use if you are taking any drug for** ■ anticoagulation (thinning the blood) ■ diabetes ■ gout ■ arthritis

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur.

**Stop use and ask a doctor if** ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days

**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.

LOT: UNVARNISHED AREA EXP: 70612KRQLR

DISTRIBUTED BY THE KROGER CO. CINCINNATI, OHIO 45202  
QUALITY GUARANTEE 800-632-6900 | www.kroger.com

0 41260 01002 6

## KROGER REGULAR STRENGTH PINK BISMUTH

bismuth subsalicylate suspension

### Product Information

**Product Type**

HUMAN OTC DRUG

**Item Code (Source)**

NDC:30142-608

Route of Administration		ORAL		
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ)		BISMUTH SUBSALICYLATE	525 mg in 30 mL	
Inactive Ingredients				
Ingredient Name			Strength	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)				
D&C RED NO. 22 (UNII: 1678RKX8RT)				
D&C RED NO. 28 (UNII: 767IPOY5NH)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)				
POTASSIUM SORBATE (UNII: 1VPU26JZZ4)				
WATER (UNII: 059QF0KO0R)				
SALICYLIC ACID (UNII: O414PZ4LPZ)				
DIMETHICONE (UNII: 92RU3N3Y1O)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
XANTHAN GUM (UNII: TTV12P4NEE)				
Product Characteristics				
Color	PINK (viscous)	Score		
Shape		Size		
Flavor		Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:30142-608-08	236 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/30/2020	
2	NDC:30142-608-12	354 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/30/2020	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC monograph final	part335		03/30/2020	

**Labeler** - KROGER COMPANY (006999528)

