HEALTH MART COUGH DM- dextromethorphan polistirex suspension Strategic Sourcing Services LLC

McKesson Cough DM Drug Facts

Active ingredient (in each 5 mL)

Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide

Purpose

Cough suppressant

Uses

temporarily relieves

- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the impulse to cough to help you get to sleep

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Allergy Alert: Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions.

Ask a doctor before use if you have

- chronic cough that lasts as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if

- side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
- cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- shake bottle well before use
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by doctor

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24
	hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24
	hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24
	hours
children under 4 years of age	do not use

Other information

- each 5 mL contains: sodium 5 mg
- store at 20° to 25°C (68° to 77°F)
- dosing cup provided

Inactive Ingredients

D&C Red #30 aluminum lake, D&C Yellow #10 aluminum lake, glycerin, high fructose corn syrup, methylparaben, natural and artificial orange flavor, polysorbate 80, polyvinyl acetate, povidone, propylparaben, purified water, sodium metabisulfite, sodium polystyrene sulfonate, sucrose, tartaric acid, tragacanth gum, triacetin, xanthan gum

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Compare to Delsym® active ingredient

12 HOUR COUGH RELIEF

Cough DM

Dextromethorphan Polistirex

Extended-Release Oral Suspension

Cough Suppressant

Day or night

Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions

Dosing cup included

Alcohol Free

Gluten Free

ORANGE-FLAVORED LIQUID

3 Fl oz (89 mL)



Drug Facts

Active ingredient (in each 5 mL) Purpose Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide... ...Cough suppress

Uses temporarily relieves
■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants ■ the impulse to cough to help you get to sleep

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MADI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MADI drug, I you do not know if your prescription drug contains an MADI, ask a doctor or pharmacist before taking this product.

Allengy Alert Contains sodium metabisulfite, a suffile that may cause

llergic-type reactions.

airrigue, type reacours.

Ask a doctor before use if you have

chronic cough that lasts as occurs with smoking, asthma or emphysema

cough that occurs with too much philegm (mucus)

Stop use and ask a doctor if

stop use and ask a doctor if

side effects to FDA at 1-800-FDA-1088.

cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or
contact a Poison Control Center right away (1-800-222-1222).

- shake bottle well before use
- I measure only with dosing cup provided I do not use dosing cup with other products I dose as follows or as directed by a doctor

adults and children	10 mL every 12 hours,
12 years of age and over	not to exceed 20 mL in 24 hours
children 6 to under	5 mL every 12 hours,
12 years of age	not to exceed 10 mL in 24 hours
children 4 to under	2.5 mL every 12 hours,
6 years of age	not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information

■ each 5 mL contains: sodium 5 mg
■ store at 20° to 25°C (68° to 77°F) ■ dosing cup provided

Drug Facts (continued)

Drug Facts (Continued)
Inactive ingredients D&C Red #30
aluminum lake, D&C Yellow #10 aluminum lake,
glycein, high fructose com syrup, methylparaben,
natural and artificial orange flavor, polysorbate 80,
polyvinyl acetate, povidone, propylparaben,
purified water, socilum metabisulifie, sodium
polystyrene sulfornate, sucrose, tratrario acid,
tragacanth gum, triacetin, xanthan gum

Questions or comments? 1-800-719-9260

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Delsym®.

DOSING

SHAKE WELL BEFORE USE.

Measure only with doeing cup provided.

Do not use doeing cup with other products. Dose 12 years to adult 10 mL EVERY 12 HOURS 6 to under 12 5 mL EVERY 12 HOURS 4 to 2.5 mL EVERY 12 HOURS under 6

nel for full dosing directions







12 HOUR COUGH RELIEF

Cough DM

Dextromethorphan Polistirex **Extended-Release Oral Suspension** Cough Suppressant

Day or night

metabisulfite, a sulfite that may cause allergic-type reactions

Dosing cup included

Alcohol Free Gluten Free



ACTUAL SIZE





3 Fl oz (89 mL)

ORANGE-FLAVORED LIQUID

MSKESSON

HEALTH MART COUGH DM

dextromethorphan polistirex suspension

Product Information

Product Type HUMAN OTC DRUG NDC:62011-0176 Item Code (Source)

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength
	DEXTRO METHO RPHAN HYDRO BRO MIDE (UNII: 9 D2RTI9 KYH)	DEXTROMETHORPHAN	30 mg
	(DEXTROMETHORPHAN - UNII:7355X3ROTS)	HYDROBROMIDE	in 5 mL

Inactive Ingredients		
Ingredient Name	Strength	
POLISTIREX (UNII: 5H9 W9 GTW27)		
D&C RED NO. 30 (UNII: 2S42T2808B)		
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)		
GLYCERIN (UNII: PDC6 A3C0 OX)		
HIGH FRUCTO SE CORN SYRUP (UNII: XY6 UN3QB6S)		
METHYLPARABEN (UNII: A2I8C7HI9T)		
POLYSORBATE 80 (UNII: 6OZP39ZG8H)		
POLYVINYL ACETATE (UNII: 32K497ZK2U)		
PO VIDO NE, UNSPECIFIED (UNII: FZ989GH94E)		
PROPYLPARABEN (UNII: Z8 IX2SC1OH)		
WATER (UNII: 059QF0KO0R)		
SO DIUM METABISULFITE (UNII: 4VON5FNS3C)		
SODIUM POLYSTYRENE SULFONATE (UNII: 1699G8679Z)		
SUCROSE (UNII: C151H8M554)		
TARTARIC ACID (UNII: W48881119H)		
TRAGACANTH (UNII: 2944357020)		
TRIACETIN (UNII: XHX3C3X673)		
XANTHAN GUM (UNII: TTV12P4NEE)		

Product Characteristics			
Color	ORANGE	Score	
Shape		Size	
Flavor	ORANGE	Imprint Code	
Contains			

	Packaging			
	# Item Code	Package Description	Marketing Start Date	Marketing End Date
ı	1 NDC:62011-0176-1	1 in 1 CARTON	08/28/2012	
ı	1	89 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091135	08/28/2012	

Labeler - Strategic Sourcing Services LLC (116956644)