## MUCUS RELIEF SEVERE CONGESTION AND COUGH MAXIMUM STRENGTHdextromethorphan hbr, guaifenesin, phenylephrine hcl liquid Freds Inc

-----

## **Drug Facts**

## Active ingredients (in each 20 mL)

Dextromethorphan HBr 20 mg Guaifenesin 400 mg Phenylephrine HCl 10 mg

## **Purposes**

Cough suppressant

Expectorant

Nasal Decongestant

### Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
  - the intensity of coughing
  - the impulse to cough to help you get to sleep
  - $\circ \;$  nasal congestion due to a cold

# Warnings

### Do not use

- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

# Ask a doctor before use if you have

- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

## When using this product,

do not use more than directed.

## Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- symptoms do not get better within 7 days or occur with a fever
- cough comes back, or occurs with fever, rash or headache that lasts

These could be signs of a serious condition.

## If pregnant or breast-feeding,

ask a health professional before use.

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### **Directions**

- do not take more than 6 doses in a 24-hour period
- measure only with dosing cup provided Do not use any other dosing device.
- keep dosing cup with product
- mL=milliliter
- shake well before using
- adults and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

### Other information

- each 20 mL contains: sodium 17 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

# Inactive ingredients

citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, sodium citrate, xanthan gum

# **Principal Display Panel**

Maximum Strength

### **Mucus Relief**

Severe Congestion and Cough

Dextromethorphan HBr-Cough suppressant-controls cough

Guaifenesin-Expectorant- thins mucus and relieves chest congestion

Phenylephrine HCI-nasal decongestant- relieves stuffy nose

ages 12 & over

FL OZ (mL)

COMPARE TO the active Ingredients of Maximum Strength Mucinex® Fast-Max® Severe Congestion & Cough\*

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Severe Congestion & Cough.

# TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY: fred's Inc.

4300 NEW GETWELL RD,

MEMPHIS, TN 38118

www.fredsinc.com

Questions or comments: 1-855-331-3733

## Package Label



TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Severe Congestion & Cough.



PLD-C283D LB003126

# Drug Facts

## Active ingredients (in each 20 mL)

Dextromethorphan HBr 20 mg......Cough suppressant
Guaifenesin 400 mg.....Expectorant

Phenylephrine HCl 10 mg......Nasal decongestant

PEEL CORNER FOR MORE DRUG FACTS

**Purposes** 

## Drug Facts (continued)

### Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the intensity of coughing
- the impulse to cough to help you get to sleep
- nasal congestion due to a cold

### Warnings

### Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

### Drug Facts (continued)

### Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
   diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

When using this product, do not use more than directed.

#### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

### Drug Facts (continued)

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- shake well before using
- adults and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

### Other information

- each 20 mL contains: sodium 17 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium

### Drug Facts (continued)

benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

PEEL CORNER FOR MORE DRUG FACTS ▲

Fred's Pharmacy Maximum Strength Mucus Relief Severe Congestion and Cough

dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55315-337
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name	<b>Basis of Strength</b>	Strength		
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL		
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL		
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL		

Inactive Ingredients	
Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging			
# Item Code	Package Description	Marketing Start Date	Marketing End Date
<b>1</b> NDC:55315-337-06	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/31/2015	06/30/2024

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M012	10/31/2015	06/30/2024	
OTC Monograph Drug	M012	10/31/2015	06/30/2024	

Revised: 11/2023 Freds Inc