

**ACETAMINOPHEN DEXTROMETHORPHAN HBR GUAIFENESIN PHENYLEPHRINE HCL- acetaminophen dextromethorphan hbr guaifenesin phenylephrine hcl tablet, coated
Strive Pharmaceuticals Inc.**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

DGH MS Mucus Relief Cold & Flu

Acetaminophen 325mg, Dextromethorphan HBr 10mg, Guaifenesin 200mg, Phenylephrine HCl 5mg

Acetaminophen 325mg

Dextromethorphan HBr 10mg

Guaifenesin 200mg

Phenylephrine HCl 5mg

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

colloidal silicon dioxide, croscarmellose sodium, di-calcium phosphate, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, stearic acid, talc, titanium dioxide

- do not take more than directed (see overdose warning)
- do not take more than 12 caplets in any 24 hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

- temporarily relieves these common cold and flu symptoms: *headache *cough *minor aches and pains *sore throat *sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose Warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adult as well as for children even if you do not notice any signs or symptoms.

Keep out of reach of children.

Overdose Warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adult as well as for children even if you do not notice any signs or symptoms.

- store between 20-25°C (68-77°F).
- avoid excessive heat, light and humidity.



Mucus Relief Cold & Flu

DG™ | health

NDC 70692-133-82

Compare to the active ingredients of Mucinex® Fast-Max® Cold & Flu ALL IN ONE!

For Ages 12 & Over

Maximum Strength

Mucus Relief Cold & Flu

**Acetaminophen/Pain Reliever/Fever Reducer
Dextromethorphan HBr/Cough Suppressant
Guaifenesin/Expectorant • Phenylephrine HCl/Nasal Decongestant**

TEMPORARILY RELIEVES

- Headache
- Sore Throat
- Nasal Congestion
- Body Pain
- Chest Congestion
- Sinus Congestion
- Fever
- Cough
- Sinus Pressure



8 Caplets

Actual Caplet Size

4 Pouches: 2 Caplets each

100%
Satisfaction
Guaranteed!
(888) 309-9030

LOT:
EXP:

A1218



3

TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR IF POUCH IS TORN OR BROKEN. KEEP OUTER PACKAGE FOR FULL INFORMATION.

DISTRIBUTED BY DOLGENCORP, LLC
100 MISSION RIDGE, GOODLETTSVILLE, TN 37072
MADE IN INDIA

CT13882 REV. 032019

*This product is not manufactured or distributed by Reckitt Benckiser, owner of the registered trademark MUCINEX® FAST-MAX® COLD & FLU ALL IN ONE

Drug Facts (continued)
Inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, d-calcium phosphate, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, stearic acid, talc, titanium dioxide
Questions or comments? 1-888-389-9030

Drug Facts (continued)
Purposes Ask a doctor before use if you have:
- high blood pressure, thyroid disease, heart disease, diabetes, liver disease, or heart disease
- cough or asthma
- trouble breathing due to an enlarged prostate gland
- asthma, chronic bronchitis, or emphysema
- persistent cough such as occurs with smoking.
Uses temporarily relieves these common cold and flu symptoms: nasal congestion, headache, cough, minor aches and pains, sore throat, sinus congestion and pressure, temporarily reduces fever, helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.
Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 12 caplets in 24 hours, which is the maximum daily amount. Do not use with other drugs containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking the product.
Other information Store between 20-25°C (68-77°F). Avoid excessive heat, light and humidity.
Directions Do not take more than directed (see **Warnings**). Do not take more than 12 caplets in any 24-hour period. Adults and children 12 years of age and over: take 2 caplets every 4 hours. Children under 12 years of age: do not use.
Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
When using this product do not use more than directed:
- If pregnant or breast-feeding, ask a health professional before use.
- If you are taking other prescription or over-the-counter drugs, ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.
- Stop use and ask a doctor if:
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache
- that lasts. These could be signs of a serious condition.
- If pregnant or breast-feeding, ask a health professional before use.
- If you get worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache
- that lasts. These could be signs of a serious condition.
When using this product do not use more than directed:
- If pregnant or breast-feeding, ask a health professional before use.
- If you are taking other prescription or over-the-counter drugs, ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.
- Stop use and ask a doctor if:
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache
- that lasts. These could be signs of a serious condition.
- If pregnant or breast-feeding, ask a health professional before use.
- If you get worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache
- that lasts. These could be signs of a serious condition.
Other information Store between 20-25°C (68-77°F). Avoid excessive heat, light and humidity.

Drug Facts
Active ingredients (in each caplet)
Acetaminophen 325 mg, Dextromethorphan HBr 10 mg, Guaifenesin 200 mg, Phenylephrine HCl 5 mg, Nasal decongestant
Contains: Cough suppressant, Expectorant, Pain reliever, Fever reducer
Contains: Cough suppressant, Expectorant, Pain reliever, Fever reducer, Nasal decongestant



ACETAMINOPHEN DEXTROMETHORPHAN HBR GUAIFENESIN PHENYLEPHRINE HCL

acetaminophen dextromethorphan hbr guaifenesin phenylephrine hcl tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70692-133
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CALCIUM PHOSPHATE (UNII: 97Z1W3NDX)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	red	Score	no score
Shape	CAPSULE	Size	20mm
Flavor		Imprint Code	S719
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	NDC:70692-133-82	8 in 1 PACKAGE; Type 0: Not a Combination Product	07/10/2019	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC monograph final	part341		07/10/2019	

Labeler - Strive Pharmaceuticals Inc. (080028013)

Revised: 7/2019

Strive Pharmaceuticals Inc.