


## **FAMOTIDINE- famotidine injection, solution** **Baxter Healthcare Corporation**

-----

### **Famotidine Injection** **in GALAXY Plastic Container** **Rx Only**

#### **DESCRIPTION**

The active ingredient in Famotidine Injection is a histamine H<sub>2</sub>-receptor antagonist. Famotidine is N'-(aminosulfonyl)-3-[[[2-[(diaminomethylene)amino]-4-thiazolyl]methyl]thio]propanimidamide. The empirical formula of famotidine is C<sub>8</sub>H<sub>15</sub>N<sub>7</sub>O<sub>2</sub>S<sub>3</sub> and its molecular weight is 337.45. Its structural formula is:



Famotidine is a white to pale yellow crystalline compound that is freely soluble in glacial acetic acid, slightly soluble in methanol, very slightly soluble in water, and practically insoluble in ethanol.

Famotidine Injection is supplied as a sterile solution, for intravenous use only, in plastic single dose GALAXY containers. Each 50 mL of the premixed, iso-osmotic intravenous injection contains 20 mg Famotidine, USP, and the following inactive ingredients: L-aspartic acid 6.8 mg, sodium chloride, USP, 450 mg, and Water for Injection. The pH ranges from 5.7 to 6.4 and may have been adjusted with additional L-aspartic acid and/or with sodium hydroxide.

The GALAXY plastic container is fabricated from a specially designed multilayer plastic. Solutions are in contact with the polyethylene layer of the container and can leach out certain chemical components of the plastic in very small amounts within the expiration period. The suitability and safety of the plastic have been confirmed in tests in animals according to the USP biological tests for plastic containers, as well as by tissue culture toxicity studies.

#### **CLINICAL PHARMACOLOGY IN ADULTS**

##### **GI Effects**

Famotidine is a competitive inhibitor of histamine H<sub>2</sub>-receptors. The primary clinically important pharmacologic activity of famotidine is inhibition of gastric secretion. Both the acid concentration and volume of gastric secretion are suppressed by famotidine, while changes in pepsin secretion are proportional to volume output.

In normal volunteers and hypersecretors, famotidine inhibited basal and nocturnal gastric secretion, as well as secretion stimulated by food and pentagastrin. After oral administration, the onset of the antisecretory effect occurred within one hour; the maximum effect was dose-dependent, occurring within one to three hours. Duration of inhibition of secretion by doses of 20 and 40 mg was 10 to 12 hours.

After intravenous administration, the maximum effect was achieved within 30 minutes. Single intravenous doses of 10 and 20 mg inhibited nocturnal secretion for a period of 10 to 12 hours. The 20 mg dose was associated with the longest duration of action in most subjects.

Single evening oral doses of 20 and 40 mg inhibited basal and nocturnal acid secretion in all subjects; mean nocturnal gastric acid secretion was inhibited by 86% and 94%, respectively, for a period of at least 10 hours. The same doses given in the morning suppressed food-stimulated acid secretion in all subjects. The mean suppression was 76% and 84%, respectively, 3 to 5 hours after administration, and 25% and 30%, respectively, 8 to 10 hours after administration. In some subjects who received the 20 mg dose, however, the antisecretory effect was dissipated within 6-8 hours. There was no cumulative effect with repeated doses. The nocturnal intragastric pH was raised by evening doses of 20 and 40 mg of famotidine to mean values of 5.0 and 6.4, respectively. When famotidine was given after breakfast, the basal daytime interdigestive pH at 3 and 8 hours after 20 or 40 mg of famotidine was raised to about 5.

Famotidine had little or no effect on fasting or postprandial serum gastrin levels. Gastric emptying and exocrine pancreatic function were not affected by famotidine.

### **Other Effects**

Systemic effects of famotidine in the central nervous system (CNS), cardiovascular, respiratory or endocrine systems were not noted in clinical pharmacology studies. Also, no antiandrogenic effects were noted (see ADVERSE REACTIONS). Serum hormone levels, including prolactin, cortisol, thyroxine (T<sub>4</sub>), and testosterone, were not altered after treatment with famotidine.

### **Pharmacokinetics**

Orally administered famotidine is incompletely absorbed and its bioavailability is 40-45%. Famotidine undergoes minimal first-pass metabolism. After oral doses, peak plasma levels occur in 1-3 hours. Plasma levels after multiple doses are similar to those after single doses. Fifteen to 20% of famotidine in plasma is protein bound. Famotidine has an elimination half-life of 2.5-3.5 hours. Famotidine is eliminated by renal (65-70%) and metabolic (30-35%) routes. Renal clearance is 250-450 mL/min, indicating some tubular excretion. Twenty-five to 30% of an oral dose and 65-70% of an intravenous dose are recovered in the urine as unchanged compound. The only metabolite identified in man is the S-oxide.

There is a close relationship between creatinine clearance values and the elimination half-life of famotidine. In patients with severe renal insufficiency, i.e., creatinine clearance less than 10 mL/min, the elimination half-life of famotidine may exceed 20 hours and adjustment of dose or dosing intervals in moderate and severe renal insufficiency may be necessary (see PRECAUTIONS, DOSAGE AND ADMINISTRATION).

In elderly patients, there are no clinically significant age-related changes in the

pharmacokinetics of famotidine. However, in elderly patients with decreased renal function, the clearance of the drug may be decreased (see PRECAUTIONS, Geriatric Use).

## Drug Interaction Studies

Human Organic Anion Transporter (OAT) 1 and 3: *In vitro* studies indicate that famotidine is a substrate for OAT1 and OAT3. Following coadministration of probenecid (1500 mg), an inhibitor of OAT1 and OAT3, with a single oral 20 mg dose of famotidine in 8 healthy subjects, the serum AUC<sub>0-10h</sub> of famotidine increased from 424 to 768 ng•hr/mL and the maximum serum concentration (C<sub>max</sub>) increased from 73 to 113 ng/mL. Renal clearance, urinary excretion rate and amount of famotidine excreted unchanged in urine were decreased. The clinical relevance of this interaction is unknown. Multidrug and Toxin Extrusion Protein 1 (MATE-1): An *in vitro* study showed that famotidine is an inhibitor of MATE-1. However, no clinically significant interaction with metformin, a substrate for MATE-1, was observed.

CYP1A2: Famotidine is a weak CYP1A2 inhibitor (see PRECAUTIONS, Drug Interactions).

## CLINICAL STUDIES

The majority of clinical study experience involved oral administration of famotidine tablets, and is provided herein for reference.

### Duodenal Ulcer

In a U.S. multicenter, double-blind study in outpatients with endoscopically confirmed duodenal ulcer, orally administered famotidine was compared to placebo. As shown in Table 1, 70% of patients treated with famotidine 40 mg h.s. were healed by week 4.

**Table 1.**

Outpatients with Endoscopically Confirmed Healed Duodenal Ulcers

	<u>Famotidine</u> 40 mg h.s. (N=89)	<u>Famotidine</u> 20 mg b.i.d. (N=84)	<u>Placebo</u> h.s. (N=97)
Week 2	*32%	*38%	17%
Week 4	*70%	*67%	31%

\* Statistically significantly different than placebo (p<0.001)

Patients not healed by week 4 were continued in the study. By week 8, 83% of patients treated with famotidine had healed versus 45% of patients treated with placebo. The incidence of ulcer healing with famotidine was significantly higher than with placebo at each time point based on proportion of endoscopically confirmed healed ulcers.

In this study, time to relief of daytime and nocturnal pain was significantly shorter for patients receiving famotidine than for patients receiving placebo; patients receiving famotidine also took less antacid than the patients receiving placebo.

### Long-Term Maintenance

Treatment of Duodenal Ulcers

Famotidine, 20 mg p.o. h.s. was compared to placebo h.s. as maintenance therapy in two double-blind, multicenter studies of patients with endoscopically confirmed healed duodenal ulcers. In the U.S. study the observed ulcer incidence within 12 months in patients treated with placebo was 2.4 times greater than in the patients treated with famotidine. The 89 patients treated with famotidine had a cumulative observed ulcer incidence of 23.4% compared to an observed ulcer incidence of 56.6% in the 89 patients receiving placebo ( $p < 0.01$ ). These results were confirmed in an international study where the cumulative observed ulcer incidence within 12 months in the 307 patients treated with famotidine was 35.7%, compared to an incidence of 75.5% in the 325 patients treated with placebo ( $p < 0.01$ ).

### Gastric Ulcer

In both a U.S. and an international multicenter, double-blind study in patients with endoscopically confirmed active benign gastric ulcer, orally administered famotidine, 40 mg h.s., was compared to placebo h.s. Antacids were permitted during the studies, but consumption was not significantly different between the famotidine and placebo groups. As shown in Table 2, the incidence of ulcer healing (dropouts counted as unhealed) with famotidine was statistically significantly better than placebo at weeks 6 and 8 in the U.S. study, and at weeks 4, 6 and 8 in the international study, based on the number of ulcers that healed, confirmed by endoscopy.

**Table 2.**

Patients with Endoscopically Confirmed Healed Gastric Ulcers

	<u>U.S. Study</u>		<u>International Study</u>	
	Famotidine 40 mg h.s. (N=74)	Placebo h.s. (N=75)	Famotidine 40 mg h.s. (N=149)	Placebo h.s. (N=145)
Week 4	45%	39%	†47%	31%
Week 6	†66%	44%	†65%	46%
Week 8	*78%	64%	†80%	54%

\*,† Statistically significantly better than placebo ( $p \leq 0.05$ ,  $p \leq 0.01$  respectively)

Time to complete relief of daytime and nighttime pain was statistically significantly shorter for patients receiving famotidine than for patients receiving placebo; however, in neither study was there a statistically significant difference in the proportion of patients whose pain was relieved by the end of the study (week 8).

### Gastroesophageal Reflux Disease (GERD)

Orally administered famotidine was compared to placebo in a U.S. study that enrolled patients with symptoms of GERD and without endoscopic evidence of erosion or ulceration of the esophagus. Famotidine 20 mg b.i.d. was statistically significantly superior to 40 mg h.s. and to placebo in providing a successful symptomatic outcome, defined as moderate or excellent improvement of symptoms (Table 3).

**Table 3.**

% Successful Symptomatic Outcome

	Famotidine <u>20 mg b.i.d.</u> (N=154)	Famotidine <u>40 mg h.s.</u> (N=149)	<u>Placebo</u> (N=73)
Week 6	82*	69	62

\*  $p \leq 0.01$  vs Placebo

By two weeks of treatment, symptomatic success was observed in a greater percentage of patients taking famotidine 20 mg b.i.d. compared to placebo ( $p \leq 0.01$ ).

Symptomatic improvement and healing of endoscopically verified erosion and ulceration were studied in two additional trials. Healing was defined as complete resolution of all erosions or ulcerations visible with endoscopy. The U.S. study comparing famotidine 40 mg p.o. b.i.d. to placebo and famotidine 20 mg p.o. b.i.d., showed a significantly greater percentage of healing for famotidine 40 mg b.i.d. at weeks 6 and 12 (Table 4).

**Table 4.**

% Endoscopic Healing - U.S. Study

	Famotidine <u>40 mg b.i.d.</u> (N=127)	Famotidine <u>20 mg b.i.d.</u> (N=125)	<u>Placebo</u> (N=66)
Week 6	48*,†	32	18
Week 12	69*,‡	54*	29

\*  $p \leq 0.01$  vs Placebo

†  $p \leq 0.05$  vs Famotidine 20 mg b.i.d.

‡  $p \leq 0.01$  vs Famotidine 20 mg b.i.d.

As compared to placebo, patients who received famotidine had faster relief of daytime and nighttime heartburn and a greater percentage of patients experienced complete relief of nighttime heartburn. These differences were statistically significant.

In the international study, when famotidine 40 mg p.o. b.i.d. was compared to ranitidine 150 mg p.o. b.i.d., a statistically significantly greater percentage of healing was observed with famotidine 40 mg b.i.d. at week 12 (Table 5). There was, however, no significant difference among treatments in symptom relief.

**Table 5.**

% Endoscopic Healing - International Study

	Famotidine <u>40 mg b.i.d.</u> (N=175)	Famotidine <u>20 mg b.i.d.</u> (N=93)	Ranitidine <u>150 mg b.i.d.</u> (N=172)
Week 6	48	52	42
Week 12	71*	68	60

\*  $p \leq 0.05$  vs Ranitidine 150 mg b.i.d.

Pathological Hypersecretory Conditions (e.g., Zollinger-Ellison Syndrome, Multiple Endocrine Adenomas)

In studies of patients with pathological hypersecretory conditions such as Zollinger-Ellison Syndrome with or without multiple endocrine adenomas, famotidine significantly inhibited gastric acid secretion and controlled associated symptoms. Orally administered doses from 20 to 160 mg q 6 h maintained basal acid secretion below 10 mEq/hr; initial doses were titrated to the individual patient need and subsequent adjustments were necessary with time in some patients. Famotidine was well tolerated at these high dose levels for prolonged periods (greater than 12 months) in eight patients, and there were no cases reported of gynecomastia, increased prolactin levels, or impotence which were considered to be due to the drug.

## CLINICAL PHARMACOLOGY IN PEDIATRIC PATIENTS

### Pharmacokinetics

Table 6 presents pharmacokinetic data from clinical trials and a published study in pediatric patients (<1 year of age; N=27) given famotidine I.V. 0.5 mg/kg and from published studies of small numbers of pediatric patients (1-15 years of age) given famotidine intravenously. Areas under the curve (AUCs) are normalized to a dose of 0.5 mg/kg I.V. for pediatric patients 1-15 years of age and compared with an extrapolated 40 mg intravenous dose in adults (extrapolation based on results obtained with a 20 mg I.V. adult dose).

**Table 6.**

Pharmacokinetic Parameters* of Intravenous Famotidine				
Age (N=number of patients)	Area Under the Curve (AUC) (ng-hr/mL)	Total Clearance (Cl) (L/hr/kg)	Volume of Distribution (V <sub>d</sub> ) (L/kg)	Elimination Half-life (T <sub>1/2</sub> ) (hours)
0-1 month <sup>†</sup> (N=10)	NA	0.13 ± 0.06	1.4 ± 0.4	10.5 ± 5.4
0-3 months <sup>‡</sup> (N=6)	2688 ± 847	0.21 ± 0.06	1.8 ± 0.3	8.1 ± 3.5
>3-12 months <sup>‡</sup> (N=11)	1160 ± 474	0.49 ± 0.17	2.3 ± 0.7	4.5 ± 1.1
1-11 yrs (N=20)	1089 ± 834	0.54 ± 0.34	2.07 ± 1.49	3.38 ± 2.60
11-15 yrs (N=6)	1140 ± 320	0.48 ± 0.14	1.5 ± 0.4	2.3 ± 0.4
Adult (N=16)	1726 <sup>§</sup>	0.39 ± 0.14	1.3 ± 0.2	2.83 ± 0.99

\* Values are presented as means ± SD unless indicated otherwise.

<sup>†</sup> Single center study.

<sup>‡</sup> Multicenter study.

<sup>§</sup> Mean value only.

Plasma clearance is reduced and elimination half-life is prolonged in pediatric patients 0-3 months of age compared to older pediatric patients. The pharmacokinetic parameters for pediatric patients, ages >3 months-15 years, are comparable to those obtained for adults.

Bioavailability studies of 8 pediatric patients (11-15 years of age) showed a mean oral bioavailability of 0.5 compared to adult values of 0.42 to 0.49. Oral doses of 0.5 mg/kg achieved AUCs of 645 ± 249 ng-hr/mL and 580 ± 60 ng-hr/mL in pediatric patients <1 year of age (N=5) and in pediatric patients 11-15 years of age, respectively, compared to 482 ± 181 ng-hr/mL in adults treated with 40 mg orally.

## Pharmacodynamics

Pharmacodynamics of famotidine were evaluated in 5 pediatric patients 2-13 years of age using the sigmoid  $E_{max}$  model. These data suggest that the relationship between serum concentration of famotidine and gastric acid suppression is similar to that observed in one study of adults (Table 7).

**Table 7.**

Pharmacodynamics of famotidine using the sigmoid  $E_{max}$  model

	<u><math>EC_{50}</math> (ng/mL)*</u>
Pediatric Patients	26 ± 13
Data from one study	
a) healthy adult subjects	26.5 ± 10.3
b) adult patients with upper GI bleeding	18.7 ± 10.8

\* Serum concentration of famotidine associated with 50% maximum gastric acid reduction. Values are presented as means ± SD.

Five published studies (Table 8) examined the effect of famotidine on gastric pH and duration of acid suppression in pediatric patients. While each study had a different design, acid suppression data over time are summarized as follows:

**Table 8.**

<u>Dosage</u>	<u>Route</u>	<u>Effect*</u>	<u>Number of Patients (age range)</u>
0.5 mg/kg, single dose	I.V.	gastric pH >4 for 19.5 hours (17.3, 21.8) <sup>†</sup>	11 (5-19 days)
0.3 mg/kg; single dose	I.V.	gastric pH >3.5 for 8.7 ± 4.7 <sup>‡</sup> hours	6 (2-7 years)
0.4-0.8 mg/kg	I.V.	gastric pH >4 for 6-9 hours	18 (2-69 months)
0.5 mg/kg, single dose	I.V.	a >2 pH unit increase above baseline in gastric pH for >8 hours	9 (2-13 years)
0.5 mg/kg b.i.d.	I.V.	gastric pH >5 for 13.5 ± 1.8 <sup>‡</sup> hours	4 (6-15 years)
0.5 mg/kg b.i.d.	oral	gastric pH >5 for 5.0 ± 1.1 <sup>‡</sup> hours	4 (11-15 years)

\* Values reported in published literature.

† Mean (95% confidence interval).

‡ Mean ± SD.

The duration of effect of famotidine I.V. 0.5 mg/kg on gastric pH and acid suppression was shown in one study to be longer in pediatric patients <1 month of age than in older pediatric patients. This longer duration of gastric acid suppression is consistent with the decreased clearance in pediatric patients <3 months of age (see Table 6).

## INDICATIONS AND USAGE

Famotidine Injection is supplied as a premixed solution in GALAXY plastic containers and is intended for intravenous use only. Famotidine Injection is indicated in some hospitalized patients with pathological hypersecretory conditions or intractable ulcers, or as an alternative to the oral dosage forms for short term use in patients who are unable to take oral medication for the following conditions:

1. *Short term treatment of active duodenal ulcer.* Most adult patients heal within 4 weeks; there is rarely reason to use famotidine at full dosage for longer than 6 to 8 weeks. Studies have not assessed the safety of famotidine in uncomplicated active duodenal ulcer for periods of more than eight weeks.
2. *Maintenance therapy for duodenal ulcer patients at reduced dosage after healing of an active ulcer.* Controlled studies in adults have not extended beyond one year.
3. *Short term treatment of active benign gastric ulcer.* Most adult patients heal within 6 weeks. Studies have not assessed the safety or efficacy of famotidine in uncomplicated active benign gastric ulcer for periods of more than 8 weeks.
4. *Short term treatment of gastroesophageal reflux disease (GERD).* Famotidine is indicated for short term treatment of patients with symptoms of GERD (see CLINICAL PHARMACOLOGY IN ADULTS, Clinical Studies).  
Famotidine is also indicated for the short term treatment of esophagitis due to GERD including erosive or ulcerative disease diagnosed by endoscopy (see CLINICAL PHARMACOLOGY IN ADULTS, Clinical Studies).
5. *Treatment of pathological hypersecretory conditions (e.g., Zollinger-Ellison Syndrome, multiple endocrine adenomas)* (see CLINICAL PHARMACOLOGY IN ADULTS, Clinical Studies).

## CONTRAINDICATIONS

Hypersensitivity to any component of this product. Cross sensitivity in this class of compounds has been observed. Therefore, Famotidine Injection should not be administered to patients with a history of hypersensitivity to other H<sub>2</sub>-receptor antagonists.

## PRECAUTIONS

### General

Symptomatic response to therapy with famotidine does not preclude the presence of gastric malignancy.

### Patients with Moderate or Severe Renal Insufficiency

CNS adverse reactions have been reported in elderly patients and patients with moderate and severe renal insufficiency. Prolonged QT intervals have also been reported in patients with moderate and severe renal insufficiency. Longer intervals between doses or lower doses may need to be used in adult and pediatric patients with moderate (creatinine clearance <50 mL/min) or severe (creatinine clearance <10 mL/min) renal insufficiency to adjust for the longer elimination half-life of famotidine (see CLINICAL PHARMACOLOGY IN ADULTS, DOSAGE AND ADMINISTRATION).



## **Drug Interactions**

### Drugs Dependent on Gastric pH for Absorption

Famotidine can reduce the absorption of other drugs, due to its effect on reducing intragastric acidity, leading to loss of efficacy of the concomitant drug. See the prescribing information for other drugs dependent on gastric pH for absorption.

### Tizanidine (CYP1A2 Substrate)

Although not studied clinically, famotidine is considered a weak CYP1A2 inhibitor. Concomitant use of tizanidine, a CYP1A2 substrate, with famotidine may lead to increased exposure of tizanidine and adverse reactions of hypotension, bradycardia or excessive drowsiness. Avoid concomitant use with famotidine. If use is clinically necessary, refer to the full prescribing information for tizanidine for dosage information.

## **Carcinogenesis, Mutagenesis, Impairment of Fertility**

In a 106 week study in rats and a 92 week study in mice given oral doses of up to 2000 mg/kg/day (approximately 2500 times the recommended human dose for active duodenal ulcer), there was no evidence of carcinogenic potential for famotidine.

Famotidine was negative in the microbial mutagen test (Ames test) using *Salmonella typhimurium* and *Escherichia coli* with or without rat liver enzyme activation at concentrations up to 10,000 mcg/plate. In *in vivo* studies in mice, with a micronucleus test and a chromosomal aberration test, no evidence of a mutagenic effect was observed.

In studies with rats given oral doses of up to 2000 mg/kg/day or intravenous doses of up to 200 mg/kg/day, fertility and reproductive performance were not affected.

## **Pregnancy**

Reproductive studies have been performed in rats and rabbits at oral doses of up to 2000 and 500 mg/kg/day, respectively, and in both species at I.V. doses of up to 200 mg/kg/day, and have revealed no significant evidence of impaired fertility or harm to the fetus due to famotidine. While no direct fetotoxic effects have been observed, sporadic abortions occurring only in mothers displaying marked decreased food intake were seen in some rabbits at oral doses of 200 mg/kg/day (250 times the usual human dose) or higher. There are, however, no adequate or well-controlled studies in pregnant women. Because animal reproductive studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

## **Nursing Mothers**

Studies performed in lactating rats have shown that famotidine is secreted into breast milk. Transient growth depression was observed in young rats suckling from mothers treated with maternotoxic doses of at least 600 times the usual human dose. Famotidine is detectable in human milk. Because of the potential for serious adverse reactions in nursing infants from famotidine, a decision should be made whether to discontinue nursing or discontinue the drug, taking into account the importance of the drug to the mother.

## **Pediatric Patients <1 year of age**

Use of famotidine in pediatric patients <1 year of age is supported by evidence from adequate and well-controlled studies of famotidine in adults, and by the following studies in pediatric patients <1 year of age.

Two pharmacokinetic studies in pediatric patients <1 year of age (N=48) demonstrated that clearance of famotidine in patients >3 months to 1 year of age is similar to that seen in older pediatric patients (1-15 years of age) and adults. In contrast, pediatric patients 0-3 months of age had famotidine clearance values that were 2- to 4-fold less than those in older pediatric patients and adults. These studies also show that the mean bioavailability in pediatric patients <1 year of age after oral dosing is similar to older pediatric patients and adults. Pharmacodynamic data in pediatric patients 0-3 months of age suggest that the duration of acid suppression is longer compared with older pediatric patients, consistent with the longer famotidine half-life in pediatric patients 0-3 months of age. (See CLINICAL PHARMACOLOGY IN PEDIATRIC PATIENTS, Pharmacokinetics and Pharmacodynamics.)

In a double-blind, randomized, treatment-withdrawal study, 35 pediatric patients <1 year of age who were diagnosed as having gastroesophageal reflux disease were treated for up to 4 weeks with famotidine oral suspension (0.5 mg/kg/dose or 1 mg/kg/dose). Although an intravenous famotidine formulation was available, no patients were treated with intravenous famotidine in this study. Also, caregivers were instructed to provide conservative treatment including thickened feedings. Enrolled patients were diagnosed primarily by history of vomiting (spitting up) and irritability (fussiness). The famotidine dosing regimen was once daily for patients < 3 months of age and twice daily for patients ≥3 months of age. After 4 weeks of treatment, patients were randomly withdrawn from the treatment and followed an additional 4 weeks for adverse events and symptomatology. Patients were evaluated for vomiting (spitting up), irritability (fussiness) and global assessments of improvement. The study patients ranged in age at entry from 1.3 to 10.5 months (mean 5.6 ± 2.9 months), 57% were female, 91% were white and 6% were black. Most patients (27/35) continued into the treatment withdrawal phase of the study. Two patients discontinued famotidine due to adverse events. Most patients improved during the initial treatment phase of the study. Results of the treatment withdrawal phase were difficult to interpret because of small numbers of patients. Of the 35 patients enrolled in the study, agitation was observed in 5 patients on famotidine that resolved when the medication was discontinued; agitation was not observed in patients on placebo (see ADVERSE REACTIONS, Pediatric Patients).

These studies suggest that a starting dose of 0.5 mg/kg/dose of famotidine oral suspension may be of benefit for the treatment of GERD for up to 4 weeks once daily in patients <3 months of age and twice daily in patients 3 months to <1 year of age; the safety and benefit of famotidine treatment beyond 4 weeks have not been established. Famotidine should be considered for the treatment of GERD only if conservative measures (e.g., thickened feedings) are used concurrently and if the potential benefit outweighs the risk.

### **Pediatric Patients 1-16 years of age**

Use of famotidine in pediatric patients 1-16 years of age is supported by evidence from adequate and well-controlled studies of famotidine in adults, and by the following studies in pediatric patients: In published studies in small numbers of pediatric patients 1-15 years of age, clearance of famotidine was similar to that seen in adults. In pediatric patients 11-15 years of age, oral doses of 0.5 mg/kg were associated with a mean area

under the curve (AUC) similar to that seen in adults treated orally with 40 mg. Similarly, in pediatric patients 1-15 years of age, intravenous doses of 0.5 mg/kg were associated with a mean AUC similar to that seen in adults treated intravenously with 40 mg. Limited published studies also suggest that the relationship between serum concentration and acid suppression is similar in pediatric patients 1-15 years of age as compared with adults. These studies suggest that the starting dose for pediatric patients 1-16 years of age is 0.25 mg/kg intravenously (injected over a period of not less than two minutes or as a 15-minute infusion) q 12 h up to 40 mg/day.

While published uncontrolled clinical studies suggest effectiveness of famotidine in the treatment of peptic ulcer, data in pediatric patients are insufficient to establish percent response with dose and duration of therapy. Therefore, treatment duration (initially based on adult duration recommendations) and dose should be individualized based on clinical response and/or gastric pH determination and endoscopy. Published uncontrolled studies in pediatric patients have demonstrated gastric acid suppression with doses up to 0.5 mg/kg intravenously q 12 h.

### **Geriatric Use**

Of the 4,966 subjects in clinical studies who were treated with famotidine, 488 subjects (9.8%) were 65 and older, and 88 subjects (1.7%) were greater than 75 years of age. No overall differences in safety or effectiveness were observed between these subjects and younger subjects. In post marketing experience, CNS adverse reactions have been reported in elderly patients with and without renal impairment receiving famotidine (see WARNINGS).

Famotidine is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, monitor renal function. Dosage adjustment in the case of moderate or severe renal impairment is necessary (see PRECAUTIONS, Patients with Moderate or Severe Renal Insufficiency and DOSAGE AND ADMINISTRATION, Dosage Adjustment for Patients with Moderate or Severe Renal Insufficiency).

### **ADVERSE REACTIONS**

The adverse reactions listed below have been reported during domestic and international clinical trials in approximately 2500 patients. In those controlled clinical trials in which famotidine tablets were compared to placebo, the incidence of adverse experiences in the group which received famotidine tablets, 40 mg at bedtime, was similar to that in the placebo group.

The following adverse reactions have been reported to occur in more than 1% of patients on therapy with famotidine in controlled clinical trials, and may be causally related to the drug: headache (4.7%), dizziness (1.3%), constipation (1.2%) and diarrhea (1.7%).

The following other adverse reactions have been reported infrequently in clinical trials or since the drug was marketed. The relationship to therapy with famotidine has been unclear in many cases. Within each category the adverse reactions are listed in order of decreasing severity:

*Body as a Whole:* fever, asthenia, fatigue

*Cardiovascular:* arrhythmia, AV block, palpitation, prolong QT interval.

*Gastrointestinal:* cholestatic jaundice, hepatitis, liver enzyme abnormalities, vomiting, nausea, abdominal discomfort, anorexia, dry mouth

*Hematologic:* rare cases of agranulocytosis, pancytopenia, leukopenia, thrombocytopenia

*Hypersensitivity:* anaphylaxis, angioedema, orbital or facial edema, urticaria, rash, conjunctival injection

*Musculoskeletal:* musculoskeletal pain including muscle cramps, arthralgia

*Nervous System/Psychiatric:* grand mal seizure; psychic disturbances, which were reversible in cases for which follow-up was obtained, including hallucinations, confusion, agitation, depression, anxiety, decreased libido; paresthesia; insomnia; somnolence. Convulsions, in patients with impaired renal function, have been reported.

*Respiratory:* bronchospasm, interstitial pneumonia

*Skin:* toxic epidermal necrolysis/Stevens Johnson syndrome, alopecia, acne, pruritus, dry skin, flushing

*Special Senses:* tinnitus, taste disorder

*Other:* rare cases of impotence and rare cases of gynecomastia have been reported; however, in controlled clinical trials, the incidences were not greater than those seen with placebo.

The adverse reactions reported for famotidine tablets may also occur with famotidine for oral suspension, famotidine injection in Galaxy plastic container, and famotidine injection. In addition, transient irritation at the injection site has been observed with famotidine injection.

## **Pediatric Patients**

In a clinical study in 35 pediatric patients <1 year of age with GERD symptoms [e.g., vomiting (spitting up), irritability (fussing)], agitation was observed in 5 patients on famotidine that resolved when the medication was discontinued.

## **OVERDOSAGE**

The adverse reactions in overdose cases are similar to the adverse reactions encountered in normal clinical experience (see ADVERSE REACTIONS). Oral doses of up to 640 mg/day have been given to adult patients with pathological hypersecretory conditions with no serious adverse effects. In the event of overdose, treatment should be symptomatic and supportive. In the event of overdose, the patient should be monitored, and supportive therapy should be employed. Due to low binding to plasma proteins, famotidine is eliminated by hemodialysis. There is a limited experience on the usefulness of hemodialysis as a treatment for famotidine overdose.

The intravenous LD<sub>50</sub> of famotidine for mice and rats ranged from 254-563 mg/kg and the minimum lethal single I.V. dose in dogs was approximately 300 mg/kg. Signs of acute intoxication in I.V. treated dogs were emesis, restlessness, pallor of mucous

membranes or redness of mouth and ears, hypotension, tachycardia and collapse. The oral LD<sub>50</sub> of famotidine in male and female rats and mice was greater than 3000 mg/kg and the minimum lethal acute oral dose in dogs exceeded 2000 mg/kg. Famotidine did not produce overt effects at high oral doses in mice, rats, cats and dogs, but induced significant anorexia and growth depression in rabbits starting with 200 mg/kg/day orally.

## **DOSAGE AND ADMINISTRATION**

In some hospitalized patients with pathological hypersecretory conditions or intractable ulcers, or in patients who are unable to take oral medication, Famotidine Injection may be administered until oral therapy can be instituted.

The recommended dosage for Famotidine Injection in adult patients is 20 mg intravenously q 12 h.

The doses and regimen for parenteral administration in patients with GERD have not been established.

### **Dosage for Pediatric Patients <1 year of age Gastroesophageal Reflux Disease (GERD)**

See PRECAUTIONS, Pediatric Patients <1 year of age.

The use of intravenous famotidine in pediatric patients <1 year of age with GERD has not been adequately studied.

### **Dosage for Pediatric Patients 1-16 years of age**

See PRECAUTIONS, Pediatric Patients 1-16 years of age.

The studies described in PRECAUTIONS, Pediatric Patients 1-16 years of age suggest that the starting dose in pediatric patients 1-16 years of age is 0.25 mg/kg intravenously (injected over a period of not less than two minutes or as a 15-minute infusion) q 12 h up to 40 mg/day.

While published uncontrolled clinical studies suggest effectiveness of famotidine in the treatment of peptic ulcer, data in pediatric patients are insufficient to establish percent response with dose and duration of therapy. Therefore, treatment duration (initially based on adult duration recommendations) and dose should be individualized based on clinical response and/or gastric pH determination and endoscopy. Published uncontrolled studies in pediatric patients 1-16 years of age have demonstrated gastric acid suppression with doses up to 0.5 mg/kg intravenously q 12 h.

### **Dosage Adjustments for Patients with Moderate or Severe Renal Insufficiency**

In adult patients with moderate (creatinine clearance <50 mL/min) or severe (creatinine clearance <10 mL/min) renal insufficiency, the elimination half-life of famotidine is increased. For patients with severe renal insufficiency, it may exceed 20 hours, reaching approximately 24 hours in anuric patients. Since CNS adverse effects have been reported in patients with moderate and severe renal insufficiency, to avoid excess accumulation of the drug in patients with moderate or severe renal insufficiency, the dose of Famotidine Injection may be reduced to half the dose, or the dosing interval may

be prolonged to 36-48 hours as indicated by the patient's clinical response.

Based on the comparison of pharmacokinetic parameters for famotidine in adults and pediatric patients, dosage adjustment in pediatric patients with moderate or severe renal insufficiency should be considered.

### **Pathological Hypersecretory Conditions (e.g., Zollinger-Ellison Syndrome, Multiple Endocrine Adenomas)**

The dosage of famotidine in patients with pathological hypersecretory conditions varies with the individual patient. The recommended adult intravenous dose is 20 mg q 12 h. Doses should be adjusted to individual patient needs and should continue as long as clinically indicated. In some patients, a higher starting dose may be required. Oral doses up to 160 mg q 6 h have been administered to some adult patients with severe Zollinger-Ellison Syndrome.

### **Famotidine Injection**

Famotidine Injection, supplied in GALAXY containers, is a 50 mL iso-osmotic solution premixed with 0.9% sodium chloride for administration as an infusion over a 15-30 minute period.

*This premixed solution is for intravenous use only using sterile equipment.*

### **DIRECTIONS FOR USE OF GALAXY CONTAINERS**

Check the container for minute leaks prior to use by squeezing the bag firmly. If leaks are found, discard solution as sterility may be impaired. Visually inspect the container. If the outlet port protector is damaged, detached, or not present, discard container as solution path sterility may be impaired. Do not add supplementary medication. Do not use unless solution is clear and seal is intact.

**CAUTION:** Do not use plastic containers in series connections. Such use could result in air embolism due to residual air being drawn from the primary container before administration of the fluid from the secondary container is complete.

### **Preparation for administration:**

1. Suspend container from eyelet support.
2. Remove protector from outlet port at bottom of container.
3. Attach administration set. Refer to complete directions accompanying set.

### **Concomitant Use of Antacids**

Antacids may be given concomitantly if needed.

### **Stability**

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration whenever solution and container permit.

Famotidine Injection, as supplied premixed in 0.9% sodium chloride in GALAXY containers, is stable through the labeled expiration date when stored under the recommended conditions. (See HOW SUPPLIED, Storage.)

## **HOW SUPPLIED**

### **FOR INTRAVENOUS USE ONLY**

Famotidine Injection containing 20 mg of famotidine per 50 mL is a clear, non-preserved, sterile solution premixed in a vehicle made iso-osmotic with sodium chloride. Famotidine Injection is supplied in a carton of 12 x 50 mL single dose GALAXY plastic containers as follows:

2G3424

50 mL container

NDC 0338-5197-41

## **Storage**

Store Famotidine Injection in GALAXY containers at room temperature (25°C/77°F). Exposure of the premixed product to excessive heat should be avoided. Brief exposure to temperatures up to 35°C (95°F) does not adversely affect the product.

## **Baxter Healthcare Corporation**

Deerfield, IL 60015 USA

Printed in USA

Baxter and Galaxy are trademarks of Baxter International Inc. or its subsidiaries.

07-19-07-644

Rev. August 2024

## **PACKAGE LABEL - PRINCIPAL DISPLAY PANEL**

**Baxter**  
**Famotidine**  
**Injection**

**20 mg**

**Rx Only**

**GALAXY**  
Single Dose  
Container

**50 mL**  
**Iso-osmotic**

**NDC 0338-5197-41**  
**Code 2G3424**  
**Sterile Nonpyrogenic**

Each 50 mL contains: 20 mg famotidine, USP, 450 mg sodium chloride, USP, and 6.8 mg L-aspartic acid. pH may have been adjusted with additional L-aspartic acid and/or sodium hydroxide. pH 5.7 to 6.4

Usual Dosage: Intravenously as directed by a physician. See package insert.

**Cautions:** Do not add supplementary medication. Must not be used in series connections.

Check for minute leaks by squeezing bag firmly. If leaks are found, discard bag as sterility may be impaired. Do not use unless solution is clear.

Recommended storage: Room temperature (25°C/77°F). Avoid excessive heat. Brief exposure up to 35°C/95°F does not adversely affect the product.

Baxter and Galaxy are registered trademarks of  
Baxter International Inc.

**Baxter Healthcare Corporation**  
Deerfield, IL 60015 USA

07-34-00-2281

**\* BAR CODE**  
**POSITION ONLY**  
303385197410

**Container Label**



**Baxter  
Famotidine  
Injection**

**20 mg**

**Rx only**

**GALAXY  
Single Dose  
Container**

**50 mL**

**Iso-osmotic**

**NDC 0338-5197-41**

**Code 2G3424**

**Sterile Nonpyrogenic**

Each 50 mL contains: 20 mg famotidine, USP, 450 mg sodium chloride, USP, and 6.8 mg L-aspartic acid. pH may have been adjusted with additional L-aspartic acid and/or sodium hydroxide. pH 5.7 to 6.4

Usual Dosage: Intravenously as directed by a physician. See package insert.

**Cautions:** Do not add supplementary medication. Must not be used in series connections.

Check for minute leaks by squeezing bag firmly. If leaks are found, discard bag as sterility may be impaired. Do not use unless solution is clear.

Recommended storage: Room temperature (25°C/77°F). Avoid excessive heat. Brief exposure up to 35°C/95°F does not adversely affect the product.

---

Baxter and Galaxy are registered trademarks of  
Baxter International Inc.

**Baxter Healthcare Corporation**

Deerfield, IL 60015 USA

07-34-00-2281

**Recommended storage: Room temperature (25°C/77°F). Avoid excessive heat. Brief exposure up to 35°C/95°F does not adversely affect the product.**

Baxter and **GALAXY** are trademarks of Baxter International Inc.

07-04-00-1261

**Recommended storage: Room temperature (25°C/77°F). Avoid excessive heat. Brief exposure up to 35°C/95°F does not adversely affect the product.**

Baxter and **GALAXY** are trademarks of Baxter International Inc.

07-04-00-1261

**Baxter**  
**Famotidine Injection**

12 x 50 mL Single Dose Containers    iso-osmotic  
Recommended storage: Room temperature (25°C/77°F)

Rx only

**20 mg**

Baxter Healthcare Corporation  
Deerfield, IL 60015 USA

NDC 0338-5197-41  
Code 2G3424

\*FOR BAR CODE POSITION ONLY

(01) 20303385197414

**Baxter**  
**Famotidine Injection**

12 x 50 mL Single Dose Containers    iso-osmotic  
Recommended storage: Room temperature (25°C/77°F)

Rx only

**20 mg**

Baxter Healthcare Corporation  
Deerfield, IL 60015 USA

NDC 0338-5197-41  
Code 2G3424

\*FOR BAR CODE POSITION ONLY

(01) 20303385197414

**GALAXY** Container

Sterile Nonpyrogenic

Each 50 mL contains: 20 mg famotidine, USP, 450 mg sodium chloride, USP, and 6.8 mg L-aspartic acid. pH may have been adjusted with additional L-aspartic acid and/or sodium hydroxide. pH 5.7 to 6.4  
Usual Dosage: Intravenously as directed by physician. See package insert.

**Cautions:** Do not add supplementary medication. Must not be used in series connections. Check for minute leaks by squeezing bag firmly. If leaks are found, discard bag as sterility may be impaired. Do not use unless solution is clear.

**GALAXY** Container

Sterile Nonpyrogenic

Each 50 mL contains: 20 mg famotidine, USP, 450 mg sodium chloride, USP, and 6.8 mg L-aspartic acid. pH may have been adjusted with additional L-aspartic acid and/or sodium hydroxide. pH 5.7 to 6.4  
Usual Dosage: Intravenously as directed by physician. See package insert.

**Cautions:** Do not add supplementary medication. Must not be used in series connections. Check for minute leaks by squeezing bag firmly. If leaks are found, discard bag as sterility may be impaired. Do not use unless solution is clear.

**Carton Label**

**Recommended storage: Room temperature (25°C/77°F). Avoid excessive heat.**

**Brief exposure up to 35°C/95°F does not adversely affect the product.**

Baxter and **GALAXY** are trademarks of Baxter International Inc. 07-04-00-1261  
07-04-00-1261

**Recommended storage: Room temperature (25°C/77°F). Avoid excessive heat.**

**Brief exposure up to 35°C/95°F does not adversely affect the product.**

Baxter and **GALAXY** are trademarks of Baxter International Inc. 07-04-00-1261  
07-04-00-1261

**Baxter NDC 0338-5197-41**

**Famotidine Injection Rx only 20 mg Code 2G3424**

**12 x 50 mL Single Dose Containers Iso-osmotic**

**Recommended storage: Room temperature (25°C/77°F)**

**Baxter Healthcare Corporation**

Deerfield, IL 60015 USA

\*FOR BAR CODE POSITION ONLY

(01) 20303385197414

**Baxter NDC 0338-5197-41**

**Famotidine Injection Rx only 20 mg Code 2G3424**

**12 x 50 mL Single Dose Containers Iso-osmotic**

**Recommended storage: Room temperature (25°C/77°F)**

**Baxter Healthcare Corporation**

Deerfield, IL 60015 USA

\*FOR BAR CODE POSITION ONLY

(01) 20303385197414

**GALAXY** Container Sterile Nonpyrogenic

Each 50 mL contains: 20 mg famotidine, USP, 450 mg sodium chloride, USP, and 6.8 mg L-aspartic acid. pH may have been adjusted with additional L-aspartic acid and/or sodium hydroxide. pH 5.7 to 6.4

Usual Dosage: Intravenously as directed by physician. See package insert.

**Cautions:** Do not add supplementary medication. Must not be used in series connections. Check for minute leaks by squeezing bag firmly. If leaks are found, discard bag as sterility may be impaired. Do not use unless solution is clear.

**GALAXY** Container Sterile Nonpyrogenic

Each 50 mL contains: 20 mg famotidine, USP, 450 mg sodium chloride, USP, and 6.8 mg L-aspartic acid. pH may have been adjusted with additional L-aspartic acid and/or sodium hydroxide. pH 5.7 to 6.4

Usual Dosage: Intravenously as directed by physician. See package insert.

**Cautions:** Do not add supplementary medication. Must not be used in series connections. Check for minute leaks by squeezing bag firmly. If leaks are found, discard bag as sterility may be impaired. Do not use unless solution is clear.

---

# FAMOTIDINE

famotidine injection, solution

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0338-5197
<b>Route of Administration</b>	INTRAVENOUS		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>FAMOTIDINE</b> (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	20 mg in 50 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	450 mg in 50 mL
<b>ASPARTIC ACID</b> (UNII: 30KYC7MIAI)	6.8 mg in 50 mL
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	
<b>WATER</b> (UNII: 059QF0KO0R)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-5197-41	12 in 1 CARTON	05/10/2001	
1		50 mL in 1 BAG; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075591	05/10/2001	

**Labeler** - Baxter Healthcare Corporation (005083209)

## Establishment

Name	Address	ID/FEI	Business Operations
Baxter Healthcare Corporation		194684502	MANUFACTURE(0338-5197) , LABEL(0338-5197) , ANALYSIS(0338-5197) , PACK(0338-5197) , STERILIZE(0338-5197)

Revised: 8/2024

Baxter Healthcare Corporation