

## **QC PINK BISMUTH STOMACH RELIEF- bismuth subsalicylate tablet, chewable CHAIN DRUG MARKETING ASSOCIATION**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **QC Pink Bismuth Subsalicylate 262mg 30 Chewable Tablets**

#### **Active ingredient (in each tablet)**

Bismuth subsalicylate 262 mg

#### **Purpose**

Upset stomach reliever and antidiarrheal

#### **Uses**

relieves:

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn
- indigestion
- nausea
- gas
- belching
- fullness

#### **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

#### **Allergy alert: Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

#### **Do not use if you have**

- an ulcer
- a bleeding problem
- bloody or black stool

## Ask a doctor before use if you have

- fever
- mucus in the stool

## Ask a doctor or pharmacist before use if you are taking any drug for

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

## When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur.

## Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

**If pregnant or breast-feeding,** ask a health professional before use.

## Keep Out of Reach of Children

In case of overdose, get medical help or contact a Poison Control Center immediately at 1-800-222-1222

## Directions

- chew completely or dissolve tablets in mouth.
- **adults and children 12 years and older:** 2 tablets (1 dose) every ½ hour to 1 hour as needed for diarrhea
- 2 tablets (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (16 tablets) in 24 hours
- use until diarrhea stops but not more than 2 days
- **children under 12 years:** ask a doctor
- drink plenty clear fluids to help prevent dehydration caused by diarrhea.

## Other information

- each tablet contains: **salicylate 102 mg**
- store at room temperature.
- avoid excessive heat
- keep this carton for future reference on full labeling

**TAMPER EVIDENT: Do not use if printed inner wrap is broken or missing.**

## **Inactive ingredients**

D&C Red No. 27 lake, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, sorbitol, sucralose.

## **Questions comments?**

**1-800-935-2362**

## **Principal Display**

QUALITY CHOICE®

**NDC# 63868-705-30**

**\*Compare to the active ingredient in Pepto-Bismol® Chewables\***

## **Pink Bismuth**

Stomach Relief

Bismuth Subsalicylate

## **Upset Stomach Reliever/Antidiarrheal**

5 Symptom Digestive Relief

Heartburn

Indigestion

Nausea

Upset Stomach

Diarrhea

Original Flavor

**30 Chewable Tablets**

**QC SATISFACTION GUARANTEED**

## **DISTRIBUTED BY:**

C.D.M.A., Inc.

43157 W. Nine Mile

Novi, MI 48376-0995

[www.qualitychoice.com](http://www.qualitychoice.com)

Questions: 248-449-9300

\*This product is not manufactured or distributed by PROCTER & GAMBLE INC., the distributor of Pepto-Bismol®



## QC PINK BISMUTH STOMACH RELIEF

bismuth subsalicylate tablet, chewable

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63868-705
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	262 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6130)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>D&amp;C RED NO. 27</b> (UNII: 2LRS185U6K)	

**SORBITOL** (UNII: 506T60A25R)  
**SUCRALOSE** (UNII: 96K6UQ3ZD4)

### Product Characteristics

<b>Color</b>	PINK	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	RP124
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-705-30	5 in 1 CARTON	02/24/2020	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part335	02/24/2020	

**Labeler** - CHAIN DRUG MARKETING ASSOCIATION (011920774)

Revised: 1/2022

CHAIN DRUG MARKETING ASSOCIATION