

**MULTI-SYMPTOM MAPAP COLD FORMULA- acetaminophen,
dextromethorphan hbr, phenylephrine hcl tablet, film coated
Major Pharmaceuticals**

Major 44-470C-CMS

Active ingredients (in each caplet)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - minor aches and pains
 - nasal congestion
 - headache
 - cough
 - sinus congestion and pressure
 - sore throat
- helps clear nasal passages
- promotes nasal and sinus drainage
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- thyroid disease
- high blood pressure
- liver disease
- heart disease
- diabetes
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- adults and children 12 years and over
 - take 2 caplets every 4 hours
 - swallow whole - do not crush, chew, or dissolve
 - do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, croscarmellose sodium, crospovidone, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, stearic acid, sucralose, talc, titanium dioxide

Questions or comments?

1-800-426-9391

Principal Display Panel

MAJOR®

NDC 0904-5786-24

MULTI-SYMP TOM

Mapap®

COLD FORMULA

CAPLETS

ACETAMINOPHEN

DEXTROMETHORPHAN HBr

PHENYLEPHRINE HCl

Pain Reliever / Fever Reducer

Cough Suppressant

Nasal Decongestant

For Relief of: Fever, Headache,
Sore Throat, Nasal Congestion, Cough

DAYTIME

ASPIRIN FREE

24 CAPLETS

with Cool
Blast Flavor

Actual Size

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

50844 REV0722J47008

Distributed by:
MAJOR® PHARMACEUTICALS
Indianapolis, IN 46268
(800) 616-2471
www.majorpharmaceuticals.com

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

6 0940578624 3
No print/No varnish
Rev 03/24 M-17 Re-order No. 006363

NDC 0904-5786-24

MAJOR®

Multipap®
COLD FORMULA
CAPLETS

ACETAMINOPHEN
DEXTROMETHORPHAN HBr
PHENYLEPHRINE HCl

Pain Reliever / Fever Reducer
Cough Suppressant
Nasal Decongestant

For Relief of: Fever, Headache,
Sore Throat, Nasal Congestion, Cough

DAYTIME
ASPIRIN FREE

24 CAPLETS with Cool Blast Flavor
Actual Size

50844 REV0722J47008
Distributed by:
MAJOR® PHARMACEUTICALS
Indianapolis, IN 46268
(800) 616-2471
www.majorpharmaceuticals.com

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Drug Facts (continued)
Questions or comments? 1-800-426-9391
TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Drug Facts (continued)
Inactive ingredients: corn starch, croscarmellose sodium, croscopolone, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, stearic acid, sucralose, talc, titanium dioxide

Drug Facts (continued)
Uses
temporarily relieves these common cold and flu symptoms:
minor aches and pains
headache
nasal congestion
cough
sore throat
sinus congestion and pressure
temporarily reduces fever
helps clear nasal passages
promotes nasal and sinus drainage

Drug Facts (continued)
Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen.
3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: blisters, rash, skin redness.
If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
If you have ever had an allergic reaction to this product or any of its ingredients
Ask a doctor before use if you have liver disease, diabetes, heart disease, thyroid disease, high blood pressure, difficulty in urination due to enlargement of the prostate gland, persistent or chronic cough such as occurs with smoking, asthma, or emphysema, or cough that occurs with too much phlegm (mucus)

Drug Facts (continued)
Directions
do not take more than directed
adults and children 12 years and over
take 2 caplets every 4 hours
swallow whole - do not crush, chew, or dissolve
do not take more than 10 caplets in 24 hours
children under 12 years: ask a doctor

Other information
TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
see end flap for expiration date and lot number

Drug Facts (continued)
Warnings
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.
When using this product do not exceed recommended dosage.
Stop use and ask a doctor if nervousness, dizziness, or sleeplessness occur
pain, nasal congestion, or cough gets worse or lasts more than 7 days
fever gets worse or lasts more than 3 days
redness or swelling is present
new symptoms occur
cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)
Active ingredients (in each caplet)
Acetaminophen 325 mg..... Pain reliever/fever reducer
Dextromethorphan HBr 10 mg..... Cough suppressant
Phenylephrine HCl 5 mg..... Nasal decongestant

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Major 44 -470C

MULTI-SYMPTOM MAPAP COLD FORMULA

acetaminophen, dextromethorphan hbr, phenylephrine hcl tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0904-5786
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL	Size	17mm
Flavor	MENTHOL	Imprint Code	44;470
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0904-5786-24	2 in 1 CARTON	07/15/2005	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/15/2005	

Labeler - Major Pharmaceuticals (191427277)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(0904-5786) , pack(0904-5786)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(0904-5786)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(0904-5786)

Revised: 5/2024

Major Pharmaceuticals