

MUPIROCIN- mupirocin ointment Proficient Rx LP


MUPIROCIN OINTMENT USP, 2%

For Dermatologic Use

Rx Only

DESCRIPTION

Each gram of Mupirocin Ointment USP, 2% contains 20 mg mupirocin in a bland water miscible ointment base (polyethylene glycol ointment, NF) consisting of polyethylene glycol 400 and polyethylene glycol 3350. Mupirocin is a naturally occurring antibiotic. The chemical name is (*E*)-(2*S*,3*R*,4*R*,5*S*)-5-[(2*S*,3*S*,4*S*,5*S*)-2,3-Epoxy-5-hydroxy-4-methylhexyl]tetrahydro-3,4-dihydroxy-β-methyl-2*H*-pyran-2-crotonic acid, ester with 9-hydroxynonanoic acid. The molecular formula of mupirocin is C₂₆H₄₄O₉ and the molecular weight is 500.62. The chemical structure is:



CLINICAL PHARMACOLOGY

Application of ¹⁴C-labeled mupirocin ointment to the lower arm of normal male subjects followed by occlusion for 24 hours showed no measurable systemic absorption (<1.1 nanogram mupirocin per milliliter of whole blood). Measurable radioactivity was present in the stratum corneum of these subjects 72 hours after application.

Following intravenous or oral administration, mupirocin is rapidly metabolized. The principal metabolite, monic acid, is eliminated by renal excretion, and demonstrates no antibacterial activity. In a study conducted in 7 healthy adult male subjects, the elimination half-life after intravenous administration of mupirocin was 20 to 40 minutes for mupirocin and 30 to 80 minutes for monic acid. The pharmacokinetics of mupirocin has not been studied in individuals with renal insufficiency.

Microbiology -

Mupirocin is an antibacterial agent produced by fermentation using the organism *Pseudomonas fluorescens*. It is active against a wide range of gram-positive bacteria including methicillin-resistant *Staphylococcus aureus* (MRSA). It is also active against certain gram-negative bacteria. Mupirocin inhibits bacterial protein synthesis by reversibly and specifically binding to bacterial isoleucyl transfer-RNA synthetase. Due to

this unique mode of action, mupirocin demonstrates no *in vitro* cross-resistance with other classes of antimicrobial agents.

Resistance occurs rarely. However, when mupirocin resistance does occur, it appears to result from the production of a modified isoleucyl-tRNA synthetase. High-level plasmid-mediated resistance (MIC > 1024 mcg/mL) has been reported in some strains of *S. aureus* and coagulase-negative staphylococci.

Mupirocin is bactericidal at concentrations achieved by topical administration. However, the minimum bactericidal concentration (MBC) against relevant pathogens is generally 8-fold to 30-fold higher than the minimum inhibitory concentration (MIC). In addition, mupirocin is highly protein bound (>97%), and the effect of wound secretions on the MICs of mupirocin has not been determined.

Mupirocin has been shown to be active against most strains of *S. aureus* and *Streptococcus pyogenes*, both *in vitro* and in clinical studies (see **INDICATIONS AND USAGE**). The following *in vitro* data are available, BUT THEIR CLINICAL SIGNIFICANCE IS UNKNOWN. Mupirocin is active against most strains of *Staphylococcus epidermidis* and *Staphylococcus saprophyticus*.

INDICATIONS AND USAGE

Mupirocin Ointment USP, 2% is indicated for the topical treatment of impetigo due to: *S. aureus* and *S. pyogenes*.

CONTRAINDICATIONS

This drug is contraindicated in individuals with a history of sensitivity reactions to any of its components.

WARNINGS

Mupirocin Ointment USP, 2% is not for ophthalmic use.

PRECAUTIONS

If a reaction suggesting sensitivity or chemical irritation should occur with the use of Mupirocin Ointment USP, 2%, treatment should be discontinued and appropriate alternative therapy for the infection instituted.

As with other antibacterial products, prolonged use may result in overgrowth of nonsusceptible organisms, including fungi.

Mupirocin Ointment USP, 2% is not formulated for use on mucosal surfaces. Intranasal use has been associated with isolated reports of stinging and drying. A paraffin-based formulation - *Bactroban Nasal® (mupirocin calcium ointment) - is available for intranasal use.

Polyethylene glycol can be absorbed from open wounds and damaged skin and is excreted by the kidneys. In common with other polyethylene glycol-based ointments, Mupirocin Ointment USP, 2% should not be used in conditions where absorption of large

quantities of polyethylene glycol is possible, especially if there is evidence of moderate or severe renal impairment.

Information for Patients -

Use this medication only as directed by your healthcare provider. It is for external use only. Avoid contact with the eyes. The medication should be stopped and your healthcare practitioner contacted if irritation, severe itching, or rash occurs. If impetigo has not improved in 3 to 5 days, contact your healthcare practitioner.

Drug Interactions -

The effect of the concurrent application of Mupirocin Ointment USP, 2% and other drug products has not been studied.

Carcinogenesis, Mutagenesis, Impairment of Fertility -

Long-term studies in animals to evaluate carcinogenic potential of mupirocin have not been conducted.

Results of the following studies performed with mupirocin calcium or mupirocin sodium *in vitro* and *in vivo* did not indicate a potential for genotoxicity: Rat primary hepatocyte unscheduled DNA synthesis, sediment analysis for DNA strand breaks, *Salmonella* reversion test (Ames), *Escherichia coli* mutation assay, metaphase analysis of human lymphocytes, mouse lymphoma assay, and bone marrow micronuclei assay in mice.

Reproduction studies were performed in male and female rats with mupirocin administered subcutaneously at doses up to 14 times a human topical dose (approximately 60 mg mupirocin per day) on a mg/m² basis and revealed no evidence of impaired fertility and reproductive performance from mupirocin.

Pregnancy:

Teratogenic Effects -

Reproduction studies have been performed in rats and rabbits with mupirocin administered subcutaneously at doses up to 22 and 43 times, respectively, the human topical dose (approximately 60 mg mupirocin per day) on a mg/m² basis and revealed no evidence of harm to the fetus due to mupirocin. There are however, no adequate and well-controlled studies in pregnant women. Because animal studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Nursing Mothers -

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when Mupirocin Ointment USP, 2% is administered to a nursing woman.

Pediatric Use -

The safety and effectiveness of Mupirocin Ointment USP, 2% have been established in the age range of 2 months to 16 years. Use of mupirocin ointment USP, 2% in these age groups is supported by evidence from adequate and well-controlled studies of mupirocin

ointment USP, 2% in impetigo in pediatric patients studied as part of the pivotal clinical trials (see **CLINICAL STUDIES**).

ADVERSE REACTIONS

The following local adverse reactions have been reported in connection with the use of mupirocin ointment USP, 2%: Burning, stinging, or pain in 1.5% of patients; itching in 1% of patients; rash, nausea, erythema, dry skin, tenderness, swelling, contact dermatitis, and increased exudate in less than 1% of patients. Systemic reactions to mupirocin ointment USP, 2% have occurred rarely.

DOSAGE AND ADMINISTRATION

A small amount of Mupirocin Ointment USP, 2% should be applied to the affected area 3 times daily. The area treated may be covered with a gauze dressing if desired. Patients not showing a clinical response within 3 to 5 days should be re-evaluated.

CLINICAL STUDIES

The efficacy of topical mupirocin ointment USP, 2% in impetigo was tested in two studies. In the first, patients with impetigo were randomized to receive either mupirocin ointment USP, 2% or vehicle placebo three times daily for 8 to 12 days. Clinical efficacy rates at end of therapy in the evaluable populations (adults and pediatric patients included) were 71% for mupirocin ointment USP, 2% (n=49) and 35% for vehicle placebo (n=51). Pathogen eradication rates in the evaluable populations were 94% for mupirocin ointment USP, 2% and 62% for vehicle placebo. There were no side effects reported in the group receiving mupirocin ointment USP, 2%.

In the second study, patients with impetigo were randomized to receive either mupirocin ointment USP, 2% three times daily or 30 to 40 mg/kg oral erythromycin ethylsuccinate per day (this was an unblinded study) for 8 days. There was a follow-up visit 1 week after treatment ended. Clinical efficacy rates at the follow-up visit in the evaluable populations (adults and pediatric patients included) were 93% for mupirocin ointment USP, 2% (n=29) and 78.5% for erythromycin (n=28). Pathogen eradication rates in the evaluable patient populations were 100% for both test groups. There were no side effects reported in the group receiving mupirocin ointment.

Pediatrics -

There were 91 pediatric patients aged 2 months to 15 years in the first study described above. Clinical efficacy rates at end of therapy in the evaluable populations were 78% for mupirocin ointment USP, 2% (n=42) and 36% for vehicle placebo (n=49). In the second study described above, all patients were pediatric except two adults in the group receiving mupirocin ointment USP, 2%. The age range of the pediatric patients was 7 months to 13 years. The clinical efficacy rate for mupirocin ointment USP, 2% (n=27) was 96%, and for erythromycin it was unchanged (78.5%).

HOW SUPPLIED

Mupirocin Ointment USP, 2% is available as follows:
22 gram tube (NDC 63187-261-22)

Store at 20-25°C (68-77°F) [see USP Controlled Room Temperature].

*Bactroban Nasal® is a registered trademark of GlaxoSmithKline.

MADE IN ISRAEL

MANUFACTURED BY PERRIGO, YERUHAM 80500, ISRAEL

DISTRIBUTED BY

PERRIGO®

ALLEGAN, MI 49010

Rev. 01/09

: 1N200 RC J3

OR

Manufactured By Perrigo

Bronx, NY 10457

Distributed By Perrigo

Allegan, MI 49010 • www.perrigo.com

Rev. 11/12

: 5R200 RC J2

Relabeled by:

Proficient Rx LP

Thousand Oaks, CA 91320

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL



Scan Here



NDC 63187-261-22

Relabeled By: Proficient Rx LP
Thousand Oaks, CA 91320

RX Only

Mupirocin 2%
22gm Ointment
Lot #:00000
NDC 63187-261-22
SN# MASTER
Exp:00/00/00

Mupirocin 2%
22gm Ointment
Lot #:00000
NDC 63187-261-22
SN# MASTER
Exp:00/00/00

Mupirocin 2%
22gm Ointment
Lot #:00000
NDC 63187-261-22
SN# MASTER
Exp:00/00/00



GTIN: 00363187261221
SN# MASTER
Exp. 00/00/00
Lot #:00000



Mupirocin 2%

22gm Ointment

Each gram contains: 20 mg mupirocin in a polyethylene glycol ointment, NF base.

See package insert

Product ID: RM026122

Mfr. By: Perrigo Yeruham, Israel Made in Israel

Store at 20°-25°C (68°-77°F)

Keep medication out of the reach of children

MUPIROCIN

mupirocin ointment

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:63187-261(NDC:45802-112)
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MUPIROCIN (UNII: D0GX8630A5) (MUPIROCIN - UNII:D0GX8630A5)	MUPIROCIN	20 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63187-261-22	1 in 1 CARTON	10/01/2014	
1		22 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA065123	10/30/2009	

Labeler - Proficient Rx LP (079196022)

Establishment

Name	Address	ID/FEI	Business Operations
Proficient Rx LP		079196022	RELABEL(63187-261)

Revised: 4/2022

Proficient Rx LP