

BUTORPHANOL TARTRATE- butorphanol tartrate injection, solution
Hospira, Inc.

Butorphanol Tartrate Injection, USP CIV

WARNING: ADDICTION, ABUSE, AND MISUSE; LIFE-THREATENING RESPIRATORY DEPRESSION; NEONATAL OPIOID WITHDRAWAL SYNDROME; and RISKS FROM CONCOMITANT USE WITH BENZODIAZEPINES OR OTHER CNS DEPRESSANTS

Addiction, Abuse, and Misuse

Butorphanol tartrate injection exposes patients and other users to the risks of opioid addiction, abuse, and misuse, which can lead to overdose and death. Assess each patient's risk prior to prescribing butorphanol tartrate injection, and monitor all patients regularly for the development of these behaviors or conditions [see WARNINGS].

Life-Threatening Respiratory Depression

Serious, life-threatening, or fatal respiratory depression may occur with use of butorphanol tartrate injection. Monitor for respiratory depression, especially during initiation of butorphanol tartrate injection or following a dose increase [see WARNINGS].

Neonatal Opioid Withdrawal Syndrome

Prolonged use of butorphanol tartrate injection during pregnancy can result in neonatal opioid withdrawal syndrome, which may be life-threatening if not recognized and treated, and requires management according to protocols developed by neonatology experts. If opioid use is required for a prolonged period in a pregnant woman, advise the patient of the risk of neonatal opioid withdrawal syndrome and ensure that appropriate treatment will be available [see WARNINGS].

Risks From Concomitant Use With Benzodiazepines Or Other CNS Depressants


Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death [see WARNINGS, PRECAUTIONS; Drug Interactions].

- Reserve concomitant prescribing of butorphanol tartrate and benzodiazepines or other CNS depressants for use in patients for whom alternative treatment options are inadequate.
- Limit dosages and durations to the minimum required.
- Follow patients for signs and symptoms of respiratory depression and sedation.

DESCRIPTION

Butorphanol tartrate is a synthetically derived opioid agonist-antagonist analgesic of the phenanthrene series. The chemical name is (-)-17-(cyclobutylmethyl) morphinan-3, 14-

diol D-(-)- tartrate (1:1) (salt). The molecular formula is $C_{21}H_{29}NO_2 \cdot C_4H_6O_6$, which corresponds to a molecular weight of 477.56 and the following structural formula:



Butorphanol tartrate is a white crystalline substance. The dose is expressed as the tartrate salt. One milligram of the salt is equivalent to 0.68 mg of the free base. The n-octanol/aqueous buffer partition coefficient of butorphanol is 180:1 at pH 7.5.

Butorphanol tartrate injection is a sterile, nonpyrogenic parenteral aqueous solution of butorphanol tartrate for intravenous or intramuscular administration.

Each milliliter (mL) contains butorphanol tartrate 1 or 2 mg; sodium citrate, dihydrate, 6.4 mg; citric acid hydrous 3.3 mg; sodium chloride 6.4 mg. The pH is 4.5 (3.0 to 5.5).

CLINICAL PHARMACOLOGY

Mechanism of Action

Butorphanol is a partial opioid agonist at the mu opioid receptor and a full agonist at the kappa opioid receptor. The principal therapeutic action of butorphanol is analgesia. Clinically, dosage is titrated to provide adequate analgesia and may be limited by adverse reactions, including respiratory and CNS depression.

The precise mechanism of the analgesic action is unknown. However, specific CNS opioid receptors for endogenous compounds with opioid-like activity have been identified throughout the brain and spinal cord and are thought to play a role in the analgesic effects of this drug.

Pharmacodynamics

The analgesic effect of butorphanol is influenced by the route of administration. Onset of analgesia is within a few minutes for intravenous administration and within 15 minutes for intramuscular injection.

Peak analgesic activity occurs within 30 to 60 minutes following intravenous and intramuscular administration.

The duration of analgesia varies depending on the pain model as well as the route of administration, but is generally 3 to 4 hours with IM and IV doses as defined by the time

50% of patients required remedication. In postoperative studies, the duration of analgesia with IV or IM butorphanol was similar to morphine, meperidine and pentazocine when administered in the same fashion at equipotent doses (see **Clinical Trials**).

Effects on the Central Nervous System

Butorphanol produces respiratory depression by direct action on brain stem respiratory centers. The respiratory depression involves a reduction in the responsiveness of the brain stem respiratory centers to both increases in carbon dioxide tension and electrical stimulation.

In human studies involving individuals without significant respiratory dysfunction, 2 mg of butorphanol IV and 10 mg of morphine sulfate IV depressed respiration to a comparable degree. At higher doses, the magnitude of respiratory depression with butorphanol is not appreciably increased; however, the duration of respiratory depression is longer. Respiratory depression noted after administration of butorphanol to humans by any route is reversed by treatment with naloxone, a specific opioid antagonist (see **OVERDOSAGE**).

Butorphanol, like other mixed agonist-antagonists with a high affinity for the kappa receptor, may produce unpleasant psychotomimetic effects in some individuals.

Nausea and/or vomiting may be produced by doses of 1 mg or more administered by any route.

In human studies of butorphanol (see **CLINICAL PHARMACOLOGY; Clinical Trials**), sedation is commonly noted at doses of 0.5 mg or more. Narcosis is produced by 10 to 12 mg doses of butorphanol administered over 10 to 15 minutes intravenously.

Butorphanol causes miosis, even in total darkness. Pinpoint pupils are a sign of opioid overdose but are not pathognomonic (e.g., pontine lesions of hemorrhagic or ischemic origins may produce similar findings). Marked mydriasis rather than miosis may be seen due to hypoxia in overdose situations.

Effects on the Gastrointestinal Tract and Other Smooth Muscle

Butorphanol causes a reduction in motility associated with an increase in smooth muscle tone in the antrum of the stomach and duodenum. Digestion of food in the small intestine is delayed and propulsive contractions are decreased. Propulsive peristaltic waves in the colon are decreased, while tone may be increased to the point of spasm, resulting in constipation. Other opioid-induced effects may include a reduction in biliary and pancreatic secretions, spasm of sphincter of Oddi, and transient elevations in serum amylase.

Effects on the Cardiovascular System

Hemodynamic changes noted during cardiac catheterization in patients receiving single 0.025 mg/kg intravenous doses of butorphanol have included increases in pulmonary artery pressure, wedge pressure and vascular resistance, increases in left ventricular end diastolic pressure and in systemic arterial pressure.

Effects on the Endocrine System

Opioids inhibit the secretion of adrenocorticotropic hormone (ACTH), cortisol, and luteinizing hormone (LH) in humans (see **ADVERSE REACTIONS**). They also stimulate prolactin, growth hormone (GH) secretion, and pancreatic secretion of insulin and glucagon.

Chronic use of opioids may influence the hypothalamic-pituitary-gonadal axis, leading to androgen deficiency that may manifest as low libido, impotence, erectile dysfunction, amenorrhea, or infertility. The causal role of opioids in the clinical syndrome of hypogonadism is unknown because the various medical, physical, lifestyle, and psychological stressors that may influence gonadal hormone levels have not been adequately controlled for in studies conducted to date (see **ADVERSE REACTIONS**).

Effects on the Immune System

Opioids have been shown to have a variety of effects on components of the immune system. The clinical significance of these findings is unknown. Overall, the effects of opioids appear to be modestly immunosuppressive.

Concentration-Efficacy Relationships

The minimum effective analgesic concentration will vary widely among patients, especially among patients who have been previously treated with potent agonist opioids. The minimum effective analgesic concentration of butorphanol for any individual patient may increase over time due to an increase in pain, the development of a new pain syndrome, and/or the development of analgesic tolerance (see **DOSAGE AND ADMINISTRATION**).

Concentration-Adverse Reaction Relationships


There is a relationship between increasing butorphanol plasma concentration and increasing frequency of dose-related opioid adverse reactions such as nausea, vomiting, CNS effects, and respiratory depression. In opioid-tolerant patients, the situation may be altered by the development of tolerance to opioid-related adverse reactions (see **DOSAGE AND ADMINISTRATION**).

Pharmacokinetics

Butorphanol tartrate injection is rapidly absorbed after IM injection and peak plasma levels are reached in 20 to 40 minutes.

Following its initial absorption/distribution phase, the single dose pharmacokinetics of butorphanol by the intravenous and intramuscular routes of administration are similar (see Figure 1).

Figure 1—Butorphanol Plasma Levels After IV and IM Administration of 2 mg Dose



Serum protein binding is independent of concentration over the range achieved in clinical practice (up to 7 ng/mL) with a bound fraction of approximately 80%.

The volume of distribution of butorphanol varies from 305 to 901 liters and total body clearance from 52 to 154 liters/hr (see Table 1).

Table 1—Mean Pharmacokinetic Parameters of Intravenous Butorphanol in Young and Elderly Subjects*

Parameters	Young	Elderly
AUC (inf) [†] (hr • ng/mL)	7.24 (1.57) [‡] (4.40–9.77) [§]	8.71 (2.02) (4.76–13.03)
Half-life (hr)	4.56 (1.67) (2.06–8.70)	5.61 (1.36) (3.25–8.79)
Volume of Distribution [¶] (L)	487 (155) (305–901)	552 (124) (305–737)
Total body Clearance (L/hr)	99 (23) (70–154)	82 (21) (52–143)

* Young subjects (n=24) are from 20 to 40 years old and elderly (n=24) are greater than 65 years of age.

[†] Area under plasma concentration-time curve after a 1 mg dose.

[‡] Mean (1S.D.).

[§] (range of observed values).

[¶] Derived from IV data.

The drug is transported across the blood-brain and placental barriers and into human milk (see **PRECAUTIONS: Labor or Delivery and Nursing Mothers**).

Butorphanol is extensively metabolized in the liver. Metabolism is qualitatively and quantitatively similar following intravenous or intramuscular administration. Oral bioavailability is only 5 to 17% because of extensive first pass metabolism of butorphanol.

The major metabolite of butorphanol is hydroxybutorphanol, while norbutorphanol is produced in small amounts. Both have been detected in plasma following administration

of butorphanol, with norbutorphanol present at trace levels at most time points. The elimination half-life of hydroxybutorphanol is about 18 hours and, as a consequence, considerable accumulation (~5-fold) occurs when butorphanol is dosed to steady state.

Elimination occurs by urine and fecal excretion. When ³H labeled butorphanol is administered to normal subjects, most (70 to 80%) of the dose is recovered in the urine, while approximately 15% is recovered in the feces.

About 5% of the dose is recovered in the urine as butorphanol. Forty-nine percent is eliminated in the urine as hydroxybutorphanol. Less than 5% is excreted in the urine as norbutorphanol.

Butorphanol pharmacokinetics in the elderly differ from younger patients (see Table 1).

In renally impaired patients with creatinine clearances <30 mL/min, the elimination half-life was approximately doubled and the total body clearance was approximately one half (10.5 hours [clearance 150 L/h] compared to 5.8 hours [clearance 260 L/h] in healthy subjects). No effect on C_{max} or T_{max} was observed after a single dose.

After intravenous administration to patients with hepatic impairment, the elimination half-life of butorphanol was approximately tripled and total body clearance was approximately one half (half-life 16.8 hours, clearance 92 L/h) compared to healthy subjects (half-life 4.8 hours, clearance 175 L/h). The exposure of hepatically impaired patients to butorphanol was significantly greater (about 2-fold) than that in healthy subjects.

(see **PRECAUTIONS: Hepatic and Renal Disease, Drug Interactions** and **Geriatric Use** and **CLINICAL PHARMACOLOGY: Individualization of Dosage**).

Clinical Trials

The effectiveness of opioid analgesics varies in different pain syndromes. Studies with butorphanol tartrate injection have been performed in postoperative (primarily abdominal and orthopedic) pain and pain during labor and delivery, as preoperative and preanesthetic medication, and as a supplement to balanced anesthesia (see below).

Use in the Management of Pain-Postoperative Pain

The analgesic efficacy of butorphanol tartrate injection in postoperative pain was investigated in several double-blind active-controlled studies involving 958 butorphanol-treated patients. The following doses were found to have approximately equivalent analgesic effect: 2 mg butorphanol, 10 mg morphine, 40 mg pentazocine and 80 mg meperidine.

After intravenous administration of butorphanol tartrate, onset and peak analgesic effect occurred by the time of first observation (30 minutes). After intramuscular administration, pain relief onset occurred at 30 minutes or less, and peak effect occurred between 30 minutes and one hour. The duration of action of butorphanol tartrate injection was 3 to 4 hours when defined as the time necessary for pain intensity to return to pretreatment level or the time to retreatment.

Preanesthetic Medication

Butorphanol tartrate injection, (2 mg and 4 mg) and meperidine (80 mg) were studied for use as preanesthetic medication in hospitalized surgical patients. Patients received a

single intramuscular dose of either butorphanol or meperidine approximately 90 minutes prior to anesthesia. The anesthesia regimen included barbiturate induction, followed by nitrous oxide and oxygen with halothane or enflurane, with or without a muscle relaxant.

Anesthetic preparation was rated as satisfactory in all 42 butorphanol injection patients regardless of the type of surgery.

Balanced Anesthesia

Butorphanol tartrate administered intravenously (mean dose 2 mg) was compared to intravenous morphine sulfate (mean dose 10 mg) as premedication shortly before thiopental induction, followed by balanced anesthesia in 50 ASA Class 1 and 2 patients. Anesthesia was then maintained by repeated intravenous doses, averaging 4.6 mg butorphanol and 22.8 mg morphine per patient.

Anesthetic induction and maintenance were generally rated as satisfactory with both butorphanol injection (25 patients) and morphine (25 patients) regardless of the type of surgery performed. Emergence from anesthesia was comparable with both agents.

Labor

(see **PRECAUTIONS**)

The analgesic efficacy of intravenous butorphanol tartrate injection was studied in pain during labor. In a total of 145 patients butorphanol (1 mg and 2 mg) was as effective as 40 mg and 80 mg of meperidine (144 patients) in the relief of pain in labor with no effect on the duration or progress of labor. Both drugs readily crossed the placenta and entered fetal circulation. The condition of the infants in these studies, determined by Apgar scores at 1 and 5 minutes (8 or above) and time to sustained respiration, showed that butorphanol had the same effects on the infants as meperidine.

In these studies neurobehavioral testing in infants exposed to butorphanol injection at a mean of 18.6 hours after delivery, showed no significant differences between treatment groups.

Individualization of Dosage

Use of butorphanol in geriatric patients, patients with renal impairment, patients with hepatic impairment and during labor requires extra caution (see below and the appropriate sections in **PRECAUTIONS**).

For pain relief the recommended initial dosage regimen of butorphanol tartrate injection is 1 mg IV or 2 mg IM with repeated doses every three to four hours as necessary. This dosage regimen is likely to be effective for the majority of patients. Dosage adjustments of butorphanol injection should be based on observations of its beneficial and adverse effects. The initial dose in the elderly and in patients with renal or hepatic impairment should generally be half the recommended adult dose (0.5 mg IV and 1 mg IM). Repeat doses in these patients should be determined by the patient's response rather than at fixed intervals but will generally be no less than 6 hours (see **PRECAUTIONS**).

The usual preoperative dose is 2 mg IM given 60 to 90 minutes before surgery or 2 mg IV shortly before induction. This is approximately equivalent in sedative effect to 10 mg morphine or 80 mg of meperidine. This single preoperative dose should be individualized based on age, body weight, physical status, underlying pathological condition, use of

other drugs, type of anesthesia to be used and the surgical procedure involved.

During maintenance in balanced anesthesia the usual incremental dose of butorphanol tartrate is 0.5 to 1 mg IV. The incremental dose may be higher, up to 0.06 mg/kg (4 mg/70 kg), depending on previous sedative, analgesic, and hypnotic drugs administered. The total dose of butorphanol injection will vary; however, patients seldom require less than 4 mg or more than 12.5 mg (approximately 0.06 to 0.18 mg/kg).

As with other opioids of this class, butorphanol injection may not provide adequate intraoperative analgesia in every patient or under all conditions. A failure to achieve successful analgesia during balanced anesthesia is commonly reflected by increases in general sympathetic tone. Consequently, if blood pressure or heart rate continue to rise, consideration should be given to adding a potent volatile liquid inhalation anesthetic or another intravenous medication.

In labor, the recommended initial dose of butorphanol tartrate is 1 or 2 mg IM or IV in mothers with fetuses of 37 weeks gestation or beyond and without signs of fetal distress. Dosage adjustments of butorphanol in labor should be based on initial response with consideration given to concomitant analgesic or sedative drugs and the expected time of delivery. A dose should not be repeated in less than four hours nor administered less than four hours prior to the anticipated delivery (see **PRECAUTIONS**).

INDICATIONS AND USAGE

Butorphanol Tartrate Injection is indicated

- as a preoperative or pre-anesthetic medication
- as a supplement to balanced anesthesia
- for the relief of pain during labor, and
- for the management of pain severe enough to require an opioid analgesic and for which alternative treatments are inadequate.

Limitations of Use

Because of the risks of addiction, abuse, and misuse, with opioids, even at recommended doses [see **WARNINGS**], reserve butorphanol tartrate for use in patients for whom alternative treatment options [e.g. non-opioid analgesics]

- Have not been tolerated, or are not expected to be tolerated
- Have not provided adequate analgesia, or are not expected to provide adequate analgesia

CONTRAINDICATIONS

Butorphanol Tartrate Injection is contraindicated in:

- Patients with significant respiratory depression (see **WARNINGS**)
- Patients with acute or severe bronchial asthma in an unmonitored setting or in the absence of resuscitative equipment (see **WARNINGS**)
- Patients with known or suspected gastrointestinal obstruction, including paralytic ileus (see **WARNINGS**)
- Patients with hypersensitivity to butorphanol tartrate or any of the formulation

excipients (e.g., anaphylaxis) (see **WARNINGS**)

WARNINGS

Addiction, Abuse, and Misuse

Butorphanol tartrate injection is a Schedule IV controlled substance. As an opioid, butorphanol tartrate exposes users to the risks of addiction, abuse, and misuse [see **DRUG ABUSE AND DEPENDENCE**].

Although the risk of addiction in any individual is unknown, it can occur in patients appropriately prescribed butorphanol tartrate. Addiction can occur at recommended dosages and if the drug is misused or abused.

Assess each patient's risk for opioid addiction, abuse, or misuse prior to prescribing butorphanol tartrate injection, and monitor all patients receiving butorphanol tartrate for the development of these behaviors or conditions. Risks are increased in patients with a personal or family history of substance abuse (including drug or alcohol abuse or addiction) or mental illness (e.g., major depression). The potential for these risks should not, however, prevent the proper management of pain in any given patient. Patients at increased risk may be prescribed opioids such as butorphanol tartrate, but use in such patients necessitates intensive counseling about the risks and proper use of butorphanol tartrate along with intensive monitoring for signs of addiction, abuse, and misuse.

Opioids are sought by drug abusers and people with addiction disorders and are subject to criminal diversion. Consider these risks when prescribing or dispensing butorphanol tartrate. Strategies to reduce these risks include prescribing the drug in the smallest appropriate quantity. Contact local state professional licensing board or state controlled substances authority for information on how to prevent and detect abuse or diversion of this product.

Life-Threatening Respiratory Depression

Serious, life-threatening, or fatal respiratory depression has been reported with the use of opioids, even when used as recommended. Respiratory depression, if not immediately recognized and treated, may lead to respiratory arrest and death. Management of respiratory depression may include close observation, supportive measures, and use of opioid antagonists, depending on the patient's clinical status [see **OVERDOSAGE**]. Carbon dioxide (CO₂) retention from opioid-induced respiratory depression can exacerbate the sedating effects of opioids.

While serious, life-threatening, or fatal respiratory depression can occur at any time during the use of butorphanol tartrate, the risk is greatest during the initiation of therapy or following a dosage increase. Monitor patients closely for respiratory depression, especially within the first 24 to 72 hours of initiating therapy with and following dosage increases of butorphanol tartrate.

To reduce the risk of respiratory depression, proper dosing and titration of butorphanol tartrate are essential [see **DOSAGE AND ADMINISTRATION**]. Overestimating the butorphanol tartrate dosage when converting patients from another opioid product can result in a fatal overdose with the first dose.

Opioids can cause sleep-related breathing disorders including central sleep apnea (CSA) and sleep-related hypoxemia. Opioid use increases the risk of CSA in a dose-dependent fashion. In patients who present with CSA, consider decreasing the opioid dosage using best practices for opioid taper [see **DOSAGE AND ADMINISTRATION**].

Neonatal Opioid Withdrawal Syndrome

Prolonged use of butorphanol tartrate during pregnancy can result in withdrawal in the neonate. Neonatal opioid withdrawal syndrome, unlike opioid withdrawal syndrome in adults, may be life-threatening if not recognized and treated, and requires management according to protocols developed by neonatology experts. If opioid use is required for a prolonged period in a pregnant woman, advise the patient of the risk of neonatal opioid withdrawal syndrome and ensure that appropriate treatment will be available [see **PRECAUTIONS; Information for Patients, Pregnancy**].

Risks from Concomitant Use with Benzodiazepines or Other CNS Depressants

Profound sedation, respiratory depression, coma, and death may result from the concomitant use of butorphanol tartrate with benzodiazepines or other CNS depressants (e.g., non-benzodiazepine sedatives/hypnotics, anxiolytics, tranquilizers, muscle relaxants, general anesthetics, antipsychotics, other opioids, alcohol). Because of these risks, reserve concomitant prescribing of these drugs for use in patients for whom alternative treatment options are inadequate.

Observational studies have demonstrated that concomitant use of opioid analgesics and benzodiazepines increases the risk of drug-related mortality compared to use of opioid analgesics alone. Because of similar pharmacological properties, it is reasonable to expect similar risk with the concomitant use of other CNS depressant drugs with opioid analgesics [see **PRECAUTIONS; Drug Interactions**].

If the decision is made to prescribe a benzodiazepine or other CNS depressant concomitantly with an opioid analgesic, prescribe the lowest effective dosages and minimum durations of concomitant use. In patients already receiving an opioid analgesic, prescribe a lower initial dose of the benzodiazepine or other CNS depressant than indicated in the absence of an opioid, and titrate based on clinical response. If an opioid analgesic is initiated in a patient already taking a benzodiazepine or other CNS depressant, prescribe a lower initial dose of the opioid analgesic, and titrate based on clinical response. Follow patients closely for signs and symptoms of respiratory depression and sedation.

Advise both patients and caregivers about the risks of respiratory depression and sedation when butorphanol tartrate is used with benzodiazepines or other CNS depressants (including alcohol and illicit drugs). Advise patients not to drive or operate heavy machinery until the effects of concomitant use of the benzodiazepine or other CNS depressant have been determined. Screen patients for risk of substance use disorders, including opioid abuse and misuse, and warn them of the risk for overdose and death associated with the use of additional CNS depressants including alcohol and illicit drugs see **PRECAUTIONS; Information for Patients, Drug Interactions**.

Life-Threatening Respiratory Depression in Patients with Chronic Pulmonary Disease or in Elderly, Cachectic, or Debilitated Patients

The use of butorphanol tartrate in patients with acute or severe bronchial asthma in an unmonitored setting or in the absence of resuscitative equipment is contraindicated.

Patients with Chronic Pulmonary Disease

Butorphanol tartrate-treated patients with significant chronic obstructive pulmonary disease or cor pulmonale, and those with a substantially decreased respiratory reserve, hypoxia, hypercapnia, or pre-existing respiratory depression are at increased risk of decreased respiratory drive including apnea, even at recommended dosages of butorphanol tartrate [see **WARNINGS**].

Elderly, Cachectic, or Debilitated Patients

Life-threatening respiratory depression is more likely to occur in elderly, cachectic, or debilitated patients because they may have altered pharmacokinetics or altered clearance compared to younger, healthier patients [see **WARNINGS**].

Monitor such patients closely, particularly when initiating and titrating butorphanol tartrate and when butorphanol tartrate is given concomitantly with other drugs that depress respiration [see **WARNINGS**]. Alternatively, consider the use of non-opioid analgesics in these patients.

Adrenal Insufficiency

Cases of adrenal insufficiency have been reported with opioid use, more often following greater than 1 month of use. Presentation of adrenal insufficiency may include non-specific symptoms and signs including nausea, vomiting, anorexia, fatigue, weakness, dizziness, and low blood pressure. If adrenal insufficiency is suspected, confirm the diagnosis with diagnostic testing as soon as possible. If adrenal insufficiency is diagnosed, treat with physiologic replacement doses of corticosteroids. Wean the patient off of the opioid to allow adrenal function to recover and continue corticosteroid treatment until adrenal function recovers. Other opioids may be tried as some cases reported use of a different opioid without recurrence of adrenal insufficiency. The information available does not identify any particular opioids as being more likely to be associated with adrenal insufficiency.

Risks of Use in Patients with Increased Intracranial Pressure, Brain Tumors, Head Injury, or Impaired Consciousness

In patients who may be susceptible to the intracranial effects of CO₂ retention (e.g., those with evidence of increased intracranial pressure or brain tumors), Butorphanol Tartrate Injection may reduce respiratory drive, and the resultant CO₂ retention can further increase intracranial pressure. Monitor such patients for signs of sedation and respiratory depression, particularly when initiating therapy with Butorphanol Tartrate Injection.

Opioids may also obscure the clinical course in a patient with a head injury. Avoid the use of Butorphanol Tartrate Injection in patients with impaired consciousness or coma.

Risks of Use in Patients with Gastrointestinal Conditions

Butorphanol Tartrate Injection is contraindicated in patients with gastrointestinal obstruction, including paralytic ileus.

Butorphanol in Butorphanol Tartrate Injection may cause spasm of the sphincter of Oddi. Opioids may cause increases in serum amylase. Monitor patients with biliary tract disease, including acute pancreatitis, for worsening symptoms.

Increased Risk of Seizures in Patients with Seizure Disorders

The butorphanol in Butorphanol Tartrate Injection may increase the frequency of seizures in patients with seizure disorders, and may increase the risk of seizures occurring in other clinical settings associated with seizures. Monitor patients with a history of seizure disorders for worsened seizure control during Butorphanol Tartrate Injection therapy.

Withdrawal

The use of Butorphanol Tartrate Injection, a mixed agonist/antagonist opioid analgesic, in patients who are receiving a full opioid agonist analgesic may reduce the analgesic effect and/or precipitate withdrawal symptoms. Avoid concomitant use of Butorphanol Tartrate Injection with a full opioid agonist analgesic. When discontinuing Butorphanol Tartrate Injection, gradually taper the dosage (**see DOSAGE AND ADMINISTRATION**). Do not abruptly discontinue Butorphanol Tartrate Injection (**see DRUG ABUSE AND DEPENDENCE**).

Cardiovascular Effects

Because butorphanol may increase the work of the heart, especially the pulmonary circuit, the use of butorphanol in patients with acute myocardial infarction, ventricular dysfunction, or coronary insufficiency should be limited to those situations where the benefits clearly outweigh the risk (see **CLINICAL PHARMACOLOGY**).

Severe hypertension has been reported rarely during butorphanol therapy. In such cases, butorphanol should be discontinued and the hypertension treated with antihypertensive drugs. In patients who are not opioid dependent, naloxone has also been reported to be effective.

PRECAUTIONS

General

Hypotension associated with syncope during the first hour of dosing with Butorphanol Tartrate Injection has been reported rarely, particularly in patients with past history of similar reactions to opioid analgesics. Therefore, patients should be advised to avoid activities with potential risks.

Risks of Driving and Operating Machinery

Butorphanol Tartrate Injection may impair the mental or physical abilities needed to perform potentially hazardous activities such as driving a car or operating machinery. Warn patients not to drive or operate dangerous machinery unless they are tolerant to the effects of Butorphanol Tartrate Injection and know how they will react to the medication (see PRECAUTIONS; Information for Patients/Caregivers).

Disorders of Respiratory Function or Control

Butorphanol may produce respiratory depression, especially in patients receiving other CNS active agents, or patients suffering from CNS diseases or respiratory impairment.

Hepatic and Renal Disease

In patients with hepatic or renal impairment, the initial dose of Butorphanol Tartrate Injection should generally be half the recommended adult dose (0.5 mg IV and 1 mg IM). Repeat doses in these patients should be determined by the patient's response rather than at fixed intervals but will generally be no less than 6 hours apart (see **CLINICAL PHARMACOLOGY: Pharmacokinetics and Individualization of Dosage** sections).

Information for Patients

Addiction, Abuse, and Misuse

Inform patients that the use of butorphanol tartrate, even when taken as recommended, can result in addiction, abuse, and misuse, which can lead to overdose and death [see **WARNINGS**]. Instruct patients not to share butorphanol tartrate with others and to take steps to protect butorphanol tartrate from theft or misuse.

Life-Threatening Respiratory Depression

Inform patients of the risk of life-threatening respiratory depression, including information that the risk is greatest when starting butorphanol tartrate or when the dosage is increased, and that it can occur even at recommended dosages [see **WARNINGS**]. Advise patients how to recognize respiratory depression and to seek medical attention if breathing difficulties develop.

Interactions with Benzodiazepines and Other CNS Depressants

Inform patients and caregivers that potentially fatal additive effects may occur if butorphanol tartrate is used with benzodiazepines or other CNS depressants, including alcohol, and not to use these concomitantly unless supervised by a health care provider [see **WARNINGS, PRECAUTIONS; Drug Interactions**].

Serotonin Syndrome

Inform patients that butorphanol tartrate could cause a rare but potentially life-threatening condition resulting from concomitant administration of serotonergic drugs. Warn patients of the symptoms of serotonin syndrome and to seek medical attention right away if symptoms develop. Instruct patients to inform their physicians if they are taking, or plan to take serotonergic medications. [see **PRECAUTIONS; Drug Interactions**]

Anaphylaxis

Inform patients that anaphylaxis has been reported with ingredients contained in Butorphanol Tartrate Injection. Advise patients how to recognize such a reaction and when to seek medical attention (see **CONTRAINDICATIONS, ADVERSE REACTIONS**).

Pregnancy

Neonatal Opioid Withdrawal Syndrome

Inform patients that prolonged use of butorphanol tartrate during pregnancy can result in neonatal opioid withdrawal syndrome, which may be life-threatening if not recognized and treated [see **WARNINGS, PRECAUTIONS; Pregnancy**]

Embryo-Fetal Toxicity

Inform female patients of reproductive potential that butorphanol tartrate can cause fetal harm and to inform the prescriber of a known or suspected pregnancy [see **PRECAUTIONS; Pregnancy**].

Lactation

Advise nursing mothers to monitor infants for increased sleepiness (more than usual), breathing difficulties, or limpness. Instruct nursing mothers to seek immediate medical care if they notice these signs [see **PRECAUTIONS; Nursing Mothers**].

Infertility

Inform patients that chronic use of opioids may cause reduced fertility. It is not known whether these effects on fertility are reversible (see **ADVERSE REACTIONS**).

Driving or Operating Heavy Machinery

Inform patients that Butorphanol Tartrate Injection may impair the mental and/or physical abilities required for the performance of potentially hazardous tasks such as driving a car or operating machinery and to avoid such tasks while taking this product, until they know how they will react to the medication.

Drug Interactions

Benzodiazepines and Other Central Nervous System (CNS) Depressants

Due to additive pharmacologic effect, the concomitant use of benzodiazepines or other CNS depressants such as alcohol, other sedatives/hypnotics, anxiolytics, tranquilizers, muscle relaxants, general anesthetics, antipsychotics, and other opioids, can increase the risk of respiratory depression, profound sedation, coma, and death.

Reserve concomitant prescribing of these drugs for use in patients for whom alternative treatment options are inadequate. Limit dosages and durations to the minimum required. Follow patients closely for signs of respiratory depression and sedation [see **WARNINGS**].

Serotonergic Drugs

The concomitant use of opioids with other drugs that affect the serotonergic neurotransmitter system, such as selective serotonin reuptake inhibitors (SSRIs), serotonin and norepinephrine reuptake inhibitors (SNRIs), tricyclic antidepressants (TCAs), triptans, 5-HT₃ receptor antagonists, drugs that effect the serotonin neurotransmitter system (e.g., mirtazapine, trazodone, tramadol), certain muscle relaxants (i.e., cyclobenzaprine, metaxalone), and monoamine oxidase (MAO) inhibitors (those intended to treat psychiatric disorders and also others, such as linezolid and intravenous methylene blue), has resulted in serotonin syndrome. [see **PRECAUTIONS; INFORMATION FOR PATIENTS**]

If concomitant use is warranted, carefully observe the patient, particularly during treatment initiation and dose adjustment. Discontinue butorphanol tartrate if serotonin syndrome is suspected.

Cytochrome P450 (CYP 450) Interactions

It is not known if the effects of Butorphanol Tartrate Injection are altered by concomitant medications that affect hepatic metabolism of drugs (CYP 450 inhibitors or inducers) (e.g., erythromycin, theophylline, etc.), but physicians should be alert to the possibility that a smaller initial dose and longer intervals between doses may be needed.

Monoamine Oxidase inhibitors (MAOIs)

No information is available about the use of butorphanol concurrently with MAO inhibitors.

Advise patient to avoid concomitant use of these drugs.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Two-year carcinogenicity studies were conducted in mice and rats given butorphanol tartrate in the diet up to 60 mg/kg/day (12 and 24 times the human daily dose of 24 mg/day based on a body surface area comparison, respectively). There was no evidence of carcinogenicity in either species in these studies.

Mutagenesis

Butorphanol was not genotoxic in the in vitro bacterial reverse mutation assay (Ames) or in an in vitro unscheduled DNA synthesis and repair assay conducted in cultured human fibroblast cells.

Impairment of Infertility

In a study where male rats were treated subcutaneously with 0.5 or 2.5 mg/kg butorphanol for 75 days prior to mating to female rats treated subcutaneously with 0.5 or 2.5 mg/kg butorphanol for 14-days prior to mating and throughout gestation and lactation, no adverse effects on fertility were noted (0.2- and 1-times the human daily dose of 24 mg based on body surface area).

In a study where male rats were treated orally with 10, 40, or 160 mg/kg for 63 days prior to mating to female rats treated orally with the same doses of butorphanol for 14 days prior to mating, reduced pregnancy rates were reported in the high dose group (65-times the human daily dose of 24 mg based on body surface area).

Pregnancy

Pregnancy Category C

Reproduction studies in mice, rats, and rabbits during organogenesis did not reveal any teratogenic potential to butorphanol. However, pregnant rats treated subcutaneously with butorphanol at 1 mg/kg (5.9 mg/m²) had a higher frequency of stillbirths than controls. Butorphanol at 30 mg/kg/oral (360 mg/m²) and 60 mg/kg/oral (720 mg/m²)

also showed higher incidences of post-implantation loss in rabbits.

There are no adequate and well-controlled studies of Butorphanol Tartrate Injection in pregnant women before 37 weeks of gestation. Butorphanol Tartrate Injection should be used during pregnancy only if the potential benefit justifies the potential risk to the infant.

Fetal/Neonatal Adverse Reactions

Prolonged use of opioid analgesics during pregnancy for medical or nonmedical purposes can result in physical dependence in the neonate and neonatal opioid withdrawal syndrome shortly after birth.

Neonatal opioid withdrawal syndrome presents as irritability, hyperactivity and abnormal sleep pattern, high pitched cry, tremor, vomiting, diarrhea and failure to gain weight. The onset, duration, and severity of neonatal opioid withdrawal syndrome vary based on the specific opioid used, duration of use, timing and amount of last maternal use, and rate of elimination of the drug by the newborn. Observe newborns for symptoms of neonatal opioid withdrawal syndrome and manage accordingly [see **WARNINGS**].

Labor or Delivery

Opioids cross the placenta and may produce respiratory depression and psychophysiologic effects in neonates. An opioid antagonist, such as naloxone, must be available for reversal of opioid-induced respiratory depression in the neonate. Butorphanol tartrate is not recommended for use in pregnant women during or immediately prior to labor, when other analgesic techniques are more appropriate. Opioid analgesics, including butorphanol tartrate, can prolong labor through actions which temporarily reduce the strength, duration, and frequency of uterine contractions. However, this effect is not consistent and may be offset by an increased rate of cervical dilation, which tends to shorten labor. Monitor neonates exposed to opioid analgesics during labor for signs of excess sedation and respiratory depression.

Butorphanol has been detected in milk following administration of Butorphanol Tartrate Injection to nursing mothers. The amount an infant would receive is probably clinically insignificant (estimated 4 µg/L of milk in a mother receiving 2 mg IM four times a day).

There have been rare reports of infant respiratory distress/apnea following the administration of Butorphanol Tartrate Injection during labor. The reports of respiratory distress/apnea have been associated with administration of a dose within 2 hours of delivery, use of multiple doses, use with additional analgesic or sedative drugs, or use in preterm pregnancies (see **OVERDOSAGE: Treatment**). In a study of 119 patients, the administration of 1 mg of IV Butorphanol Tartrate Injection during labor was

associated with transient (10–90 minutes) sinusoidal fetal heart rate patterns, but 16 was not associated with adverse neonatal outcomes. In the presence of an abnormal fetal heart rate pattern, Butorphanol Tartrate Injection should be used with caution.

Nursing Mothers

Butorphanol has been detected in milk following administration of Butorphanol Tartrate Injection to nursing mothers. The amount an infant would receive is probably clinically insignificant (estimated 4 µg/L of milk in a mother receiving 2 mg IM four times a day).

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for butorphanol tartrate and any potential adverse effects on the breastfed infant from butorphanol tartrate or from the underlying maternal condition.

Infants exposed to butorphanol tartrate through breast milk should be monitored for excess sedation and respiratory depression. Withdrawal symptoms can occur in breastfed infants when maternal administration of an opioid analgesic is stopped, or when breast-feeding is stopped.

Pediatric Use

Butorphanol is not recommended for use in patients below 18 years of age because safety and efficacy have not been established in this population.

Geriatric Use

Elderly patients (aged 65 years or older) may have increased sensitivity to butorphanol tartrate. In general, use caution when selecting a dosage for an elderly patient, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function and of concomitant disease or other drug therapy.

Respiratory depression is the chief risk for elderly patients treated with opioids, and has occurred after large initial doses were administered to patients who were not opioid-tolerant or when opioids were co-administered with other agents that depress respiration. Titrate the dosage of butorphanol tartrate slowly in geriatric patients [see **WARNINGS**].

This drug is known to be substantially excreted by the kidney, and the risk of adverse reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

ADVERSE REACTIONS

Clinical Trial Experience

A total of 1658 patients were studied in premarketing clinical trials of Butorphanol Tartrate Injection. In nearly all cases the type and incidence of side effects with butorphanol were those commonly observed with opioid analgesics.

The adverse experiences described below are based on data from short- and long-term clinical trials in patients receiving Butorphanol Tartrate Injection.

The most frequently reported adverse experiences across all clinical trials with Butorphanol Tartrate Injection and Nasal Spray were somnolence (43%), dizziness (19%), nausea and/or vomiting (13%). The following adverse experiences were reported at a frequency of 1% or greater in clinical trials and were considered to be probably related to the use of butorphanol:

Body as a Whole: Asthenia/Lethargy, Headache, Sensation of Heat

Cardiovascular: Vasodilation, Palpitations

Digestive: Anorexia, Constipation, Dry Mouth, Nausea and/or Vomiting, Stomach Pain

Nervous: Anxiety, Confusion, Dizziness, Euphoria, Floating Feeling, Insomnia, Nervousness, Paresthesia, Somnolence, Tremor

Respiratory: Cough, Dyspnea

Skin and Appendages: Sweating, Pruritus

Special Senses: Blurred Vision, Ear Pain, Tinnitus, Unpleasant Taste

The following adverse experiences were reported with a frequency of less than 1% in clinical trials and were considered to be probably related to the use of butorphanol:

Cardiovascular: Hypotension, Syncope

Nervous: Abnormal Dreams, Agitation, Dysphoria, Hallucinations, Hostility, Withdrawal Symptoms

Skin and Appendages: Rash/Hives

Urogenital: Impaired Urination

The following infrequent additional adverse experiences were reported in a frequency of less than 1% of the patients studied in short-term butorphanol tartrate nasal sprays trials and under circumstances where the association between these events and butorphanol administration is unknown. They are being listed as alerting information for the physician due to their clinical significance:

Body as a Whole: Edema

Cardiovascular: Chest Pain, Hypertension, Tachycardia

Nervous: Depression

Respiratory: Shallow Breathing

Postmarketing Experience

The following adverse reactions have been identified during post approval use of Butorphanol Tartrate Injection. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

- Serotonin syndrome: Cases of serotonin syndrome, a potentially life-threatening condition, have been reported during concomitant use of opioids with serotonergic drugs.
- Adrenal insufficiency: Cases of adrenal insufficiency have been reported with opioid use, more often following greater than one month of use.
- Anaphylaxis: Anaphylaxis has been reported with ingredients contained in Butorphanol Tartrate Injection.
- Androgen deficiency: Cases of androgen deficiency have occurred with chronic use of opioids (see **CLINICAL PHARMACOLOGY**).

DRUG ABUSE AND DEPENDENCE

Controlled Substance

Butorphanol Tartrate Injection contains butorphanol, a Schedule IV controlled substance.

Abuse

Butorphanol Tartrate Injection contains butorphanol, a substance with a high potential for abuse similar to other opioids including fentanyl, hydrocodone, hydromorphone, methadone, morphine, oxycodone, oxymorphone, and tapentadol. Butorphanol Tartrate Injection can be abused and is subject to misuse, addiction, and criminal diversion (see **WARNINGS**). All patients treated with opioids require careful monitoring for signs of abuse and addiction, since use of opioid analgesic products carries the risk of addiction even under appropriate medical use.

Prescription drug abuse is the intentional non-therapeutic use of a prescription drug, even once, for its rewarding psychological or physiological effects.

Drug addiction is a cluster of behavioral, cognitive, and physiological phenomena that develop after repeated substance use and includes: a strong desire to take the drug, difficulties in controlling its use, persisting in its use despite harmful consequences, a higher priority given to drug use than to other activities and obligations, increased tolerance, and sometimes a physical withdrawal.

"Drug-seeking" behavior is very common in persons with substance use disorders. Drug-seeking tactics include emergency calls or visits near the end of office hours, refusal to undergo appropriate examination, testing, or referral, repeated "loss" of prescriptions, tampering with prescriptions and reluctance to provide prior medical records or contact information for other treating health care provider(s). "Doctor shopping" (visiting multiple prescribers) to obtain additional prescriptions is common among drug abusers and people suffering from untreated addiction. Preoccupation with achieving adequate pain relief can be appropriate behavior in a patient with poor pain control.

Abuse and addiction are separate and distinct from physical dependence and tolerance. Health care providers should be aware that addiction may not be accompanied by concurrent tolerance and symptoms of physical dependence in all addicts. In addition, abuse of opioids can occur in the absence of true addiction.

Butorphanol tartrate, like other opioids, can be diverted for non-medical use into illicit channels of distribution. Careful record-keeping of prescribing information, including quantity, frequency, and renewal requests, as required by state and federal law, is strongly advised.

Proper assessment of the patient, proper prescribing practices, periodic re-evaluation of therapy, and proper dispensing and storage are appropriate measures that help to limit abuse of opioid drugs.

Risks Specific to Abuse of Butorphanol Tartrate Injection

Abuse of Butorphanol Tartrate Injection poses a risk of overdose and death. The risk is increased with concurrent abuse of Butorphanol Tartrate Injection with alcohol and other central nervous system depressants.

Parenteral drug abuse is commonly associated with transmission of infectious diseases

such as hepatitis and HIV.

Dependence

Both tolerance and physical dependence can develop during chronic opioid therapy. Tolerance is the need for increasing doses of opioids to maintain a defined effect such as analgesia (in the absence of disease progression or other external factors). Tolerance may occur to both the desired and undesired effects of drugs, and may develop at different rates for different effects.

Physical dependence results in withdrawal symptoms after abrupt discontinuation or a significant dosage reduction of a drug. Withdrawal also may be precipitated through the administration of drugs with opioid antagonist activity (e.g., naloxone, nalmefene), mixed agonist/antagonist analgesics (pentazocine, butorphanol, nalbuphine), or partial agonists (buprenorphine). Physical dependence may not occur to a clinically significant degree until after several days to weeks of continued opioid usage.

Butorphanol tartrate should not be abruptly discontinued [see **DOSAGE AND ADMINISTRATION**]. If butorphanol tartrate is abruptly discontinued in a physically-dependent patient, a withdrawal syndrome may occur. Some or all of the following can characterize this syndrome: restlessness, lacrimation, rhinorrhea, yawning, perspiration, chills, myalgia, and mydriasis. Other signs and symptoms also may develop, including: irritability, anxiety, backache, joint pain, weakness, abdominal cramps, insomnia, nausea, anorexia, vomiting, diarrhea, or increased blood pressure, respiratory rate, or heart rate.

Infants born to mothers physically dependent on opioids will also be physically dependent and may exhibit respiratory difficulties and withdrawal signs [see **PRECAUTIONS; Pregnancy**].

OVERDOSAGE

Clinical Presentation

Acute overdose with butorphanol tartrate can be manifested by respiratory depression, somnolence progressing to stupor or coma, skeletal muscle flaccidity, cold and clammy skin, constricted pupils, and, in some cases, pulmonary edema, bradycardia, hypotension, partial or complete airway obstruction, atypical snoring, and death. Marked mydriasis rather than miosis may be seen with hypoxia in overdose situations.

Treatment of Overdose

In case of overdose, priorities are the reestablishment of a patent and protected airway and institution of assisted or controlled ventilation, if needed. Employ other supportive measures (including oxygen and vasopressors) in the management of circulatory shock and pulmonary edema as indicated. Cardiac arrest or arrhythmias will require advanced life-support techniques.

The opioid antagonist, naloxone or nalmefene, are specific antidotes to respiratory depression resulting from opioid overdose. For clinically significant respiratory or circulatory depression secondary to butorphanol tartrate overdose, administer an opioid antagonist. As butorphanol is a mixed opioid agonist/antagonist, larger doses of

naloxone or nalmefene may be needed to reverse the effects of an overdose.

Opioid antagonists should not be administered in the absence of clinically significant respiratory or circulatory depression secondary to butorphanol tartrate overdose.

Because the duration of opioid reversal is expected to be less than the duration of action of butorphanol in Butorphanol Tartrate Injection, carefully monitor the patient until spontaneous respiration is reliably re-established. If the response to an opioid antagonist is suboptimal or only brief in nature, administer additional antagonist as directed by the product's prescribing information.

In an individual physically dependent on opioids, administration of the recommended usual dosage of the antagonist will precipitate an acute withdrawal syndrome. The severity of the withdrawal symptoms experienced will depend on the degree of physical dependence and the dose of the antagonist administered. If a decision is made to treat serious respiratory depression in the physically dependent patient, administration of the antagonist should be begun with care and by titration with smaller than usual doses of the antagonist.

DOSAGE AND ADMINISTRATION

Important Dosage and Administration Instructions

Use the lowest effective dosage for the shortest duration consistent with individual patient treatment goals (see **WARNINGS**).

Initiate the dosing regimen for each patient individually, taking into account the patient's severity of pain, patient response, prior analgesic treatment experience, and risk factors for addiction, abuse, and misuse (see **WARNINGS**).

Monitor patients closely for respiratory depression, especially within the first 24 to 72 hours of initiating therapy and following dosage increases with Butorphanol Tartrate Injection and adjust the dosage accordingly (see **WARNINGS**).

Initial Dosage

Factors to be considered in determining the dose are age, body weight, physical status, underlying pathological condition, use of other drugs, type of anesthesia to be used, and surgical procedure involved. Use in the elderly, patients with hepatic or renal disease, or in labor requires extra caution (see **PRECAUTIONS** section and **CLINICAL PHARMACOLOGY: Individualization of Dosage** section). The following doses are for patients who do not have impaired hepatic or renal function and who are not on CNS active agents.

Use for Pain

Intravenous

The usual recommended single dose for IV administration is 1 mg repeated every three to four hours as necessary. The effective dosage range, depending on the severity of pain, is 0.5 to 2 mg repeated every three to four hours.

Intramuscular

The usual recommended single dose for IM administration is 2 mg in patients who will be able to remain recumbent, in the event drowsiness or dizziness occurs. This may be repeated every three to four hours, as necessary. The effective dosage range depending on the severity of pain is 1 to 4 mg repeated every three to four hours. There are insufficient clinical data to recommend single doses above 4 mg.

Use as Preoperative/Preanesthetic Medication

The preoperative medication dosage of butorphanol tartrate injection should be individualized (see **CLINICAL PHARMACOLOGY: Individualization of Dosage** section). The usual adult dose is 2 mg IM, administered 60 to 90 minutes before surgery. This is approximately equivalent in sedative effect to 10 mg morphine or 80 mg meperidine.

Use in Balanced Anesthesia

The usual dose of butorphanol tartrate injection is 2 mg IV shortly before induction and/or 0.5 to 1 mg IV in increments during anesthesia. The increment may be higher, up to 0.06 mg/kg (4 mg/70 kg), depending on previous sedative, analgesic, and hypnotic drugs administered. The total dose of butorphanol injection will vary; however, patients seldom require less than 4 mg or more than 12.5 mg (approximately 0.06 to 0.18 mg/kg).

Labor

In patients at full term in early labor a 1 to 2 mg dose of butorphanol tartrate IV or IM may be administered and repeated after 4 hours. Alternative analgesia should be used for pain associated with delivery or if delivery is expected to occur within 4 hours.

If concomitant use of butorphanol with drugs that may potentiate its effects is deemed necessary (see **PRECAUTIONS: Drug Interactions**) the lowest effective dose should be employed.

Dosage Modifications in Elderly Patients and Patients with Renal or Hepatic Impairment

The initial dose sequence in elderly patients and patients with hepatic or renal impairment should be limited to 1 mg followed, if needed, by 1 mg in 90 to 120 minutes. The repeat dose sequence should be determined by the patient's response rather than at fixed times but will generally be no less than at 6 hours intervals [see **CLINICAL PHARMACOLOGY: Individualization of Dosage** and **PRECAUTIONS**].

Titration and Maintenance of Therapy

Individually titrate Butorphanol Tartrate Injection to a dose that provides adequate analgesia and minimizes adverse reactions. Continually reevaluate patients receiving Butorphanol Tartrate Injection to assess the maintenance of pain control and the relative incidence of adverse reactions, as well as monitoring for the development of addiction, abuse, or misuse (see **WARNINGS**). Frequent communication is important among the prescriber, other members of the healthcare team, the patient, and the caregiver/family during periods of changing analgesic requirements, including initial titration.

If the level of pain increases after dosage stabilization, attempt to identify the source of

increased pain before increasing the Butorphanol Tartrate Injection dosage. If unacceptable opioid-related adverse reactions are observed, consider reducing the dosage. Adjust the dosage to obtain an appropriate balance between management of pain and opioid-related adverse reactions.

Discontinuation of Butorphanol Tartrate Injection

When a patient who has been taking Butorphanol Tartrate Injection regularly and may be physically dependent no longer requires therapy with Butorphanol Tartrate Injection, taper the dose gradually, by 25% to 50% every 2 to 4 days, while monitoring carefully for signs and symptoms of withdrawal. If the patient develops these signs or symptoms, raise the dose to the previous level and taper more slowly, either by increasing the interval between decreases, decreasing the amount of change in dose, or both. Do not abruptly discontinue Butorphanol Tartrate Injection in a physically-dependent patient (see WARNINGS, DRUG ABUSE AND DEPENDENCE).

Safety and Handling

Butorphanol tartrate injection is supplied in sealed delivery systems that have a low risk of accidental exposure to health care workers. Ordinary care should be taken to avoid aerosol generation while preparing a syringe for use. Following skin contact, rinsing with cool water is recommended.

The disposal of Schedule IV controlled substances must be consistent with State and Federal Regulations.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

HOW SUPPLIED

Butorphanol Tartrate Injection, USP is supplied as follows:

Unit of Sale	Concentration (Total Butorphanol Concentration Per Container)
NDC 0409-1623-01 Carton of 10 - 1 mL Single-Dose Glass Fliptop Vials	1 mg/mL (1 mg/1 mL)
NDC 0409-1626-01 Carton of 10 - 1 mL Single-Dose Glass Fliptop Vials	2 mg/mL (2 mg/1 mL)
NDC 0409-1626-02 Carton of 10 - 2 mL Single-Dose Glass Fliptop Vials	2 mg/mL (4 mg/2 mL)

Store at 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature.]

Protect from light.

Distributed by Hospira Inc., Lake Forest, IL 60045 USA

LAB-0831-4.0

Revised: 6/2019

PRINCIPAL DISPLAY PANEL - 2 mg/mL Vial Label

1 mL

Single-dose

Rx only

BUTORPHANOL


TARTRATE Inj., USP

CIV

2 mg/mL

For I.M. or I.V. use

Dist. by Hospira, Inc., Lake Forest, IL 60045 USA



PRINCIPAL DISPLAY PANEL - 2 mg/mL Vial Carton


1 mL Single-dose Fliptop Vials

NDC 0409-1626-01
Contains 10 of NDC 0409-1626-21

BUTORPHANOL TARTRATE
Injection, USP 2 mg/mL
Rx only
CIV

For Intramuscular or Intravenous use Discard unused portion.

Hospira



PRINCIPAL DISPLAY PANEL - 4 mg/2 mL Vial Label


2 mL
Single-dose

Rx only

CIV

**BUTORPHANOL
TARTRATE**
Injection, USP
4 mg/2 mL (2 mg/mL)

Dist. by Hospira, Inc., Lake Forest, IL 60045 USA




PRINCIPAL DISPLAY PANEL - 4 mg/2 mL Vial Carton

2 mL Single-dose Fliptop Vials
 NDC 0409-1626-02
 Contains 10 of NDC 0409-1626-42

BUTORPHANOL TARTRATE
 Injection, USP 4 mg/2 mL (2 mg/mL)
 Rx only
 CIV

For Intramuscular or Intravenous use Discard unused portion.

Hospira



PRINCIPAL DISPLAY PANEL - 1 mg/mL Vial Label


1 mL
Single-dose

Rx only

CIV

BUTORPHANOL
TARTRATE Inj., USP
1 mg/mL

For I.M. or I.V. use
Dist. by Hospira, Inc., Lake Forest, IL 60045 USA



PRINCIPAL DISPLAY PANEL - 1 mg/mL Vial Carton

1 mL Single-dose Fliptop Vials
NDC 0409-1623-01
Contains 10 of NDC 0409-1623-21


BUTORPHANOL TARTRATE
Injection, USP
1 mg/mL

For Intramuscular or Intravenous use
Discard unused portion.

CIV

Rx only

Hospira



BUTORPHANOL TARTRATE

butorphanol tartrate injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0409-1626
Route of Administration	INTRAMUSCULAR, INTRAVENOUS	DEA Schedule	CIV

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BUTORPHANOL TARTRATE (UNII: 2L7I72RUHN) (BUTORPHANOL - UNII:QV897JC36D)	BUTORPHANOL TARTRATE	2 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	6.4 mg in 1 mL
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	3.3 mg in 1 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X)	6.4 mg in 1 mL
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0409-1626-01	10 in 1 CARTON	03/17/2006	

1	NDC:0409-1626-21	1 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		
2	NDC:0409-1626-02	10 in 1 CARTON	12/20/2005	
2	NDC:0409-1626-42	2 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074626	12/20/2005	

BUTORPHANOL TARTRATE

butorphanol tartrate injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0409-1623
Route of Administration	INTRAMUSCULAR, INTRAVENOUS	DEA Schedule	CIV

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BUTORPHANOL TARTRATE (UNII: 2L7I72RUHN) (BUTORPHANOL - UNII:QV897JC36D)	BUTORPHANOL TARTRATE	1 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	6.4 mg in 1 mL
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	3.3 mg in 1 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X)	6.4 mg in 1 mL
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0409-1623-01	10 in 1 CARTON	09/23/2005	
1	NDC:0409-1623-21	1 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
-----------	---------------------------------	-----------------	---------------

Category	Citation	Date	Date
ANDA	ANDA074626	09/23/2005	

Labeler - Hospira, Inc. (141588017)

Establishment

Name	Address	ID/FEI	Business Operations
Hospira, Inc.		093132819	ANALYSIS(0409-1623, 0409-1626) , MANUFACTURE(0409-1623, 0409-1626) , PACK(0409-1623, 0409-1626) , LABEL(0409-1623, 0409-1626)

Establishment

Name	Address	ID/FEI	Business Operations
Hospira, Inc.		030606222	ANALYSIS(0409-1623, 0409-1626)

Establishment

Name	Address	ID/FEI	Business Operations
Hospira, Inc.		827731089	ANALYSIS(0409-1623, 0409-1626)

Revised: 2/2023

Hospira, Inc.