**GOOD NEIGHBOR PHARMACY TUSSIN CF- dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution**
**Amerisource Bergen**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

---

**Amerisource Bergen Tussin CF Drug Facts**

**Active ingredients (in each 10 mL)**

Dextromethorphan HBr, USP 20 mg

Guaifenesin, USP 200 mg

Phenylephrine HCl, USP 10 mg

**Purposes**

Cough suppressant

Expectorant

Nasal decongestant

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
- nasal congestion
- cough due to minor throat and bronchial irritation

**Warnings**

**Do not use**

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

**When using this product**

**do not use more than directed**

**Stop use and ask a doctor if**

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

| age | dose |
|---|---|
| adults and children 12 years and over | 10 mL every 4 hours |
| children under 12 years | do not use |

**Other information**

- **each 10 mL contains:** sodium 5 mg
- store at 20-25ºC (68-77ºF). Do not refrigerate.

**Inactive ingredients**

anhydrous citric acid, edetate disodium, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

**Questions or comments?**

**1-800-719-9260**

**Principal Display Panel**

Compare to Robitussin® Multi-Symptom Cold active ingredients

ADULT

Tussin CF

cough suppressant

(dextromethorphan HBr)
expectorant (guaifenesin)
nasal decongestant
(phenylephrine HCl)
Multi-Symptom Cold
Relieves:
Nasal Congestion
Cough
Mucus
Peak Cold
Non-Drowsy
Alcohol Free
For Ages 12 & Over
8 fl oz (237 mL)

NDC 24385-904-34

Compare to Robitussin® Multi-Symptom Cold active ingredients*

GOOD NEIGHBOR PHARMACY®

ADULT

# Tussin CF

cough suppressant (dextromethorphan HBr)
expectorant (guaifenesin)
nasal decongestant (phenylephrine HCl)

## Multi-Symptom Cold

Relieves:
- Nasal Congestion
- Cough
- Mucus

Peak Cold Non-Drowsy

Alcohol Free
For Ages 12 & Over

8 fl oz (237 mL)

NDC 24385-904-34

Compare to Robitussin® Multi-Symptom Cold active ingredients*

GOOD NEIGHBOR PHARMACY®

ADULT

Tussin CF

cough suppressant (dextromethorphan HBr)
expectorant (guaifenesin)
nasal decongestant (phenylephrine HCl)

Multi-Symptom Cold

Relieves:
- Nasal Congestion
- Cough
- Mucus

Peak Cold Non-Drowsy

Alcohol Free
For Ages 12 & Over

8 fl oz (237 mL)



GOOD NEIGHBOR PHARMACY®

ADULT

# Tussin CF

cough suppressant (dextromethorphan HBr)
expectorant (guaifenesin)
nasal decongestant (phenylephrine HCl)

**PARENTS:**
Learn about teen medicine abuse
www.StopMedicineAbuse.org

## Drug Facts

| Active ingredients (in each 10 mL) | Purposes |
|---|---|
| Dextromethorphan HBr, USP 20 mg | Cough suppressant |
| Guaifenesin, USP 200 mg | Expectorant |
| Phenylephrine HCl, USP 10 mg | Nasal decongestant |

### Uses
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
  - nasal congestion
  - cough due to minor throat and bronchial irritation

### Warnings
**Do not use** if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

**When using this product do not use more than directed**

**Stop use and ask a doctor if**
- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition. ▶

## Drug Facts (continued)

**If pregnant or breast-feeding,** ask a health professional before use.
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### Directions
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

| age | dose |
|---|---|
| adults and children 12 years and over | 10 mL every 4 hours |
| children under 12 years | do not use |

### Other information
- **each 10 mL contains:** sodium 5 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

### Inactive ingredients
anhydrous citric acid, edetate disodium, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

### Questions or comments?
1-800-719-9260

*Good Neighbor Pharmacy® Tussin CF is not manufactured or distributed by Pfizer, distributor of Robitussin® Multi-Symptom Cold.

GLUTEN FREE

Distributed By
AmerisourceBergen
1300 Morris Drive
Chesterbrook, PA 19087
Visit us at
www.goodneighborpharmacy.com

SATISFACTION GUARANTEED

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING




# GOOD NEIGHBOR PHARMACY TUSSIN CF

dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution

| Product Information | | | |
|---|---|---|---|
| **Product Type** | HUMAN OTC DRUG | **Item Code (Source)** | NDC:24385-904 |
| **Route of Administration** | ORAL | | |

| Active Ingredient/Active Moiety | | |
|---|---|---|
| **Ingredient Name** | **Basis of Strength** | **Strength** |
| **DEXTROMETHORPHAN HYDROBROMIDE** (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 20 mg in 10 mL |
| **GUAIFENESIN** (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ) | GUAIFENESIN | 200 mg in 10 mL |
| **PHENYLEPHRINE HYDROCHLORIDE** (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | PHENYLEPHRINE HYDROCHLORIDE | 10 mg in 10 mL |

| Inactive Ingredients | |
|---|---|
| **Ingredient Name** | **Strength** |
| **ANHYDROUS CITRIC ACID** (UNII: XF417D3PSL) | |
| **EDETATE DISODIUM** (UNII: 7FLD91C86K) | |
| **FD&C RED NO. 40** (UNII: WZB9127XOA) | |
| **GLYCERIN** (UNII: PDC6A3C0OX) | |
| **PROPYLENE GLYCOL** (UNII: 6DC9Q167V3) | |
| **WATER** (UNII: 059QF0KO0R) | |
| **SODIUM BENZOATE** (UNII: OJ245FE5EU) | |
| **SODIUM CITRATE, UNSPECIFIED FORM** (UNII: 1Q73Q2JULR) | |
| **SORBITOL** (UNII: 506T60A25R) | |
| **SUCRALOSE** (UNII: 96K6UQ3ZD4) | |

| Product Characteristics | | | |
|---|---|---|---|
| **Color** | RED | **Score** | |
| **Shape** | | **Size** | |
| **Flavor** | CHERRY | **Imprint Code** | |
| **Contains** | | | |

**Packaging**

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|---|---|---|---|
| **1** | NDC:24385-904-26 | 1 in 1 CARTON | 04/11/2006 | |
| **1** | | 118 mL in 1 BOTTLE; Type 0: Not a Combination Product | | |
| **2** | NDC:24385-904-34 | 1 in 1 CARTON | 03/27/2006 | |
| **2** | | 237 mL in 1 BOTTLE; Type 0: Not a Combination Product | | |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---|---|---|---|
| OTC monograph final | part341 | 03/27/2006 | |

## Labeler - Amerisource Bergen (007914906)

Revised: 11/2019 Amerisource Bergen