

**ACETAMINOPHEN 500MG CAFFEINE 60MG PYRILAMINE MALEATE 15MG-  
acetaminophen 500mg caffeine 60mg pyrilamine maleate 15mg tablet  
WALGREENS CO.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Walgreens Menstrual Complete Pain Relief  
Acetaminophen 500mg, Caffeine 60mg, Pyrilamine Maleate 15mg**

Acetaminophen 500mg

Caffeine 60mg

Pyrilamine maleate 15mg

ammonium hydroxide, croscarmellose sodium, D&C red No. 33 aluminum lake, ethyl cellulose, FD&C blue No. 1 aluminum lake, gelatin, glycerin, hypromellose, iron oxide black, light mineral oil, magnesium stearate, microcrystalline cellulose, povidone, pregelatinised starch, propylene glycol, shellac glaze, sorbitol sorbitan, talc, titanium dioxide, triacetin

- Do not take more than the recommended dose
- Adults and children 12 years and over: take 2 gelcaps with water, repeat every 6 hours, as needed; do not exceed 6 gelcaps per day
- Children under 12 years: consult a doctor

For the temporary relief of these symptoms associated with menstrual periods:

Cramps, bloating water-weight gain, headache, backache, muscle aches, fatigue.

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take more than 6 gelcaps in 24 hours, which is the maximum daily amount for this product, with other drugs contains acetaminophen, 3 or more alcoholic drinks every day while using this product.

**Allergy alert:** Acetaminophen may cause severe skin reactions or severe allergic reactions. Symptoms may include: skin reddening, blisters, rash, hives, facial swelling, asthma (wheezing), shock. If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Do not use** with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist, if you have ever had an allergic reaction to this product or any of its ingredients.

**Ask a doctor before use** if you have liver disease, glaucoma, difficulty in urination due to enlargement of the prostate gland, a breathing problem such as emphysema or chronic bronchitis.

**Ask a doctor or pharmacist before use** if you are taking the blood thinning drug warfarin, taking sedatives or tranquilizers.

**When using this product**, you may get drowsy, avoid alcoholic drinks, excitability

may occur, especially in children, alcohol, sedatives, and tranquilizers may increase drowsiness, be careful when driving a motor vehicle or operating machinery, limit the use of caffeine-containing medications, foods, or beverages because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat. The recommended dose of this product contains about as much caffeine as a cup of coffee.

**Stop use and ask a doctor** if new symptoms occur, redness or swelling is present, pain gets worse or lasts for more than 10 days, fever gets worse or lasts for more than 3 days.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children**

**Overdose Warning:** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Pain reliever

Nighttime sleep-aid

Store at room temperature between 20-25 °C (68-77 °F)

**Other Information** Store at room temperature 68-77°F (20-25°C).

**Inactive ingredients** aluminum hydroxide, croscarmellose sodium, DcRed no. 33 aluminum lake, ethylcellulose, FD&C colorless powder, polyethylene glycol, polyethylene glycol, sorbitol sorbitan, tac, titanium dioxide, triacetin.

**Questions or comments?** 1-888-577-8033

**Directions** ■ do not take more than the recommended dose ■ adults and children 12 years and older: ■ take 2 gelscaps with water ■ repeat every 6 hours as needed ■ do not exceed 6 gelscaps per day ■ children under 12 years: consult a doctor.

**Drug Facts (continued)**

**Do not use** ■ with any other drug containing acetaminophen (prescription or over-the-counter). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland ■ a breathing problem such as emphysema or chronic bronchitis.

**Warnings** ■ Severe liver damage may occur if you take more than 6 gelscaps in 24 hours, which is the maximum daily amount for this product. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. ■ Allergy alert: acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include: ■ skin redness ■ hives ■ rash ■ facial swelling ■ asthma (wheezing) ■ shock ■ If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Drug Facts (continued)**

**Do not use** ■ with any other drug containing acetaminophen (prescription or over-the-counter). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland ■ a breathing problem such as emphysema or chronic bronchitis.

**Warnings** ■ Severe liver damage may occur if you take more than 6 gelscaps in 24 hours, which is the maximum daily amount for this product. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. ■ Allergy alert: acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include: ■ skin redness ■ hives ■ rash ■ facial swelling ■ asthma (wheezing) ■ shock ■ If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Drug Facts (continued)**

**Do not use** ■ with any other drug containing acetaminophen (prescription or over-the-counter). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland ■ a breathing problem such as emphysema or chronic bronchitis.

**Warnings** ■ Severe liver damage may occur if you take more than 6 gelscaps in 24 hours, which is the maximum daily amount for this product. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. ■ Allergy alert: acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include: ■ skin redness ■ hives ■ rash ■ facial swelling ■ asthma (wheezing) ■ shock ■ If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Drug Facts (continued)**

**Do not use** ■ with any other drug containing acetaminophen (prescription or over-the-counter). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland ■ a breathing problem such as emphysema or chronic bronchitis.

**Warnings** ■ Severe liver damage may occur if you take more than 6 gelscaps in 24 hours, which is the maximum daily amount for this product. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. ■ Allergy alert: acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include: ■ skin redness ■ hives ■ rash ■ facial swelling ■ asthma (wheezing) ■ shock ■ If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Drug Facts (continued)**

**Do not use** ■ with any other drug containing acetaminophen (prescription or over-the-counter). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland ■ a breathing problem such as emphysema or chronic bronchitis.

**Warnings** ■ Severe liver damage may occur if you take more than 6 gelscaps in 24 hours, which is the maximum daily amount for this product. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. ■ Allergy alert: acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include: ■ skin redness ■ hives ■ rash ■ facial swelling ■ asthma (wheezing) ■ shock ■ If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Drug Facts (continued)**

**Do not use** ■ with any other drug containing acetaminophen (prescription or over-the-counter). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland ■ a breathing problem such as emphysema or chronic bronchitis.

**Warnings** ■ Severe liver damage may occur if you take more than 6 gelscaps in 24 hours, which is the maximum daily amount for this product. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. ■ Allergy alert: acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include: ■ skin redness ■ hives ■ rash ■ facial swelling ■ asthma (wheezing) ■ shock ■ If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Drug Facts (continued)**

**Do not use** ■ with any other drug containing acetaminophen (prescription or over-the-counter). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland ■ a breathing problem such as emphysema or chronic bronchitis.

**Warnings** ■ Severe liver damage may occur if you take more than 6 gelscaps in 24 hours, which is the maximum daily amount for this product. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. ■ Allergy alert: acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include: ■ skin redness ■ hives ■ rash ■ facial swelling ■ asthma (wheezing) ■ shock ■ If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Drug Facts (continued)**

**Do not use** ■ with any other drug containing acetaminophen (prescription or over-the-counter). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland ■ a breathing problem such as emphysema or chronic bronchitis.

**Warnings** ■ Severe liver damage may occur if you take more than 6 gelscaps in 24 hours, which is the maximum daily amount for this product. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. ■ Allergy alert: acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include: ■ skin redness ■ hives ■ rash ■ facial swelling ■ asthma (wheezing) ■ shock ■ If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Drug Facts (continued)**

**Do not use** ■ with any other drug containing acetaminophen (prescription or over-the-counter). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland ■ a breathing problem such as emphysema or chronic bronchitis.

**Warnings** ■ Severe liver damage may occur if you take more than 6 gelscaps in 24 hours, which is the maximum daily amount for this product. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. ■ Allergy alert: acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include: ■ skin redness ■ hives ■ rash ■ facial swelling ■ asthma (wheezing) ■ shock ■ If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Drug Facts (continued)**

**Do not use** ■ with any other drug containing acetaminophen (prescription or over-the-counter). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland ■ a breathing problem such as emphysema or chronic bronchitis.

**Warnings** ■ Severe liver damage may occur if you take more than 6 gelscaps in 24 hours, which is the maximum daily amount for this product. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. ■ Allergy alert: acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include: ■ skin redness ■ hives ■ rash ■ facial swelling ■ asthma (wheezing) ■ shock ■ If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Drug Facts (continued)**

**Do not use** ■ with any other drug containing acetaminophen (prescription or over-the-counter). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland ■ a breathing problem such as emphysema or chronic bronchitis.

**Warnings** ■ Severe liver damage may occur if you take more than 6 gelscaps in 24 hours, which is the maximum daily amount for this product. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. ■ Allergy alert: acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include: ■ skin redness ■ hives ■ rash ■ facial swelling ■ asthma (wheezing) ■ shock ■ If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Drug Facts (continued)**

**Do not use** ■ with any other drug containing acetaminophen (prescription or over-the-counter). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland ■ a breathing problem such as emphysema or chronic bronchitis.

**Warnings** ■ Severe liver damage may occur if you take more than 6 gelscaps in 24 hours, which is the maximum daily amount for this product. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. ■ Allergy alert: acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include: ■ skin redness ■ hives ■ rash ■ facial swelling ■ asthma (wheezing) ■ shock ■ If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Drug Facts (continued)**

**Do not use** ■ with any other drug containing acetaminophen (prescription or over-the-counter). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland ■ a breathing problem such as emphysema or chronic bronchitis.

**Warnings** ■ Severe liver damage may occur if you take more than 6 gelscaps in 24 hours, which is the maximum daily amount for this product. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. ■ Allergy alert: acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include: ■ skin redness ■ hives ■ rash ■ facial swelling ■ asthma (wheezing) ■ shock ■ If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Drug Facts (continued)**

**Do not use** ■ with any other drug containing acetaminophen (prescription or over-the-counter). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland ■ a breathing problem such as emphysema or chronic bronchitis.

**Warnings** ■ Severe liver damage may occur if you take more than 6 gelscaps in 24 hours, which is the maximum daily amount for this product. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. ■ Allergy alert: acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include: ■ skin redness ■ hives ■ rash ■ facial swelling ■ asthma (wheezing) ■ shock ■ If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Drug Facts (continued)**

**Do not use** ■ with any other drug containing acetaminophen (prescription or over-the-counter). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland ■ a breathing problem such as emphysema or chronic bronchitis.

**Warnings** ■ Severe liver damage may occur if you take more than 6 gelscaps in 24 hours, which is the maximum daily amount for this product. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. ■ Allergy alert: acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include: ■ skin redness ■ hives ■ rash ■ facial swelling ■ asthma (wheezing) ■ shock ■ If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**TAMPER EVIDENT: Do not use if package is opened or if blister unit is torn, broken or shows any sign of tampering.**

DISTRIBUTED BY: WALGREENS CO.  
20 WALTON RD., DEERFIELD, IL 60015

**100% SATISFACTION GUARANTEED**

WALGREENS  
walmart.com ©2016 Walgreen Co.  
MADE IN U.S.A.

ITEM 542303 W10058-0519-L

1 917420839

ORIG0419-F

**Drug Facts**

**Active ingredients (in each gelcap)**

Acetaminophen 500 mg  
Caffeine 60 mg  
Pyrilamine maleate 15 mg

**Purpose**

Pain reliever

**Uses** for the temporary relief of these symptoms associated with menstrual periods: ■ cramps ■ bloating ■ water-weight gain ■ headache ■ backache ■ muscle aches ■ fatigue

**Walgreens** Compare to Midol® Complete active ingredients!<sup>††</sup>

NDC 0363-2310-24

**NEW**

# Menstrual Complete Pain Relief

ACETAMINOPHEN 500 mg / PAIN RELIEVER  
CAFFEINE 60 mg / DIURETIC  
PYRILAMINE MALEATE 15 mg / ANTIHISTAMINE

**ASPIRIN FREE GELCAPS**

- Relieves cramps, bloating, fatigue, headaches & backaches.

**24 GELCAPS** ACTUAL SIZE

**RECOMMENDED** Health excellence you only can't see.

**PHARMACY**

**Walgreens** Health since 1929®

Walgreens Pharmacist Survey  
†† This product is not manufactured or distributed by Bayer Healthcare LLC, owner of the registered trademark Midol® Complete.  
REI-0320 CT0363231024

LOT: \_\_\_\_\_  
EXP. DT: \_\_\_\_\_

# 15MG

acetaminophen 500mg caffeine 60mg pyrilamine maleate 15mg tablet

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-2310
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>CAFFEINE</b> (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	60 mg
<b>PYRILAMINE MALEATE</b> (UNII: R35D29L3ZA) (PYRILAMINE - UNII:HPE317O9TL)	PYRILAMINE MALEATE	15 mg
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>D&amp;C RED NO. 33</b> (UNII: 9DBA0SBB0L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>FERROSFERRIC OXIDE</b> (UNII: XM0M87F357)	
<b>LIGHT MINERAL OIL</b> (UNII: N6K5787QVP)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>HYDROXYETHYL CELLULOSE, UNSPECIFIED</b> (UNII: T4V6TWG28D)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>SHELLAC</b> (UNII: 46N107B710)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	

## Product Characteristics

<b>Color</b>	blue (DARK BLUE and LIGHT BLUE)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	S79
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

<b>1</b>	NDC:0363-2310-24	24 in 1 PACKAGE; Type 0: Not a Combination Product	07/05/2019	
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>		<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph not final	part343		07/05/2019	

**Labeler - WALGREENS CO. (008965063)**

Revised: 7/2023

WALGREENS CO.