

MOTION SICKNESS- meclizine hcl tablet
H E B

HEB 44-403

Active ingredient (in each tablet)

Meclizine HCl 25 mg

Purpose

Antiemetic

Uses

for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness

Warnings

Do not use

for children under 12 years of age unless directed by a doctor.

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- glaucoma
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- take the first dose 1/2 to 1 hour before starting activity
- adults and children 12 years and over: 1 to 2 tablets once daily, or as directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from heat and humidity
- use by expiration date on package

Inactive ingredients

colloidal silicon dioxide, D&C yellow #10 aluminum lake, lactose monohydrate, magnesium stearate, pregelatinized starch

Questions or comments?

1-800-426-9391

Principal Display Panel

H-E-B®

NDC 37808-903-12

**Compare to
Dramamine®**

Less Drowsy
active ingredient*

**Motion
Sickness**

Meclizine HCl 25 mg

Antiemetic

Helps prevent Nausea
& Dizziness due to
Motion Sickness for
up to 24 hours

- For Ages 12 and Over
- 24 Hours of Protection
- Less Drowsy Formula

Actual
Size

100 TABLETS

TAMPER EVIDENT: DO NOT USE IF IMPRINTED

SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

*This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Dramamine® Less Drowsy. 50844 ORG042340312

100%
GUARANTEE
promise

MADE WITH PRIDE AND CARE FOR H-E-B®, SAN ANTONIO, TX 78204

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts	Purpose	Warnings	100% GUARANTEE
Active ingredient (in each tablet) Meclizine HCl 25 mg.....Antiemetic	Uses for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness	Do not use for children under 12 years of age unless directed by a doctor. Ask a doctor before use if you have <ul style="list-style-type: none"> ■ difficulty in urination due to enlargement of the prostate gland ■ glaucoma ■ a breathing problem such as emphysema or chronic bronchitis 	<p style="font-size: x-small;">*This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Dramamine® Less Drowsy. 50844 ORG042340312 12110-1903</p> <p style="font-size: x-small;">MADE WITH PRIDE AND CARE FOR H-E-B® SAN ANTONIO, TX 78204</p>

STOP PEEING

Questions or comments? 1-800-426-9391	Inactive ingredients colloidal silicon dioxide, D&C yellow #10 aluminum lake, lactose monohydrate, magnesium stearate, pregelatinized starch	Other information <ul style="list-style-type: none"> ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ protect from heat and humidity ■ use by expiration date on package 	Directions <ul style="list-style-type: none"> ■ do not take more than directed ■ take first dose ½ to 1 hour before starting activity ■ adults and children 12 years and over: 1 to 2 tablets once daily, or as directed by a doctor
---	--	---	---

Drug Facts (continued)

When using this product

- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

HEB 44-403A

MOTION SICKNESS

meclizine hcl tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37808-903
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MECLIZINE HYDROCHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570)	MECLIZINE HYDROCHLORIDE	25 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
STARCH, CORN (UNII: O8232NY3S)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	9mm
Flavor		Imprint Code	44;403
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-903-12	100 in 1 BOTTLE; Type 0: Not a Combination Product	06/24/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M009	06/24/2019	

Labeler - H E B (007924756)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(37808-903)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(37808-903) , pack(37808-903)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(37808-903)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(37808-903)

Revised: 8/2023

H E B