

**ACETAMINOPHEN 250MG ASPIRIN 250MG CAFFEINE 65MG- acetaminophen 250mg aspirin 250mg caffeine 65mg tablet
WALGREENS CO.**

**Walgreens Headache Relief Geltabs
Acetaminophen 250mg, Aspirin 250mg, Caffeine 65mg**

Drug Facts

Active ingredients (in each geltab)

Acetaminophen 250 mg

Aspirin 250 mg (NSAID*)

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever

Pain reliever

Pain reliever aid

Uses

Temporarily relieves minor aches and pains due to:

• headache • a cold • arthritis • muscular aches • toothache • premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 geltabs in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDS (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasional rapid heartbeat.

Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer

- with any other drug containing acetaminophen (prescription or nonprescription). if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if • you have liver disease • stomach bleeding warning applies to you • you have a history of stomach problems such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma

Ask a doctor or pharmacist before use if you are taking

- a prescription drug for diabetes, gout, or arthritis
- any other drug, or are under a doctor's care for any serious condition

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- painful area is red or swollen
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- any new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed (see overdose warning)**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 geltabs every 6 hours; not more than 8 geltabs in 24 hours
- children under 12 years: ask a doctor

Other information

- store at room temperature between 20-25°C(68-77°F)
- avoid excessive heat, cold and humidity

Inactive ingredients

corn starch, D&C yellow #10 aluminum lake, edible black ink, ethyl cellulose, FD&C blue #1 aluminum lake, FD&C red #40 aluminum lake, gelatin, glycerin, hypromellose, isopropyl alcohol, lactose monohydrate, light mineral oil, microcrystalline cellulose, pregelatinized potato starch, purified water, pvpk-30, sorbitol sorbitan solution, stearic acid, talc, titanium dioxide, triacetin

Questions or comments?

1-888-577-8033 Monday-Friday 8am-4pm EST

Compare to the active ingredients in Excedrin® Extra Strength††

Extra Strength

READ AND KEEP OUTER CARTON FOR COMPLETE PRODUCT WARNINGS AND INFORMATION

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

†Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

††This product is not manufactured or distributed by GSK Consumer Healthcare SARL, owner of the registered trademark Excedrin®.

DISTRIBUTED BY: **WALGREEN CO.**
DEERFIELD, IL 60015

100% SATISFACTION GUARANTEED
walgreens.com 1-800-925-4733

©2021 Walgreen Co.

PRODUCT OF UNITED ARAB EMIRATES

Packaging

[illegible]



Compare to the active ingredients in Excedrin® Extra Strength™

Headache Relief

ACETAMINOPHEN / ASPIRIN (NSAID) / CAFFEINE / PAIN RELIEVER / PAIN RELIEVER AID

Extra Strength

80
GELTABS

ACTUAL SIZE

W3ORG1221-F

Drug Facts

Active ingredients (in each geltab)	Purposes
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID)*	Pain reliever
Caffeine 65 mg	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Uses

Temporarily relieves minor aches and pains due to:

- headache
- a cold
- arthritis
- muscular aches
- toothache
- premenstrual and menstrual cramps

Warnings

Ray's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, it changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Ray's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause skin reactions. Symptoms may include:

- skin redness
- hives
- itch
- if a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 geltabs in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasional rapid heartbeat.

Drug Facts (continued)

Do not use if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/reducer with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have liver disease, stomach bleeding warning applies to you, you have a history of stomach problems such as heartburn, you have high blood pressure, heart disease, liver cirrhosis, or kidney disease, you are taking a diuretic, you have asthma.

Ask a doctor or pharmacist before use if you are taking a prescription drug for diabetes, gout, or arthritis, any other drug, or are under a doctor's care for any serious condition.

Stop use and ask a doctor if an allergic reaction occurs. Seek medical help right away, you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better, ringing in the ears or loss of hearing occurs, painful area is red or swollen, pain gets worse or lasts for more than 10 days, fever gets worse or lasts for more than 3 days, any new symptoms occur. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed (see overdose warning)
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 geltabs every 6 hours; not more than 8 geltabs in 24 hours
- children under 12 years: ask a doctor

120 GELTABS



Compare to the active ingredients in Excedrin® Extra Strength™

Headache Relief

ACETAMINOPHEN / ASPIRIN (NSAID) / CAFFEINE / PAIN RELIEVER / PAIN RELIEVER AID

Extra Strength

120
GELTABS

Drug Facts

Active ingredients (in each geltab)	Purposes
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID)*	Pain reliever
Caffeine 65 mg	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Uses

Temporarily relieves minor aches and pains due to:

- headache
- a cold
- arthritis
- muscular aches
- toothache
- premenstrual and menstrual cramps

Warnings

Ray's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, it changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Ray's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause skin reactions. Symptoms may include:

- skin redness
- hives
- itch
- if a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 geltabs in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasional rapid heartbeat.

Drug Facts (continued)

Do not use if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/reducer with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have liver disease, stomach bleeding warning applies to you, you have a history of stomach problems such as heartburn, you have high blood pressure, heart disease, liver cirrhosis, or kidney disease, you are taking a diuretic, you have asthma.

Ask a doctor or pharmacist before use if you are taking a prescription drug for diabetes, gout, or arthritis, any other drug, or are under a doctor's care for any serious condition.

Stop use and ask a doctor if an allergic reaction occurs. Seek medical help right away, you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better, ringing in the ears or loss of hearing occurs, painful area is red or swollen, pain gets worse or lasts for more than 10 days, fever gets worse or lasts for more than 3 days, any new symptoms occur. These could be signs of a serious condition.


If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed (see overdose warning)
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 geltabs every 6 hours; not more than 8 geltabs in 24 hours
- children under 12 years: ask a doctor



DRUG FACTS

READ AND KEEP OUTER CARTON FOR COMPLETE PRODUCT WARNINGS AND INFORMATION
TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each gellab)	Purposes
Acetaminophen 250 mg.....	Pain reliever
Aspirin 250 mg (NSAID)*.....	Pain reliever
Caffeine 65 mg.....	Pain reliever aid
*nonsteroidal anti-inflammatory drug	

Uses

Temporarily relieves minor aches and pains due to:
 ■ headache ■ a cold ■ arthritis ■ muscular aches ■ toothache
 ■ premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause skin reactions.

Symptoms may include:

■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

■ more than 8 gellabs in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen

■ 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
 ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

■ have 3 or more alcoholic drinks every day while using this product
 ■ take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasional rapid heartbeat.

Drug Facts (continued)

Do not use ■ if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if ■ you have liver disease ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic
 ■ you have asthma

Ask a doctor or pharmacist before use if you are taking

■ a prescription drug for diabetes, gout, or arthritis
 ■ any other drug, or are under a doctor's care for any serious condition

Stop use and ask a doctor if

■ an allergic reaction occurs. Seek medical help right away.
 ■ you experience any of the following signs of stomach bleeding:
 ■ feel faint ■ vomit blood ■ have bloody or black stools
 ■ have stomach pain that does not get better
 ■ ringing in the ears or loss of hearing occurs
 ■ painful area is red or swollen
 ■ pain gets worse or lasts for more than 10 days
 ■ fever gets worse or lasts for more than 3 days
 ■ any new symptoms occur
 These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

■ do not use more than directed (see overdose warning)
 ■ drink a full glass of water with each dose
 ■ adults and children 12 years and over: take 2 gellabs every 6 hours; not more than 8 gellabs in 24 hours
 ■ children under 12 years: ask a doctor

Drug Facts (continued)

Other information

■ store at room temperature between 20-25°C (68-77°F)
 ■ avoid excessive heat, cold and humidity

Inactive ingredients

corn starch, D&C yellow #10 aluminum lake, edible black ink, ethyl cellulose, FD&C blue #1 aluminum lake, FD&C red #40 aluminum lake, gelatin, glycerin, hypromellose, isopropyl alcohol, lactose

lake, gelatin, glycerin, hypromellose, isopropyl alcohol, lactose monohydrate, light mineral oil, microcrystalline cellulose, pregelatinized potato starch, purified water, pvpk-30, sorbitol sorbitan solution, stearic acid, talc, titanium dioxide, triacetin

Questions or comments?

1-888-577-8033 Monday-Friday 8am-4pm EST

ACETAMINOPHEN 250MG ASPIRIN 250MG CAFFEINE 65MG

acetaminophen 250mg aspirin 250mg caffeine 65mg tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-1320
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
ETHYLCELLULOSE, UNSPECIFIED (UNII: 7Z8S9VYZ4B)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
FD&C RED NO. 40 ALUMINUM LAKE (UNII: 6T47AS764T)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
STARCH, POTATO (UNII: 8I089SAH3T)	
WATER (UNII: 059QF0KO0R)	
POVIDONE K30 (UNII: U725QWY32X)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	green (Dark Green - White)	Score	no score
Shape	ROUND	Size	13mm
Flavor		Imprint Code	S77
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-1320-12	120 in 1 BOTTLE; Type 0: Not a Combination Product	09/24/2019	
2	NDC:0363-1320-80	80 in 1 BOTTLE; Type 0: Not a Combination Product	09/24/2019	
3	NDC:0363-1320-23	20 in 1 BOTTLE; Type 0: Not a Combination Product	09/24/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	09/24/2019	

Labeler -
WALGREENS CO. (008965063)