COLD AND FLU RELIEF- acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule, liquid filled Chain Drug Consortium, LLC

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg

Dextromethoprhan HBr 10 mg

Phenlyephrine HCl 5 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - fever
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks everyday while using this product

Allery alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if the user has

- liver disease
- diabetes
- heart disease
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product,

do not exceed recommended dosage.

Stop use and ask a doctor if

- pain, cough or nasal congestion gets worse or lasts more than 7 days
- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs, with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)
- do not take more than 4 doses in 24-hour
- adults and children 12 years and over: take 2 softgels with water every 4 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- When using Daytime and Nighttime products, carefully read each label to ensure correct dosing

Other information

- store between 15-30°C (59-86°)
- avoid excessive heat

Inactive ingredients

FD&C yellow #6, FD&C red #40, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol white ink

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Principal Display Panel

*COMPARE TO HE ACTIVE INGREDIENTS IN VICKS® DAYQUIL® COLD & FLU LIQUICAPS®

Daytime, Multi-Symptom

Cold & Flu Relief

Acetaminophen 325 mg

Pain reliever/Fever reducer

DEXTROMETHORPHAN HBr 10 mg

Cough suppressant

PHENYLEPHRINE HCI 5 mg

Nasal Decongestant

Relieves: Aches •Fever •Sore Throat Cough • Nasal Congestion

Non-drowsy, Alcohol-free, Antihistamine-free

Softgels**

(**Liquid-filled Capsules)

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOW ANY SIGNS OR TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® DayQuil® and LiquiCaps® are registered trademarks of The Procter & Gamble Company.

Distributed by: Pharmacy Value Alliance, LLC

407 East Lancaster Avenue,

Wayne, PA 19087

Package Label



When using this product, do not exceed recommended dosage.	TS3 M92-MA9 vsbing-vsbrom 8595-537-778-1 lls0	
isk a doctor or pharmacist before use if you are taking the blood binning drug wartarin.	Questions or comments?	
persistent or chronic cough such as occurs with smoking, asthma, or emphysema cough that occurs with too much phlegm (mucus)	Inscrive ingredients FD&C red #40, FD&C yellow #6, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purfiled water, sorbitan, sorbitol, white ink	
is to cortor before use it you have m liver disease m diabetes heart disease m thyroid disease m high blood pressure trouble uninating due to an enlarged prostate gland	Other information arote between 15-30'C (59-86'F) arote between 12-30'C (59-86'F)	
with any other drug containing acetaminophen (preacription or nonpreacription). If you are not sure whether a drug contrains acetaminophen, ask a doctor or pharmacist. If you are now taking a preacription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or (certain drugs for depression, psychiatric, or emotional conditions, or do not know if your prescription drug contains an MAOI, ask a doctor or do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	Directions = do not take more than directed (see Overdose warning) = do not take more than 4 doses in 24 hours = do not take more than 4 doses in 24 hours = adults and children 12 years and over: take 2 softgels with water every 4 hours = swallow whole; do not crush, chew, or dissolve = swallow whole; do not crush, chew, or dissolve = children under 12 years: do not use = children under 12 years: do not use = children using other Daytime or Nighttime products, carefully read each = able to ensure correct dosing = able to ensure correct dosing	
ney include: sin reddening bilisters resh is skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 ays, is accompanied or followed by fever, headache, rash, nausea, or omiting, consult a doctor promptly. Jo not use	If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Oulck medical attention is critical for Center (1-800-222-1222) right away. Oulck medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
Warmings Jver waming: This product contains acetaminophen. Severe liver damage nay occur if you take: more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product 3 or more alcoholic drinks every day while using this product 3 or more alcoholic drinks every day while using this product 10 or more alcoholic drinks every day while using this product 10 or more alcoholic drinks every day while using this product 10 or more alcoholic drinks every day while using this product 10 or more alcoholic drinks every day while using this product 10 or more alcoholic drinks are severe and the skin reactions. Symptoms	Stop use and ask a doctor if a pain, cough, or nasal congestion gets worse or lasts more than 7 days a nervousness, diszinees, or sleeplessness occur a fever gets worse or lasts more than 3 days a fever gets worse or lasts more than 7 days a fever gets worse or lasts more than a days a fever	
Drug Facts (continued)	Drug Facts (continued)	

14

when using this product, do not exceed recommended dosage.

LC2 MUC-IMMC APRILL-APRILION CCCC-CCI-LID-L HPD



COLD AND FLU RELIEF acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule, liquid filled **Product Information** HUMAN OTC DRUG NDC:68016-797 **Product Type** Item Code (Source) **Route of Administration** ORAL **Active Ingredient/Active Moiety Ingredient Name Basis of Strength** Strength ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII: 36209ITL9D) ACETAMINOPHEN 325 mg DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) DEXTROMETHORPHAN 10 mg (DEXTROMETHORPHAN - UNII:7355X3ROTS) **HYDROBROMIDE** PHENYLEPHRINE HYDROCHLORIDE (UNII: 04/A59TNS)) (PHENYLEPHRINE -PHENYLEPHRINE 5 mg UNII:1WS297W6MV) HYDROCHLORIDE

Inactive Ingredients				
	Ingredien	t Name		Strength
FD&C YELLOW NO. 6 (UNII: I	H77VEI93A8)			
GLYCERIN (UNII: PDC6A3C00)	()			
PROPYLENE GLYCOL (UNII: 6	DC9Q167V3)			
WATER (UNII: 059QF0KO0R)				
LECITHIN, SOYBEAN (UNII: 1	DI56QDM62)			
FD&C RED NO. 40 (UNII: WZ E	39127XOA)			
GELATIN (UNII: 2G86QN327L)				
POLYETHYLENE GLYCOL, U	NSPECIFIED (UNII: 3	WJQ0SDW1A)		
SORBITOL (UNII: 506T60A25R)			
POVIDONE (UNII: FZ989GH94	E)			
SORBITAN (UNII: 6092ICV9RU)			
LIGHT MINERAL OIL (UNII: NO	K5787QVP)			
MANNITOL (UNII: 30WL53L364	A)			
Product Characterist				
Color		Score	no 4	score
	orange			
Shape	OVAL	Size	20n	nm

Fla	vor Imprint Code			PC9	
Co	ontains				
Pa	ackaging				
#	ltem Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:68016-797- 16	16 in 1 CARTON	09/02/2022		
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product			
Μ	arketing I	nformation			
	_	Application Number or Monograph	Marketing Start	Marketing End	
	Marketing Category	Citation	Date	Date	

Labeler - Chain Drug Consortium, LLC (101668460)

Revised: 3/2024

Chain Drug Consortium, LLC