

**ASPIRIN LOW DOSE- aspirin tablet, coated**  
**SAVE-A-LOT FOOD STORES, LTD.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Being Well 44-600A**

**Active ingredient (in each tablet)**

Aspirin 81 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever

**Uses**

for the temporary relief of minor aches and pains or as recommended by your doctor

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- have 3 or more alcoholic drinks every day while using this product
- have had stomach ulcers or bleeding problems
- take more or for a longer time than directed

**Do not use**

if you are allergic to aspirin or any other pain reliever/fever reducer.

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you have asthma

- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - vomit blood
  - have bloody or black stools
  - feel faint
  - have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- new symptoms occur
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts more than 10 days
- redness or swelling is present

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

***Directions***

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: ask a doctor

***Other information***

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

***Inactive ingredients***

corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

***Questions or comments?***

**1-800-426-9391**

***Principal Display Panel***

No print/No varnish  
Lot & Exp date

COMPARE TO THE ACTIVE INGREDIENT IN BAYER® LOW DOSE ASPIRIN†

# Aspirin LOW DOSE

(NSAID), 81 mg

ACTUAL SIZE

SAFETY COATED

ASPIRIN  
REGIMEN

120

Delayed Release  
TABLETS

†This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Bayer® Low Dose Aspirin.

DISTRIBUTED BY:  
SAVE-A-LOT FOOD STORES, LTD.  
EARTH CITY, MO 63045  
savealot.com

B-1811-600A-32-R  
REV0915060032

Drug Facts (continued)

uniform child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

**Inactive ingredients** corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

**Questions or comments?**  
1-800-426-9391

50844 REV0915060032

Drug Facts (continued)

- have had stomach ulcers or bleeding problems
- take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.

**Ask a doctor before use if**

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you have asthma
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are taking a prescription drug for**

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - vomit blood
  - have bloody or black stools
  - feel faint
- have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- new symptoms occur
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts more than 10 days
- redness or swelling is present

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the

Drug Facts

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

**Active ingredient (in each tablet)**  
Aspirin 81 mg (NSAID)\* ..... Pain reliever  
\*nonsteroidal anti-inflammatory drug

**Uses** for the temporary relief of minor aches and pains or as recommended by your doctor

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:  

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- have 3 or more alcoholic drinks every day while using this product

Drug Facts (continued)

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

**Active ingredient (in each tablet)**  
Aspirin 81 mg (NSAID)\* ..... Pain reliever  
\*nonsteroidal anti-inflammatory drug

**Uses** for the temporary relief of minor aches and pains or as recommended by your doctor

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:  

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- have 3 or more alcoholic drinks every day while using this product

Being Well 44-600A

COMPARE TO THE ACTIVE INGREDIENT IN BAYER® LOW DOSE ASPIRIN†

**Aspirin LOW DOSE**  
(NSAID), 81 mg

PAIN RELIEVER

SAFETY COATED

Being Well®

- Adult Low Strength Aspirin Regimen
- Enteric Safety Coated

120

Delayed Release

**TABLETS**

**TAMPER EVIDENT: DO NOT USE IF  
IMPRINTED SAFETY SEAL UNDER  
CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Bayer® Low Dose Aspirin.

50844 REV0915C60032

**DISTRIBUTED BY:  
SAVE-A-LOT FOOD STORES, LTD.  
EARTH CITY, MO 63045  
savealot.com**

**Save a Lot  
promise  
100% MONEY BACK  
GUARANTEE**

<b>ASPIRIN LOW DOSE</b>			
aspirin tablet, coated			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:46994-600
<b>Route of Administration</b>	ORAL		
<b>Active Ingredient/Active Moiety</b>			
	<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
	ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg
<b>Inactive Ingredients</b>			
	<b>Ingredient Name</b>		<b>Strength</b>
	STARCH, CORN (UNII: O8232NY3SJ)		
	D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)		
	HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)		
	METHACRYLIC ACID (UNII: 1CS02G8656)		
	MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)		
	POLYDEXTROSE (UNII: VH2XOU12IE)		
	POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)		
	SHELLAC (UNII: 46N107B71O)		
	SODIUM BICARBONATE (UNII: 8MDF5V39QO)		
	SODIUM LAURYL SULFATE (UNII: 368GB5141J)		
	TALC (UNII: 7SEV7J4R1U)		
	TITANIUM DIOXIDE (UNII: 15FIX9V2JP)		
	TRIETHYL CITRATE (UNII: 8Z96QXD6UM)		

FD&C YELLOW NO. 6 (UNI: H77VEI93A8)	
SILICON DIOXIDE (UNI: ETJ7Z6XBU4)	
TRIACETIN (UNI: XHX3C3X673)	

Product Characteristics			
Color	YELLOW	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	L
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46994-600-32	1 in 1 CARTON	05/01/2011	
1		120 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part343	05/01/2011	

**Labeler** - SAVE-A-LOT FOOD STORES, LTD. (095484572)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(46994-600)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	PACK(46994-600)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	MANUFACTURE(46994-600) , PACK(46994-600)