

RANITIDINE- ranitidine tablet, film coated
Walgreen Company

ACTIVE INGREDIENT (IN EACH TABLET)

Ranitidine 75 mg (as ranitidine hydrochloride, USP 84 mg)

PURPOSE

Acid reducer

USES

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

WARNINGS

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor
- with other acid reducers

Ask a doctor before use if you have

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Stop use and ask a doctor if

- your heartburn continues or worsens
- need to take this product for more than 14 days

If pregnant or breast-feeding

Ask a health professional before use.

Keep Out of Reach of Children

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

DIRECTIONS

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water

- to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
- can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

OTHER INFORMATION

- **TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.**
- store at 20° - 25° C (68° - 77° F)
- avoid excessive heat or humidity
- this product is sodium and sugar free

INACTIVE INGREDIENTS

colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, talc, titanium dioxide

QUESTIONS?

Call **1-800-406-7984**

PRINCIPAL DISPLAY PANEL

Well at *Walgreens*

NDC 0363-0352-30

Wal-Zan[®] 75

Ranitidine Tablets, USP 75 mg

Acid Reducer

- **Prevents & relieves heartburn associated with acid indigestion & sour stomach**

30 TABLETS

(30 DOSES)

Compare to Zantac 75[®] active ingredient^{‡‡}

DISTRIBUTED BY: WALGREEN CO.

5104612/0913

READ THE DIRECTIONS, CONSUMER INFORMATION LEAFLET AND WARNINGS BEFORE USE. KEEP THE CARTON. IT CONTAINS IMPORTANT INFORMATION.

*Walgreens Pharmacist Survey Study, November 2012.
 All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Zantac 75.

DISTRIBUTED BY: **WALGREEN CO.** ITEM 319464
 200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
 walgreens.com
 ©2013 Walgreen Co.
MADE IN INDIA



Walgreens
 Way to Well
 Contribution
 Support local preventive
 wellness with Walgreens. With
 each purchase of Walgreens
 Brand Health & Wellness
 Product, it goes to the
 Walgreens Way to Well Fund?
 up to \$3 million annually. Visit
 Walgreens.com/waytowell

Drug Facts (continued)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

Purpose
 (in each tablet)
 Active ingredient
 Ranitidine 75 mg.....Acid reducer
 (as ranitidine hydrochloride, USP 64 mg)

Uses
 relieves heartburn associated with acid indigestion and sour stomach
 prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings
 Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers
 Do not use
 if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 with other acid reducers
 Ask a doctor before use if you have
 frequent chest pain
 frequent wheezing, particularly with heartburn
 unexplained weight loss
 nausea or vomiting
 stomach pain
 had heartburn over 3 months. This may be a sign of a more serious condition
 heartburn with lightheadedness, sweating or dizziness
 chest pain or shoulder pain with stress or fatigue
 neck or shoulders, or lightheadedness

Directions
 adults and children 12 years and over:
 to relieve symptoms, swallow 1 tablet with a glass of water
 to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn
 can be used up to twice daily (do not take more than 2 tablets in 24 hours)
 children under 12 years: ask a doctor

Other information
 TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOLIO UNDER CAP IS OPEN OR TORN.
 store at 20° - 25° C (68° - 77° F)
 avoid excessive heat or humidity
 this product is sodium and sugar free

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, talc, titanium dioxide

Questions? call 1-800-406-7984

Expiration Date:
 Batch No.

NON VARNISH

ORG0913-F



Actual Size





WALGREENS PHARMACIST RECOMMENDED*

Wal-Zan® 75
 Ranitidine Tablets, USP 75 mg
 Acid Reducer

- Prevents & relieves heartburn associated with acid indigestion & sour stomach

30



NDC 0368-0362-30



RANITIDINE

ranitidine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0352
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
RANITIDINE HYDROCHLORIDE (UNII: BK76465IHM) (RANITIDINE - UNII:884KT10YB7)	RANITIDINE	75 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	pink	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	OR;606
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0352-30	1 in 1 CARTON		
1		30 in 1 BOTTLE		
2	NDC:0363-0352-08	1 in 1 CARTON		
2		80 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA201745	09/24/2013	

Labeler - Walgreen Company (008965063)

Registrant - Ranbaxy Pharmaceuticals Inc. (937890044)

Establishment

Name	Address	ID/FEI	Business Operations
Shasun Pharmaceuticals Limited		915786829	manufacture(0363-0352)

Revised: 10/2013

Walgreen Company