

## **IVERMECTIN- ivermectin paste**

### **Dechra Veterinary Products**

-----

#### **Ivermectin Paste**

Approved by FDA under ANADA # 200-390

1.87% Anthelmintic and Boticide  
Removes worms and bots with a single dose.

Each Syringe Contains 0.21 oz. (6.08 g) IVERMECTIN PASTE

Net Wt: 0.21 oz (6.08 g)  
*Contents will treat up to 1250 lbs body weight.*

#### **For Oral Use in Horses Only**

**INDICATIONS:** Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. Ivermectin Paste provides effective treatment and control of the following parasites in horses. **Large Strongyles** (adults) – *Strongylus vulgaris* (also early forms in blood vessels), *S. edentatus* (also tissue stages), *S. equinus*, *Triodontophorus* spp. including *T. brevicauda* and *T. serratus* and *Craterostomum acuticaudatum*; **Small Strongyles** (adults, including those resistant to some benzimidazole class compounds) – *Coronocylus* spp. including *C. coronatus*, *C. labiatus* and *C. labratus*, *Cyathostomum* spp. including *C. catinatum* and *C. pateratum*, *Cylicocylus* spp. including *C. insigne*, *C. leptostomum*, *C. nassatus* and *C. brevicapsulatus*, *Cylicodontophorus* spp., *Cylicostephanus* spp. including *C. calicatus*, *C. goldi*, *C. longibursatus* and *C. minutus*, and *Petrovinema poculatum*; **Small Strongyles** – Fourth-stage larvae; **Pinworms** (adults and fourth-stage larvae) – *Oxyuris equi*; **Ascarids** (adults and third- and fourth-stage larvae) – *Parascaris equorum*; **Hairworms** (adults) – *Trichostrongylus axei*; **Large-mouth Stomach Worms** (adults) – *Habronema muscae*; **Bots** (oral and gastric stages) – *Gasterophilus* spp. including *G. intestinalis* and *G. nasalis*; **Lungworms** (adults and fourth-stage larvae) – *Dictyocaulus arnfieldi*; **Intestinal Threadworms** (adults) – *Strongyloides westeri*; **Summer Sores** caused by *Habronema* and *Draschia* spp. cutaneous third-stage larvae; Dermatitis caused by neck threadworm microfilariae, *Onchocerca* sp.

#### **DOSAGE:**

**Ivermectin Paste 1.87%                      Net Wt: 0.21 oz (6.08 g)**

This syringe contains sufficient paste to treat one 1250 lb horse at the recommended dose rate of 91 mcg Ivermectin per lb (200 mcg/kg) body weight. Each weight marking on the syringe plunger delivers enough paste to treat 250 lb body weight.

#### **ADMINISTRATION:**

**(1) While holding plunger, turn the knurled ring on the plunger 1/4 turn to the left and slide it so the side nearest the barrel is at the prescribed weight marking.**

**(2) Lock the ring in place by making a 1/4 turn to the right.**

(3) Make sure that the horse's mouth contains no feed.

(4) Remove the cover from the tip of the syringe.

(5) Insert the syringe tip into the horse's mouth at the space between the teeth.

(6) Depress the plunger as far as it will go, depositing paste on the back of the tongue.

(7) Immediately raise the horse's head for a few seconds after dosing.

**PARASITE CONTROL PROGRAM:** All horses should be included in a regular parasite control program with particular attention being paid to mares, foals and yearlings. Foals should be treated initially at 6 to 8 weeks of age, and routine treatment repeated as appropriate. Consult your veterinarian for a control program to meet your specific needs. Ivermectin Paste effectively controls gastrointestinal nematodes and bots of horses. Regular treatment will reduce the chances of verminous arteritis caused by *Strongylus vulgaris*.

**PRODUCT ADVANTAGES: Broad-spectrum Control-** Ivermectin Paste 1.87% kills important internal parasites, including bots and the arterial stages of *S. vulgaris*, with a single dose. Ivermectin Paste is a potent antiparasitic agent that is neither a benzimidazole nor an organophosphate.

**ANIMAL SAFETY:** Ivermectin Paste 1.87% may be used in horses of all ages, including mares at any stage of pregnancy. Stallions may be treated without adversely affecting their fertility.

**WARNING: Do not use in horses intended for human consumption. Not for use in humans. Keep this and all drugs out of reach of children.**

Refrain from smoking and eating when handling. Wash hands after use. Avoid contact with eyes.

The Safety Data Sheet (SDS) contains more detailed occupational safety information.

**PRECAUTIONS:** Ivermectin Paste 1.87% has been formulated specifically for use in horses **only**.

This product should not be used in other animal species as severe adverse reactions, including fatalities in dogs, may result.

**Environmental Safety:** Ivermectin and excreted ivermectin residues may adversely affect aquatic organisms. Do not contaminate ground or surface water. Dispose of the syringe in an approved landfill or by incineration.

**INFORMATION FOR HORSE OWNERS:** Swelling and itching reactions after treatment with Ivermectin Paste have occurred in horses carrying heavy infections of neck threadworm microfilariae (*Onchocerca* sp.) . These reactions were most likely the result of microfilariae dying in large numbers. Symptomatic treatment may be advisable. Consult your veterinarian should any such reactions occur. Healing of summer sores involving extensive tissue changes may require other appropriate therapy in conjunction with treatment with Ivermectin Paste. Reinfection, and measures for its prevention, should also be considered. Consult your veterinarian if the condition does not improve.

**CONTACT INFORMATION:**

To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Dechra at (866) 933-2472. For additional information about adverse drug experience reporting for animal drugs, contact FDA at

1-888-FDA-VETS or online at <http://www.fda.gov/reportanimalae>

**STORAGE:**

Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F).

**Manufactured for:**

Dechra Veterinary Products, LLC  
Overland Park, KS 66211

Rev. 02/2024

NDC 17033-455-60

**Ivermectin  
Paste 1.87%**  
(ivermectin paste)

For Oral Use in  
Horses Only

Anthelmintic and  
Boticide

Approved by FDA under ANADA # 200-390

Net Wt.  
0.21 oz  
(6.08g)

 **Dechra**

Apple  
Flavor

For treatment of Large Strongyles, Small Strongyles, Pinworms, Roundworms (Ascarids), Hairworms, Neck Threadworms, Large-mouth Stomach Worms, Bots. See carton or attached labeling for complete indications and use directions. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

**WARNING:** Do not use in horses intended for human consumption. Not for use in humans. Keep this and all drugs out of reach of children. Refrain from smoking or eating when handling. Wash hands after use. Avoid contact with eyes.

**STORAGE INFORMATION:** Store at 68°F - 77°F (20°C - 25°C). Excursions between 59°F - 86°F (15°C - 30°C) are permitted.

**Manufactured for:**  
Dechra Veterinary Products,  
7015 College Boulevard, Suite 525,  
Overland Park, KS 66211 USA  
[www.dechra-us.com](http://www.dechra-us.com)

Lot No./Exp. Date:  
Rev. September 2023

**IVERMECTIN**

ivermectin paste

**Product Information**

<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:17033-455
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>IVERMECTIN</b> (UNII: 8883YP2R6D) (IVERMECTIN - UNII:8883YP2R6D)	IVERMECTIN	1.87 g in 100 g

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17033-455-60	6.08 g in 1 SYRINGE		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200390	05/04/2023	

**Labeler** - Dechra Veterinary Products (362142734)

Revised: 2/2024

Dechra Veterinary Products