

ERY- erythromycin swab
Padagis Israel Pharmaceuticals Ltd

Ery 2% Pads
(Erythromycin Pledgets USP, 2%)

Rx Only


For Dermatologic Use Only

Not for Ophthalmic Use

DESCRIPTION

Ery 2% Pads contain erythromycin, USP for topical dermatologic use. Erythromycin is a macrolide antibiotic produced from a strain of *Saccaropolyspora erythraea* (formerly *Streptomyces erythreus*). It is a base and readily forms salts with acids.

Chemically, erythromycin is C₃₇H₆₇NO₁₃. It has the following structural formula:



The chemical name for erythromycin is (3R*,4S*,5S*,6R*,7R*,9R*,11R*,12R*,13S*,14R*)-4-[(2,6-Dideoxy-3-C-methyl-3-O-methyl-α-L-ribo-hexopyranosyl)oxy]-14-ethyl-7,12,13-trihydroxy-3,5,7,9,11,13-hexamethyl-6-[[3,4,6-trideoxy-3-(dimethylamino)-β-D-xyllo-hexopyranosyl]oxy] oxacyclotetradecane-2,10-dione.

Erythromycin has the molecular weight of 733.94. It is a white or slightly yellow, crystalline powder, slightly soluble in water, soluble in alcohol, in chloroform, and in ether. It is odorless or practically odorless. It has a pH range between 8.0 and 10.5 in a methanol and water solution prepared by diluting 1 volume of a methanol solution, containing 40 mg per mL, with 19 volumes of water.

Each mL of expressible liquid contains 20 mg erythromycin in a base of dehydrated alcohol, propylene glycol and citric acid to adjust pH. Each pledget is filled to contain 0.8 mL of Erythromycin Topical Solution 2%.

CLINICAL PHARMACOLOGY

The exact mechanism by which erythromycin reduces lesions of acne vulgaris is not fully known; however, the effect appears to be due in part to the antibacterial activity of the drug.

MICROBIOLOGY

Erythromycin acts by inhibition of protein synthesis in susceptible organisms by reversibly binding to 50S ribosomal subunits, thereby inhibiting translocation of aminoacyl transfer-RNA and inhibiting polypeptide synthesis. Antagonism has been demonstrated *in vitro* between erythromycin, lincomycin, chloramphenicol, and clindamycin.

INDICATIONS AND USAGE

Ery 2% Pads are indicated for the topical treatment of acne vulgaris.

CONTRAINDICATIONS

Ery 2% Pads are contraindicated in those individuals who have shown hypersensitivity to any of its components.

WARNINGS

Pseudomembranous colitis has been reported with nearly all antibacterial agents, including erythromycin, and may range in severity from mild to life-threatening. Therefore, it is important to consider this diagnosis in patients who present with diarrhea subsequent to the administration of antibacterial agents.

Treatment with antibacterial agents alters the normal flora of the colon and may permit overgrowth of clostridia. Studies indicate that a toxin produced by *Clostridium difficile* is one primary cause of "antibiotic-associated colitis".

After the diagnosis of pseudomembranous colitis has been established, therapeutic measures should be initiated. Mild cases of pseudomembranous colitis usually respond to drug discontinuation alone. In moderate to severe cases, consideration should be given to management with fluids and electrolytes, protein supplementation and treatment with an antibacterial drug clinically effective against *C. difficile* colitis.

PRECAUTIONS

General -

For topical use only; not for ophthalmic use. Concomitant topical acne therapy should be used with caution because a possible cumulative irritancy effect may occur, especially with the use of peeling, desquamating, or abrasive agents.

The use of antibiotic agents may be associated with the overgrowth of antibiotic-resistant organisms. If this occurs, discontinue use and take appropriate measures.

Avoid contact with eyes and all mucous membranes.

Information for Patients -

Patients using Ery 2% Pads should receive the following information and instructions:

1. This medication is to be used as directed by the physician. It is for external use only. Avoid contact with the eyes, nose, mouth, and all mucous membranes.
2. This medication should not be used for any disorder other than that for which it was prescribed.
3. Patients should not use any other topical acne medication unless otherwise directed by their physician.
4. Patients should report to their physician any signs of local adverse reactions.

Carcinogenesis, Mutagenesis, Impairment of Fertility -

No animal studies have been performed to evaluate the carcinogenic and mutagenic potential or effects on fertility of topical erythromycin. However, long-term (2-year) oral studies in rats with erythromycin ethylsuccinate and erythromycin base did not provide evidence of tumorigenicity. There was no apparent effect on male or female fertility in rats fed erythromycin (base) at levels up to 0.25% of diet.

Pregnancy:

Teratogenic Effects:

Pregnancy Category B -

There was no evidence of teratogenicity or any other adverse effect on reproduction in female rats fed erythromycin base (up to 0.25% diet) prior to and during mating, during gestation and through weaning of two successive litters.

There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used in pregnancy only if clearly needed. Erythromycin has been reported to cross the placental barrier in humans, but fetal plasma levels are generally low.

Nursing Women -

It is not known whether erythromycin is excreted in human milk after topical application. However, erythromycin is excreted in human milk following oral and parenteral erythromycin administration. Therefore, caution should be exercised when erythromycin is administered to a nursing woman.

Pediatric Use -

Safety and effectiveness of this product in pediatric patients have not been established.

ADVERSE REACTIONS

The following local adverse reactions have been reported occasionally: peeling, dryness,

itching, erythema, and oiliness. Irritation of the eyes and tenderness of the skin have also been reported with topical use of erythromycin. Generalized urticarial reactions, possibly related to the use of erythromycin, which required systemic steroid therapy have been reported.

DOSAGE AND ADMINISTRATION

The Ery 2% Pads should be rubbed over the affected area twice a day (morning and evening) after skin is thoroughly washed with warm water and soap and patted dry. Acne lesions on the face, neck, shoulders, chest, and back may be treated in this manner. Additional pledgets may be used, if needed. Each pledget should be used once and discarded. Wash hands after application. Close jar tightly after each use. Drying and peeling may be controlled by reducing the frequency of applications.

HOW SUPPLIED

Ery 2% Pads are available as follows:

A plastic jar containing 60 pledgets (NDC 45802-**962**-72)

Each pledget is filled to contain 0.8 mL of Erythromycin Topical Solution 2%.

STORAGE

Keep jar tightly closed.

Store at 20-25°C (68-77°F) [see USP Controlled Room Temperature].

Made in Israel

Manufactured by Perrigo

Yeruham 80500, Israel

Distributed By

 **PERRIGO**®

Allegan, MI 49010

www.perrigo.com

Rev. 08/11

6E500 RC J4

Principal Display Panel

Rx Only

Ery 2% Pads (Erythromycin Pledgets USP, 2%)

For External Use

Avoid Contact with Eyes

60 Pledgets

Usual Dosage: See package insert for complete product information. Keep jar tightly closed. Keep out of the reach of children. Store at 20-25°C (68-77°F) [see USP Controlled Room Temperature].

Instructions for use:

1. Wash affected area with warm water and soap and then pat dry.
2. Use pledget to apply to affected areas twice a day. Keep pledget away from eyes, nose, mouth and other mucous membranes. If medication accidentally enters eyes, rinse thoroughly with tap water. Additional pledgets may be used, if needed. Each pledget should be used once and discarded.
3. Close jar tightly after each use.
4. Wash hands after use.

Rx Only

NDC 45802-962-72

Ery 2% Pads
(Erythromycin Pledgets USP, 2%)

For External Use
Avoid Contact with Eyes

60 Pledgets

Perrigo®

Each mL of Erythromycin Topical Solution 2% contains 20 mg erythromycin in a base of dehydrated alcohol, propylene glycol and citric acid to adjust pH. Each pledget is filled to contain 0.8 mL of Erythromycin Topical Solution 2%. If protective seal is broken or missing return to place of purchase.

Made in Israel
Manufactured By Perrigo, Yeruham, Israel
Distributed By Perrigo Allegan, MI 49010
www.perrigo.com
Rev 09-16
: 6E572 RC F6

Lot # / Exp date area

The following image is a placeholder representing the product identifier that is either affixed or imprinted on the drug package label during the packaging operation.

S/N [insert product's serial number]
 Lot [insert product's lot number]
 Exp [insert product's expiration date]

ERY			
erythromycin swab			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:45802-962
Route of Administration	TOPICAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ERYTHROMYCIN (UNII: 63937KV33D) (ERYTHROMYCIN - UNII:63937KV33D)		ERYTHROMYCIN	20 mg in 1 mL
Inactive Ingredients			
Ingredient Name			Strength
ALCOHOL (UNII: 3K9958V90M)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:45802-962-72	60 in 1 JAR	07/07/2008	
1		0.8 mL in 1 APPLICATOR; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA064126	07/07/2008	

Labeler - Padagis Israel Pharmaceuticals Ltd (600093611)

Revised: 12/2021

Padagis Israel Pharmaceuticals Ltd